

**FINAL REPORT OF THE THIRTY SEVENTH
NORTH CAROLINA LAYER PERFORMANCE
AND MANAGEMENT TEST¹**

Vol. 37, No. 5
October 2009

The North Carolina Layer Performance and Management Test is conducted under the auspices of the Cooperative Extension Service at North Carolina State University and the North Carolina Department of Agriculture and Consumer Services. The flock is maintained at the Piedmont Research Station, Salisbury, North Carolina. Mr. Joe Hampton is the Piedmont Research Station Superintendent; Mr. Kelly Snider is Poultry Unit Manager; Pam Jenkins is the Statistical Research Assistant; and Dr. K. E. Anderson is Project Leader. The purpose of this program is to assist poultry industry personnel in North Carolina, across the country, and internationally in the evaluation of commercial layer stocks and management systems.

The data presented herein provides the analyses of the first production cycle, the molt, the second production cycle, and of the combined 1st and 2nd cycle performance for the 37th North Carolina Layer Performance and Management Test. Performance summary tables are available for each strain, and molt treatment used as well as for the combined results.

For further information contact:

Dr. Kenneth E. Anderson
Poultry Science Department
North Carolina State University
Box 7608
Raleigh, NC 27695-7608
Tel: (919) 515-5527
Fax: (919) 515-7070
Email: ken_anderson@ncsu.edu

¹The use of trade names in this publication does not imply endorsement by the North Carolina Cooperative Extension Service of the products named nor criticism of similar ones not mentioned.

**37th NORTH CAROLINA LAYER PERFORMANCE AND
MANAGEMENT TEST
Volume 37 No. 5**

Final Report on First Laying Cycle, Molt, Second Cycle and Combined Results

Entries and Strains:

A total of ten white egg and six brown egg strains were entered for a total of sixteen strains that were accepted in accordance with the rules and regulations of the test. The strain names and egg color designations are shown in Table 1.

Table 1. Strain name and egg color designation

Strain	Egg Color Designation
Hy-Line W-36	White
Hy-Line W-98	White
Hy-Line CV-22	White
Shaver White	White
Dekalb TX	White
Lohmann LSL-Lite	White
H&N Nick Chick	White
Bovans White	White
Hisex White	White
Bovans Robust	White
ISA Brown	Brown
Hy-Line Brown	Brown
Hy-Line S. Brown	Brown
Bovans Brown	Brown
Hisex Brown	Brown
Dekalb Amber Link	Brown

In the layer test, a minimum of 760 white and brown egg pullets/strain were placed at the initiation of the layer portion of the test. However, if the number of pullets reared for a given strain was below the prescribed numbers, the pullets that were available were divided as equally as possible among the levels and replicates within the layer house, and the numbers actually placed into the layer test were recorded appropriately.

Dates of Importance:

The eggs were placed into trays and set on May 15, 2007 at the North Carolina Dept. of Agriculture and Consumer Services, Piedmont Research Station's Poultry Unit at Salisbury, NC. The flock was hatched on June 6, 2007 and were reared in grow cages at the Unit. They were then moved to the laying facilities on September 26-28, 2007 during their 17th week of age.

First cycle production records commenced on October 3, 2007 (17 weeks of age), through the molt period which was induced on September 10, 2008 [Tables 14-19]. The molt records commenced on September 10, 2008 (66 weeks of age), and ended on October 8, 2008 (70 weeks of age) [Tables 20-27]. The Second cycle commenced on October 8, 2008 and ended on June 15, 2009 (105 weeks of age) [Tables 28-39]. This report includes production data summarized from 17 to 66 weeks (1st cycle), 66 to 70 weeks (molt), and 70 to 105 weeks (2nd cycle) for the production of eggs in cages by molt program. Tables referring to the Non-Molted hens [Tables 40-45], Non-Anorexic Molt 20% body weight loss hens [Tables 46-51], and Non-Anorexic Molt 25% body weight loss hens [Tables 52-57] along with the overall data tabulations are located in Tables 58-69. A table showing changes in body weights from 17 to 66 wk of age and the weight loss during the molt period and ending body weights are included in Tables 70-73.

Pullet Housing:

The chicks were randomly assigned to the growing cages with white egg and brown egg replicates being intermingled throughout the house. The white egg strains occupied approximately 59 % of the house and brown egg strains occupied the other 42 % of the house. All strains were assigned to be represented as equally as possible in each room, row, and cage level.

The chicks from the brown egg strains destined for a range study were randomly assigned to the growing pens throughout House 6. The results of the range study were reported in the 37th NCLP&MT Single Cycle Report, (Vol. 37, No. 4).

House 8--is an environmentally controlled closed brood-grow facility with 3 banks of quad-deck cages in each room. Each room was assigned a number, each side of each bank was assigned a row number, each cage section within each row and level/row has been assigned a replicate number, and for statistical analysis pairs of rows were designated as blocks. Thus, each block consisted of two rows containing 24 replicates on all levels. This allows for a total of 3,744 pullets per room resulting in a total pullet count for this test in House 8 using 3 rearing rooms of 11,232. The white and brown-egg strains were randomly assigned to the replicates in the house. Entrant strains were assigned to the replicates in a restricted randomized manner with the restrictions being that all strains were approximately equally represented in all rows, levels, and rooms. The chicks were brooded in the same cage during the entire 17 wk rearing period. Paper was placed on the cage floor for the first 7 days within each of the replicate series within each row. Each cage within the replicate was filled with 13 white-egg or brown-egg (i.e. 13 chicks per 24" x 26" cage) pullets on the day of hatch for a rearing allowance of 48 in², 4.7 cm (1.8 in) of feeder space/bird and 1:6.5 nipple drinkers to bird ratio. The same numbers of pullets were grown in each replicate for both white and brown-egg strains. The room dividers were removed for this test so that all birds were essentially reared in a contiguous house.

Layer Housing:

The hens were randomly assigned to the replicate cages with white egg and brown egg strains being intermingled throughout the houses. The white egg strains occupied 60% of the house and brown egg strains occupied the other 40%. All strains were assigned to be represented as equally as possible in all rows, and levels.

House 4 is a high rise, environmentally controlled facility with three banks of Quad-deck (4-tier) high cages. There are a total of 216 replicates in house 4 which can support 5,184 hens. The replicate blocks contain cages that are either 61 or 81 cm wide.

House 5 is a standard height totally enclosed force ventilated laying house with a scraper pit manure handling system. It has 2 banks of tri-deck cages and two banks with quad-deck (4 levels) cages. There are a total of 252 replicates in house 5 which can support 6,048 hens.

In both houses, each side of a bank was designated as a row and each row was divided into 9 8-foot replicates/level. The replicates are equipped with feed hoppers to supply and monitor feed consumption for each individual replicate and the feed is distributed by an automatic feeding system. The cage density in both was dictated by the cage size. That is, each replicate contained cages that were either 61 or 81.2 cm wide and 40.6 cm deep, which allowed for a constant density of 64 in² (413 cm²), at 6 or 8 hens/cage, respectively. The white-egg and brown-egg strains were assigned to the replicates in a restricted randomized manner, with the restrictions being that all strains were approximately equally represented in all rows, levels and cage sizes.

Test Design:

The arrangement for the laying test involved a completely randomized design and the main effects were set up in a factorial arrangement. The main effects within Houses 4 and 5 were strain, population, and molt treatment. Following are general descriptions of the main effects:

Strain

The samples of fertile eggs were provided directly by the breeders involved [Table 74]. All eggs were set and hatched concurrently. A total of ten white egg strains and six brown egg strains participated in the test. See the 37th Hatch Report (Vol. 37, No. 1) for details.

Density

In Houses 4 and 5, all individual cages within each block contained either the brown or the white egg layers. Thus the replicate consisted of 24 hens per replicate, the hens were contained in 61 x 40.6 cm cages for 4 cages with 6 hens/cage or 81.2 x 40.6 cm cages for 3 cages with 8 hens/cage. See Table 2 for density, feed and water space allocations.

Table 2. Population and Density Allocations in Houses 4 and 5

White and Brown Hens per Cage	Cage Size Width Depth	Floor Space per Bird	Feeder Space per Bird	Water Nipples per Cage
3	61 cm x 40.7 cm	413 cm ² (64 in ²)	10.2 cm 4.0 in	1
4	81.2 cm x 40.7 cm	413 cm ² (64 in ²)	10.2 cm 4.0 in	1

Table 3. Laying House

Age	Date	House 4 (Light Hours)	House 5 (Light Hours)
Housing Pullets	Sept. 26-Oct 3, 2007	10.0	10.0
17 Weeks ¹	Oct. 3, 2007	11.0	11.0
18 Weeks	Oct. 10, 2007	11.5	11.5
19 Weeks	Oct. 17, 2007	12.0	12.0
20 Weeks	Oct. 24, 2007	12.5	12.5
21 Weeks	Oct. 30, 2007	13.0	13.0
22 Weeks	Nov. 7, 2007	13.5	13.5
23 Weeks	Nov. 14, 2007	14.0	14.0
24 Weeks	Nov. 21, 2007	14.25	14.25
25 Weeks	Nov 28, 2007	14.5	14.5
26 Weeks	Dec. 5, 2007	14.75	14.75
27 Weeks	Dec. 12, 2007	15.0	15.0
28 Weeks	Dec. 19, 2007	15.25	15.25
29 Weeks	Dec. 26, 2007	15.5	15.5
30 Weeks	Jan. 2, 2008	15.75	15.75
31 Weeks	Jan. 9, 2008	16.0	16.0
Through 66 Weeks	Sept 11, 2008	16.0	16.0

Layer Management (Molting):

The molt experiment was conducted utilizing all hens involved in the layer test. Participating strains were randomly divided into three groups such that all strains, populations, and levels were approximately equally represented. In this test, each group received one of the following treatments during the molt period commencing at 66 wks of age.

Table 4. Molt Program Names and Treatment Codes

Program Name	Brief Description	Treatment Code
Full Fed Control	Not Molted	NM
Non-anorexic molt program with 20% wt loss	LP/LE Diet no fasting	NA20
Non-anorexic molt program with 25% wt loss	LP/LE Diet no fasting	NA25

Full Fed Control (NM): The replicates assigned to the full fed control group were maintained according to the standard management program as outlined previously in rows 1 and 2. The laying house was partitioned the length of the house with an opaque curtain such that the lighting program was consistent for maximum egg production.

Non-anorexic molt program (NA): The hens were fed a diet, which contained low protein, low energy, and had supplemental Ca for skeletal maintenance. When birds in the replicates were being weighed and they reached target weight, that replicate and their sister replicates were returned to full feed. The induced molt was started at 66 wks of age. This Non-anorexic molt diet was low in energy and was designed to keep hens out of production and provide balanced nutrition for body maintenance only. The diet had low bulk density, such that a full trailer load will only weigh 2/3 of a normal full load.

Procedural steps:

Day -7 Sample of birds were weighed to determine the pre-molt weight. Target weight loss (20 or 25 % body weight) was calculated using the pre-molt weight.

Day 0 Remove morbid birds before initiation of the molt program. The NA program was initiated with the remaining layer feed being removed and replaced with the NA molt diet combined with a daylight hour reduction. Controlled light housing, reduce the day length to 8 hr [See Table 5].

Day +7 Sample of birds were weighed 7 days after diet change to determine body weights.

Day +9 Sample of birds were weighed 9 days after diet change to determine body weight. Weight loss per day calculated using 7 and 9 day body weights and target date for the % weight loss determined. When the target date for the % body weight loss is determined the hens were not be weighed until target date at which time they were provided as resting feed if body weight loss had been achieved.

Day +28 Birds were fed layer diet and light stimulated.

Table 5. Molting Lighting Schedules

Age	Date	House 4 (Light Hours)	5 (Light Hours)
Through 66 Weeks	Sept. 11, 2008	16.0	16.0
66 Weeks	Sept. 11, 2008	8 hr	8 hr
69 Weeks	Oct. 2, 2008	12.0	12.0
70 weeks	Oct. 9, 2008	13.0	13.0
71 Weeks	Oct. 16, 2008	14.0	14.0
72 Weeks	Oct. 23, 2008	15.0	15.0
73 Weeks through end of test (105 wk)	Oct. 30, 2008 to June 15, 2009	16.0	16.0

Specific monitored criteria for all of the molt programs include the following.

The goal was for the birds to attain approximately 20 or 25% body weight loss.

Maintain house temperature at 80± 5° F, but the birds should not pant. Please react to environmental temperatures.

Actual house conditions and the flock's reaction to the NCSU Non-Fasting Molting Program may affect how the light stimulation will actually be given.

Hens should have ceased egg production by Day 6-10 of the molt program. However, the hens should be allowed to consume all of the feed provided between feedings. The molting ration is designed to keep hens out of production, and to provide for skeletal and muscle maintenance. Livability was excellent with this program. The diet is bulky, such that a full trailer load will only weigh 2/3 of a normal full load. Please keep this fact in mind when ordering feed. Diet E will bring hens back into peak production. Feed intake and egg size will determine which diet to progress toward.

Layer Nutrition:

Layer diets are identified as Diets D, E, F, G, H, I, M, N, and O which consist of a pre-lay diet and a series of layer diets formulated to assure a daily protein, mineral and amino acid intake as shown below. Feed was offered *ad libitum* in accordance with the guidelines that all birds should receive acceptable nutrient intake at all times depending on the bird's age and production rate as shown in the Laying House Feeding Program Table.

The diets provided during the molt, consisted of a low protein/energy diet and a Resting Diet described in the Molt Diets Table which follow. The molt diets were formulated to provide the layer with the nutrients needed to maintain a static body weight with no egg production.

Table 6. Minimum Daily Intake of Nutrients Per Bird at Various Stages of Production in the 37th NCLP&MT

Production Stage	Pre-Peak > 87%	87-80%	80-70%	<70%
White Egg Layers				
Protein ¹ (g/day)	19	18	17	16
Calcium (g/day)	4.0	4.1	4.2	4.3
Lysine (mg/day)	820	780	730	690
TSAA (mg/day)	700	670	630	590
Brown Egg Layers				
Protein ¹ (g/day)	20	19	18	17
Calcium (g/day)	4.0	4.0	4.1	4.2
Lysine (mg/day)	830	820	780	730
TSAA (mg/day)	710	700	670	630

¹ If the egg production is higher than predicted values protein intake should be increased by 1%
 Note: House temperatures dictate the body maintenance demand of the hen. If the house temperature is 75 to 80°F feed protein content should be increased accordingly to compensate for metabolic heat needed to maintain a homeostatic body temperature. If the house temperature is at or above 85°F no adjustment is needed.

Table 7. 37th NCLP&MT Laying House Feeding Program

Rate of Production	Consumption Per 100 Birds/Day (kg)	Diet Fed	
		White Egg Strains	Brown Egg Strains
Weeks 17-26	< 9.52	D	D
Pre-Peak and > 87%	< 9.52	D	D
	9.57-10.39	F	E
	10.43-11.29	H	G
	11.34-12.20	I	H
	12.25-13.11	M	I
	>13.15	N	M
80-87%	< 9.52	F	E
	9.57-10.39	G	F
	10.43-11.29	I	H
	11.34-12.20	M	I
	12.25-13.11	N	M
	>13.15	O	N
70-80%	< 9.52	H	G
	9.57-10.39	I	H
	10.43-11.29	M	I
	11.34-12.20	N	M
	12.25-13.11	O	N
	>13.15	O	O
< 70%	< 9.52	H	G
	9.57-10.39	I	H
	10.43-11.29	N	M
	11.34-12.20	O	N
	12.25-13.11	O	O
	>13.15	O	O

Note: Low house temperatures and egg production higher than breeder guides for any given hen age will require an adjustment to the dietary phase feeding program to ensure the hens are in a positive nutrient status.

Table 8. 37th NCLP&MT Laying Periods Feed Formulations D through H

Ingredients	D	E	F	G	H
Corn	866.71	925.46	997.91	1068.19	1131.97
Soybean meal	663.18	621.10	552.33	499.80	457.65
Wheat Midds					
Fat (Tallow)	110.88	102.43	87.73	74.61	64.32
Gluten Meal 60%	95.83	88.37	100.00	99.23	90.80
D.L. Methionine	3.08	2.89	2.52	2.26	2.48
Lysine 78.8%					
Soybean Hulls					
Ground Limestone	132.42	133.70	135.07	134.02	132.50
Coarse Limestone	75.00	75.00	75.00	75.00	75.00
Bi-Carbonate	3.00	3.00	3.00	3.00	3.00
Phosphate Mono/D	36.77	34.73	32.84	30.36	28.79
Salt	6.00	5.99	5.95	5.93	5.92
Vit. premix	1.00	1.00	1.00	1.00	1.00
Min. premix	1.00	1.00	1.00	1.00	1.00
Mold Inhibitor	1.00	1.00	1.00	1.00	1.00
T-Premix	1.00	1.00	1.00	1.00	1.00
.06% Selenium Premix	1.00	1.00	1.00	1.00	1.00
Choline Cl 60%	2.14	2.33	2.65	2.59	2.57
Calculated Analysis					
Protein %	22.0	21.0	20.00	19.00	18.0
ME kcal/kg	2926.0	2926.0	2926.0	2926.0	2926.0
Calcium %	4.45	4.45	4.45	4.40	4.35
T. Phos. %	0.71	0.68	0.65	0.61	0.59
Lysine %	1.15	1.09	1.00	0.93	0.87
TSAA %	0.89	0.85	0.81	0.77	0.75

Table 9. 37th NCLP&MT Laying Periods Feed Formulations I through O

Ingredients	I	M	N	O
Corn	1199.47	1258.28	1309.81	1371.93
Soybean meal	406.08	363.91	340.24	333.87
Wheat Midds				
Fat (Tallow)	52.26	43.80	38.85	14.71
Gluten Meal 60%	89.84	82.64	61.54	25.79
D.L. Methionine	2.02	1.62	1.75	1.80
Lysine 78.8%				
Soybean Hulls				
Ground Limestone	158.82	160.10	161.33	167.71
Coarse Limestone	50.00	50.00	50.00	50.00
Bi-Carbonate	3.00	3.00	3.00	3.00
Phosphate Mono/D	26.79	24.75	22.60	20.30
Salt	5.90	5.89	5.89	5.89
Vit. premix	1.00	1.00	1.00	1.00
Min. premix	1.00	1.00	1.00	1.00
Mold Inhibitor	1.00	1.00	1.00	1.00
T-Premix	1.00	1.00	1.00	1.00
.06% Selenium Premix	1.00	1.00	1.00	1.00
Choline Cl 60%	0.83	1.02		
Calculated Analysis				
Protein %	17.00	16.00	15.00	14.00
ME kcal/kg	2926.0	2926.0	2926.0	2860.0
Calcium %	4.35	4.35	4.35	4.45
T. Phos. %	0.56	0.52	0.49	0.47
Lysine %	0.80	0.74	0.70	0.68
TSAA %	0.70	0.65	0.62	0.58

Table 10. 37th NCLP&MT Laying Periods Feed Formulations Molt and Resting Diets

Ingredient	Molt Diets	
	<u>Low ME</u>	<u>Resting</u>
Corn	702.50	1427.70
Corn Gluten Meal		
Soybean Hulls	1164.77	226.00
Soybean Meal 48%		117.00
Wheat Midds	18.26	186.50
Coarse Limestone	17.78	16.50
Phosphate Mono/D	69.84	4.00
Bentonite		
Salt	9.16	5.00
Methionine	2.69	1.30
Choline Chloride		
Vit. premix	1.00	1.00
Min. premix	1.00	1.00
T - Premix	1.00	1.00
Fat	9.99	10.00
MYC-OUT 65	1.00	2.00
.06% Sel Premix	1.00	1.00
Total	2000	2000
Calculated Analysis		
Protein %	9.92	11.75
Me kcal/kg	1650	2859
Calcium %	1.33	3.80
T. Phos %	0.88	0.44
Lysine %	0.42	0.55
TSAA %	0.35	0.49

Data Collection Schedule and Procedures:

Age at 50% Production (Maturity)—The first day at which the birds in the individual replicates achieved 50% production.

Egg Production--All eggs that had the potential of being marketed were credited toward the test unit's (replicate) egg production, regardless of the shell condition at the time of collection. All eggs were collected and recorded daily. Egg production was summarized at twenty-eight day intervals, and was calculated and reported on a Hen-Housed and Hen-Day basis.

Egg Weight--At twenty-eight day intervals, all eggs produced in the previous 24-hour period were weighed and sorted by size (See egg size distribution). Percentages of eggs within each size category, average egg weight (g), and egg mass (g) were calculated and reported.

Egg Quality--At twenty-eight day intervals, all eggs produced within the previous 24 hours were examined by candling light and graded according to current USDA standards for egg quality. Eggs were graded in the pilot processing facility and handled as they would be in a commercial off-line facility.

Egg Size Distribution--At twenty-eight day intervals, all eggs produced within the previous 24 hours were weighed and sorted according to current USDA standards for egg size.

Egg Income--Egg income was calculated using current year regional average prices for farm value of eggs based on egg production and quality evaluation.

Feed Consumption and Conversion--All feed offered for consumption was recorded for each replicate. At twenty-eight day intervals, feed not consumed was weighed back and feed consumption was calculated. Daily feed intake (kg/100 hens/day) was calculated and reported for each strain.

Feed Costs--Feed costs were based on the actual current feed prices for each feed delivery which were calculated and summarized for the complete production cycle.

Body weights—Birds were weighed and weights recorded at housing (17 wk), end of 1st cycle (66 wks), and at the start of the 2nd cycle (70 wk). Body weight gain for the 1st cycle was calculated and reported for each strain. In the Molt period lowest body weight, percent weight loss, 70 wk body weights were taken or calculated and reported for each strain.

Mortality--All mortalities were recorded daily, and obvious accidents were not included in reported mortalities.

Statistical Analyses and Separation of Means:

All data were subjected to ANOVA utilizing the GLM procedure of SAS, with main effects of strain and density. Separate analyses were conducted for white and brown egg strains. Significant differences ($P < 0.01$) within white and brown egg strains are noted by differing letters among columns of means. The layer houses data for houses 4 and 5 were pooled in this analysis. First and second order interactions were tested for significance. LS Means which is part of the GLM Procedure were separated via the PDIF option.

DESCRIPTION OF DATA TABLE STATISTICS

First cycle performance of white and brown egg strains are shown on Tables 14 to 19. The molt period performance and weight loss data of the white and brown egg strains are shown on Tables 20 to 27. There is also a report on the productivity of range hens and caged hens maintained for a single production cycle from 17-82 weeks of age (NCLP&MT Report Vol. 37, No. 4)

Breeder (Strain):

Short identification codes of the breeder and strain of the stock were developed. See more complete information following data tables in Table 74.

Hen Housed Eggs per Bird:

The total number of eggs produced divided by the number of birds housed at 119 days.

Hen Day Egg Production:

The average daily number of eggs produced per 100 hens per day.

Egg Mass:

The average daily egg mass production in grams per hen day.

Mortality:

The percentage of birds which died between 119 through 462, 462 through 490, and 490 through 735 days of age which occurred during the First-Cycle, Molt, and Second-Cycle periods, respectively are reported separately.

Feed Consumption:

The kilograms of feed consumed daily per 100 hens (housed or hen days).

Feed Conversion:

The grams of egg produced per gram of feed consumed.

Egg Weight:

The average egg weight (gms) for each period sampled. Weight of all eggs collected from previous 24 hours divided by the number of eggs collected.

Egg Income:

The calculated income per hen housed at 119 days, from egg production using current year regional average egg prices 10/3/2007 to 10/8/2008 and 10/09/2008 to 6/15/2009.

Table 11. Current year regional average egg prices 10/3/2007 to 6/15/2009.

Grade	Size	\$\$/Dozen 1 st Cycle	\$\$/Dozen 2 nd Cycle
A	Extra Large	1.45	1.24
A	Large	1.41	1.22
A	Medium	1.21	0.99
A	Small	0.96	0.81
A	Pee Wee	0.48	0.40
B	All	0.75	0.65
Checks	All	0.75	0.65

Grade Information:

The average grade of all eggs sampled according to USDA grading standards over all sampling periods. Grades are established by personnel trained in USDA grading standards.

Egg Size Distribution:

Following are the size classifications used for establishing the USDA egg size grading. There has been blending of egg size in this test with the weight cutoff between medium and large being 23.5. This maximizes the number of USDA large eggs

just as would occur in a commercial plant. The proportion of the eggs falling into the following size categories are reported in the tables.

Table 12. USDA Egg Weights Used To Establish The Egg Size Distribution Weighted for Large Eggs.

Size Category	Ounces/Dozen
Pee Wee	< 18
Small	18 – 21
Medium	21 - 23.5
Large	23.5 – 27
Extra Large	> 27

Feed Cost:

The calculated feed cost per hen housed at 119 days, using the pounds/diet consumed and the average price of each diet per ton.

Table 13. The Average Contract Feed Price For Feed Purchases During The First Cycle.

<u>Diets</u>	<u>Price Per Ton 1st Cycle</u>	<u>Price Per Ton 2nd Cycle</u>
D	300.30	
E	339.27	332.13
F	366.90	325.28
G	374.27	327.45
H	360.61	305.96
I	382.53	293.86
M		283.50
N		274.80
O		259.57
Molt Diet LP/LE	328.60	
Resting	299.20	

Metric Conversions:

1 lb = 453.6 g	1 g = .03527 oz
1 lb = .4536 kg	1 kg = 2.204 lb
1 oz = 28.35 g	1 g = 1000 mg
	1 kg = 1000 g

TABLE 14. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (119-462 DAYS)

Breeder	Population ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
Hy-Line	6	9.5	0.52	276.5	82.5	49.4	2.3	140
W-36	8	9.5	0.52	272.5	82.7	49.9	2.9	140
	Average	9.5 ^C	0.52 ^A	274.5 ^B	82.6 ^D	49.6 ^D	2.6 ^C	140 ^{AB}
Hy-Line	6	10.1	0.51	276.7	82.7	51.6	3.7	137
W-98	8	10.3	0.50	274.2	82.3	51.4	3.8	136
	Average	10.2 ^B	0.51 ^{AB}	275.4 ^B	82.5 ^D	51.5 ^C	3.7 ^C	136 ^C
Hy-Line	6	10.3	0.50	271.8	83.4	52.1	7.3	137
CV-22	8	10.1	0.51	276.2	83.8	52.3	8.5	136
	Average	10.2 ^B	0.51 ^{AB}	274.0 ^B	83.6 ^{CD}	52.2 ^{ABC}	7.9 ^{AB}	136 ^C
Shaver	6	9.7	0.51	277.2	84.5	49.8	6.6	141
White	8	9.6	0.52	271.6	85.2	50.4	9.1	140
	Average	9.6 ^C	0.52 ^A	274.4 ^B	84.8 ^{BC}	50.1 ^D	7.8 ^{AB}	141 ^A
DeKalb	6	10.3	0.50	282.4	85.6	52.0	3.4	140
TX	8	10.0	0.50	276.9	84.5	50.8	6.6	139
	Average	10.2 ^B	0.50 ^B	279.6 ^{AB}	85.1 ^B	51.4 ^C	5.0 ^{BC}	140 ^{AB}
Lohmann	6	10.6	0.50	283.0	87.3	52.9	7.9	140
LSL-Lite	8	10.8	0.49	287.9	87.8	53.3	7.4	140
	Average	10.7 ^A	0.49 ^B	285.5 ^A	87.5 ^A	53.1 ^A	7.6 ^{AB}	140 ^{AB}
H&N	6	10.1	0.51	286.1	87.4	51.4	7.0	139
Nick Chick	8	10.3	0.51	282.8	88.4	52.4	9.3	140
	Average	10.2 ^B	0.51 ^{AB}	284.5 ^A	87.9 ^A	51.9 ^{BC}	8.1 ^{AB}	139 ^B
Bovans	6	10.4	0.51	284.6	87.6	53.1	8.9	139
White	8	10.3	0.51	285.2	88.1	53.0	9.5	140
	Average	10.3 ^{AB}	0.51 ^{AB}	284.9 ^A	87.9 ^A	53.1 ^A	9.2 ^A	139 ^B
Hisex	6	10.4	0.50	286.1	87.2	52.9	6.1	140
White	8	10.4	0.50	283.2	86.4	52.6	8.5	139
	Average	10.4 ^{AB}	0.50 ^B	284.6 ^A	86.8 ^A	52.8 ^{AB}	7.3 ^{AB}	140 ^{AB}
Bovans	6	10.0	0.51	276.7	83.7	51.8	4.9	140
Robust	8	10.1	0.51	277.5	84.4	52.2	4.6	140
	Average	10.1 ^B	0.51 ^{AB}	277.1 ^B	84.0 ^{BC}	52.0 ^{ABC}	4.7 ^{BC}	140 ^{AB}
All	6	10.1	0.51	280.1	85.2	51.7	5.8	139
Strains	8	10.1	0.51	278.8	85.4	51.8	7.0	139

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 15. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (119-462 DAYS)

Breeder	Population ¹	Egg Weight	Peewee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
Hy-Line	6	59.3	0.0	1.5	14.9	27.8	55.7
W-36	8	59.7	0.0	1.4	13.3	25.7	59.6
	Average	59.5 ^C	0.0	1.5 ^{ABCD}	14.1 ^{AB}	26.7 ^{BC}	57.6 ^B
Hy-Line	6	62.0	0.0	0.5	9.5	18.6	71.4
W-98	8	62.2	0.0	0.5	10.6	17.2	71.6
	Average	62.1 ^{AB}	0.0	0.5 ^{EF}	10.0 ^{DE}	17.9 ^D	71.5 ^A
Hy-Line	6	62.2	0.0	0.3	8.8	18.8	71.9
CV-22	8	62.2	0.0	0.3	9.0	19.1	71.4
	Average	62.2 ^A	0.0	0.3 ^F	8.9 ^E	19.0 ^D	71.7 ^A
Shaver	6	58.1	0.0	2.8	14.7	36.1	46.4
White	8	58.6	0.0	1.6	14.6	36.0	47.9
	Average	58.4 ^D	0.0	2.2 ^A	14.6 ^{AB}	36.0 ^A	47.1 ^C
DeKalb	6	60.1	0.0	1.3	13.1	25.1	60.5
TX	8	59.5	0.0	1.5	12.7	27.6	58.3
	Average	59.8 ^C	0.0	1.4 ^{BCD}	12.9 ^{ABC}	26.3 ^{BC}	59.4 ^B
Lohmann	6	59.9	0.0	1.4	11.5	28.3	58.8
LSL-Lite	8	60.0	0.0	1.4	11.9	25.4	61.3
	Average	60.0 ^C	0.0	1.4 ^{BCD}	11.7 ^{CD}	26.9 ^{BC}	60.1 ^B
H&N	6	58.2	0.1	1.6	16.0	35.3	47.0
Nick Chick	8	58.7	0.0	1.8	13.9	32.1	52.1
	Average	58.5 ^D	0.0	1.7 ^{ABC}	14.9 ^A	33.7 ^A	49.6 ^C
Bovans	6	60.1	0.0	0.9	11.7	28.4	59.0
White	8	59.5	0.0	1.3	13.1	28.9	56.7
	Average	59.8 ^C	0.0	1.1 ^{CDE}	12.4 ^{BCD}	28.6 ^B	57.8 ^B
Hisex	6	60.0	0.0	2.2	11.4	24.7	61.8
White	8	60.2	0.0	2.0	11.6	25.5	60.8
	Average	60.1 ^C	0.0	2.1 ^{AB}	11.5 ^{CD}	25.1 ^C	61.3 ^B
Bovans	6	61.3	0.0	1.1	11.2	20.6	67.2
Robust	8	61.3	0.0	0.7	10.5	21.7	67.1
	Average	61.3 ^B	0.0	0.9 ^{DEF}	10.8 ^{CDE}	21.1 ^D	67.2 ^A
All	6	60.1	0.0	1.4	12.3	26.4	60.0
Strains	8	60.2	0.0	1.3	12.1	25.9	60.7

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E,F - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 16. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (119-462 DAYS)

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line	6	92.6	4.5	2.8	0.2	30.75	12.30
W-36	8	89.2	4.5	3.4	0.3	29.44	12.12
	Average	90.9	4.5 ^C	3.1	0.2	30.09	12.21 ^C
Hy-Line	6	86.9	6.6	3.6	0.1	29.76	12.98
W-98	8	84.3	6.6	3.0	0.2	28.50	13.21
	Average	85.6	6.6 ^A	3.3	0.1	29.13	13.10 ^{AB}
Hy-Line	6	91.5	5.4	3.1	0.1	30.51	12.97
CV-22	8	91.3	4.6	4.0	0.1	30.98	12.88
	Average	91.4	5.0 ^{BC}	3.5	0.1	30.74	12.92 ^B
Shaver	6	92.0	5.2	2.7	0.1	30.52	12.32
White	8	91.4	5.3	3.1	0.2	29.94	11.80
	Average	91.7	5.2 ^{BC}	2.9	0.1	30.23	12.06 ^C
DeKalb	6	91.2	5.7	2.8	0.3	31.25	13.13
TX	8	91.1	5.9	2.7	0.3	30.59	12.73
	Average	91.1	5.8 ^{AB}	2.8	0.3	30.92	12.93 ^B
Lohmann	6	93.3	4.7	1.9	0.1	31.79	13.25
LSL-Lite	8	92.4	4.9	2.5	0.2	32.18	13.71
	Average	92.9	4.8 ^{BC}	2.2	0.2	31.98	13.48 ^A
H&N	6	92.6	4.7	2.6	0.1	31.72	12.77
Nick Chick	8	89.1	5.4	2.5	0.2	30.31	12.71
	Average	90.9	5.0 ^{BC}	2.6	0.1	31.01	12.74 ^B
Bovans	6	87.6	6.1	3.0	0.2	30.41	13.08
White	8	89.9	5.0	2.5	0.1	30.97	12.86
	Average	88.7	5.6 ^{ABC}	2.7	0.2	30.69	12.97 ^B
Hisex	6	88.5	5.7	2.4	0.3	30.67	13.19
White	8	87.1	6.3	3.4	0.3	30.15	13.23
	Average	87.8	6.0 ^{AB}	2.9	0.3	30.41	13.21 ^{AB}
Bovans	6	87.8	5.7	3.3	0.2	29.77	12.89
Robust	8	88.2	5.6	3.3	0.2	30.01	12.85
	Average	88.0	5.6 ^{ABC}	3.3	0.2	29.89	12.87 ^B
All	6	90.4	5.4	2.8	0.2	30.71	12.89
Strains	8	89.4	5.4	3.0	0.2	30.30	12.81

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 17. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (119-462 DAYS)

Breeder (Strain)	Population ¹	Feed	Feed	Eggs	Egg	Egg	Age at	
		Consumption (kg/100/hen/d)	Conversion (g egg/g feed)	Per Bird Housed	Production (HD%)	Mass (g/HD)	Mortality (%)	Production (Days)
ISA	6	10.2	0.51	280.6	85.4	52.7	5.6	141
Brown	8	10.1	0.51	278.6	84.6	52.1	7.5	141
	Average	10.1 ^C	0.51 ^A	279.6 ^B	85.0 ^B	52.4 ^B	6.5	141 ^A
Hy-Line	6	9.9	0.52	278.6	83.3	52.4	4.7	140
Brown	8	9.9	0.52	274.6	82.5	51.7	3.0	140
	Average	9.9 ^C	0.52 ^A	276.6 ^B	82.9 ^C	52.0 ^B	3.9	140 ^{BC}
Hy-Line	6	10.2	0.49	280.3	84.7	50.2	6.9	140
Silver Brown	8	10.1	0.50	279.6	84.6	50.3	5.4	140
	Average	10.2 ^{BC}	0.49 ^B	279.9 ^{AB}	84.6 ^{BC}	50.2 ^C	6.2	140 ^{BC}
Bovans	6	10.7	0.51	284.3	86.1	55.4	3.3	140
Brown	8	10.4	0.52	281.2	85.7	55.0	5.3	141
	Average	10.5 ^{AB}	0.52 ^A	282.8 ^{AB}	85.9 ^{AB}	55.2 ^A	4.3	141 ^{AB}
Hisex	6	10.5	0.52	286.8	86.8	54.7	5.5	139
Brown	8	10.6	0.51	286.8	87.1	54.8	6.4	140
	Average	10.6 ^A	0.51 ^A	286.8 ^A	86.9 ^A	54.7 ^A	6.0	139 ^C
DeKalb	6	10.6	0.49	282.0	85.8	52.6	5.5	140
Amber Link	8	10.6	0.48	279.9	85.1	51.8	6.5	140
	Average	10.6 ^A	0.49 ^B	281.0 ^{AB}	85.5 ^{AB}	52.2 ^B	6.0	140 ^{BC}
All	6	10.4	0.51	282.1	85.4	53.0	5.3	140
Strains	8	10.3	0.51	280.1	84.9	52.6	5.7	140

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 18. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (119-462 DAYS)

Breeder	Population ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
ISA	6	61.0	0.0	1.2	10.2	22.7	65.8
Brown	8	60.8	0.0	1.4	10.4	22.8	65.4
	Average	60.9 ^C	0.0	1.3 ^A	10.3 ^B	22.8 ^B	65.6 ^B
Hy-Line	6	62.3	0.0	0.4	6.4	20.3	72.9
Brown	8	62.0	0.0	0.5	7.4	19.9	72.2
	Average	62.1 ^B	0.0	0.4 ^B	6.9 ^C	20.1 ^{BC}	72.5 ^A
Hy-Line Silver	6	58.7	0.0	1.0	13.2	33.4	52.4
Brown	8	59.0	0.0	0.9	14.1	32.4	52.5
	Average	58.9 ^D	0.0	0.9 ^{AB}	13.6 ^A	32.9 ^A	52.4 ^C
Bovans	6	63.6	0.0	0.7	7.3	14.4	77.4
Brown	8	63.4	0.0	0.2	7.0	15.4	77.2
	Average	63.5 ^A	0.0	0.5 ^B	7.2 ^C	14.9 ^D	77.3 ^A
Hisex	6	62.5	0.0	0.3	7.2	18.4	74.0
Brown	8	62.3	0.0	0.7	8.5	18.2	72.5
	Average	62.4 ^B	0.0	0.5 ^B	7.9 ^C	18.3 ^{CD}	73.3 ^A
DeKalb	6	60.7	0.0	1.8	10.0	22.1	66.0
Amber Link	8	60.2	0.0	1.5	11.2	24.4	62.8
	Average	60.4 ^C	0.0	1.6 ^A	10.6 ^B	23.2 ^B	64.4 ^B
All	6	61.5	0.0	0.9	9.1	21.9	68.1
Strains	8	61.3	0.0	0.9	9.8	22.2	67.1

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 19. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (119-462 DAYS)

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	6	91.0	6.1	2.7	0.1	31.18	13.06
Brown	8	87.8	6.3	2.5	0.2	29.88	12.93
	Average	89.4	6.2 ^A	2.6 ^{BC}	0.2	30.53	12.99 ^{BC}
Hy-Line	6	91.7	4.5	3.7	0.1	31.49	12.86
Brown	8	92.1	4.8	3.1	0.0	31.01	12.80
	Average	91.9	4.7 ^B	3.4 ^{AB}	0.0	31.25	12.83 ^C
Hy-Line	6	89.9	4.7	1.9	0.0	30.21	13.14
Silver Brown	8	92.8	4.4	2.8	0.0	31.24	12.98
	Average	91.3	4.6 ^B	2.3 ^C	0.0	30.72	13.06 ^{ABC}
Bovans	6	89.8	6.3	3.8	0.1	31.64	13.74
Brown	8	90.5	5.9	3.4	0.1	31.52	13.23
	Average	90.2	6.1 ^A	3.6 ^A	0.1	31.58	13.49 ^{AB}
Hisex	6	84.3	6.6	3.2	0.1	29.95	13.55
Brown	8	87.8	6.2	3.3	0.1	31.08	13.61
	Average	86.0	6.4 ^A	3.3 ^{ABC}	0.1	30.52	13.58 ^A
DeKalb	6	86.0	6.1	3.0	0.1	29.69	13.59
Amber Link	8	88.4	5.7	3.2	0.2	30.16	13.58
	Average	87.2	5.9 ^{AB}	3.1 ^{ABC}	0.2	29.93	13.59 ^A
All	6	88.8	5.7	3.1	0.1	30.69	13.32
Strains	8	89.9	5.6	3.0	0.1	30.81	13.19

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 20. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder	Population ¹	Feed Consumption	Eggs Per Bird Housed	Egg Production	Mortality	Egg Income	Feed Costs	Days to 0% Production
(Strain)		(kg/100hens/d)		(HD%)	(%)	(\$/hen)	(\$/hen)	(Days)
Hy-Line	6	6.1	8.5	31.3	0.8	0.79	0.56	6
W-36	8	6.4	8.6	32.0	0.4	0.78	0.58	7
	Average	6.2 ^B	8.6	31.7	0.6	0.78	0.57	6 ^B
Hy-Line	6	6.8	8.7	32.4	0.6	0.73	0.63	7
W-98	8	7.1	8.7	32.3	0.6	0.79	0.65	7
	Average	6.9 ^A	8.7	32.4	0.6	0.76	0.64	7 ^{AB}
Hy-Line	6	6.7	8.5	33.7	1.7	0.84	0.58	7
CV-22	8	7.1	8.1	31.8	1.8	0.73	0.60	7
	Average	6.9 ^{AB}	8.3	32.7	1.8	0.78	0.59	7 ^{AB}
Shaver	6	7.1	9.0	34.2	1.9	0.83	0.62	8
White	8	7.6	8.1	33.0	2.0	0.70	0.62	7
	Average	7.3 ^A	8.5	33.6	2.0	0.77	0.62	7 ^{AB}
DeKalb	6	6.7	9.3	34.9	1.3	0.83	0.61	8
TX	8	7.1	8.6	33.6	3.0	0.74	0.61	8
	Average	6.9 ^A	9.0	34.3	2.1	0.78	0.61	8 ^A
Lohmann	6	7.2	9.2	35.7	1.1	0.83	0.63	7
LSL-Lite	8	7.5	9.2	36.9	2.8	0.81	0.64	7
	Average	7.3 ^A	9.2	36.3	2.0	0.82	0.63	7 ^{AB}
H&N	6	7.0	8.6	33.5	2.1	0.80	0.60	7
Nick Chick	8	6.8	8.8	36.3	1.4	0.78	0.57	6
	Average	6.9 ^A	8.7	34.9	1.7	0.79	0.59	7 ^{AB}
Bovans	6	7.4	8.3	34.1	4.9	0.68	0.61	6
White	8	7.1	8.3	34.6	3.0	0.75	0.59	6
	Average	7.2 ^A	8.3	34.3	3.9	0.72	0.60	6 ^B
Hisex	6	7.2	7.8	31.3	2.3	0.68	0.61	7
White	8	7.1	8.5	34.7	3.2	0.78	0.57	7
	Average	7.1 ^A	8.1	33.0	2.8	0.73	0.59	7 ^{AB}
Bovans	6	6.9	8.7	33.0	0.8	0.82	0.62	7
Robust	8	6.9	8.6	32.6	1.2	0.77	0.62	7
	Average	6.9 ^A	8.7	32.8	1.0	0.79	0.62	7 ^{AB}
All	6	6.9	8.7	33.4	1.7	0.78	0.61	7
Strains	8	7.1	8.6	33.8	1.9	0.76	0.61	7

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 21. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder (Strain)	Molt Program ¹	Feed Consumption (kg/100hens/d)	Eggs		Egg Mortality (%)	Egg Income (\$/hen)	Feed Costs (\$/hen)	Days to 0% Production (Days)
			Per Bird Housed	Egg Production (HD%)				
Hy-Line	NM	9.1	20.5	75.5 ^e	0.5 ^{ef}	2.34	0.95	.
W-36	NA20	5.1	2.7	9.9 ^f	0.4 ^f	0.00	0.42	6
	NA25	4.5	2.6	9.6 ^f	0.8 ^{ef}	0.01	0.36	6
Hy-Line	NM	10.5	20.6	76.4 ^{de}	0.0 ^f	2.18	1.10	.
W-98	NA20	5.5	2.6	9.5 ^f	0.6 ^{ef}	0.05	0.44	7
	NA25	4.8	3.0	11.2 ^f	1.1 ^{ef}	0.04	0.38	8
Hy-Line	NM	10.4	19.5	76.6 ^{de}	1.5 ^{ef}	2.16	1.01	.
CV-22	NA20	5.5	2.6	9.9 ^f	1.1 ^{ef}	0.06	0.42	7
	NA25	4.7	2.9	11.7 ^f	2.6 ^{def}	0.12	0.34	7
Shaver	NM	10.9	19.7	77.3 ^{cde}	1.2 ^{ef}	2.13	1.04	.
White	NA20	5.8	2.9	11.8 ^f	3.3 ^{de}	0.08	0.41	8
	NA25	5.4	3.0	11.7 ^f	1.3 ^{ef}	0.09	0.41	7
DeKalb	NM	10.4	21.2	80.8 ^{bcd}	2.8 ^{def}	2.30	1.05	.
TX	NA20	5.6	3.0	11.6 ^f	2.0 ^{def}	0.05	0.43	8
	NA25	4.7	2.7	10.4 ^f	1.7 ^{def}	0.01	0.36	8
Lohmann	NM	10.9	22.0	86.6 ^a	1.7 ^{def}	2.37	1.07	.
LSL-Lite	NA20	5.6	2.8	11.3 ^f	2.8 ^{def}	0.07	0.39	7
	NA25	5.5	2.9	11.0 ^f	1.3 ^{ef}	0.03	0.44	7
H&N	NM	10.6	20.7	83.6 ^{ab}	3.0 ^{def}	2.29	1.02	.
Nick Chick	NA20	5.4	2.7	10.4 ^f	1.6 ^{ef}	0.02	0.40	7
	NA25	4.8	2.8	10.7 ^f	0.7 ^{ef}	0.06	0.34	7
Bovans	NM	11.2	20.0	82.1 ^{abc}	1.1 ^{ef}	2.15	1.07	.
White	NA20	5.5	2.4	10.3 ^f	6.6 ^c	0.00	0.38	6
	NA25	5.1	2.5	10.5 ^f	4.1 ^{cd}	0.00	0.35	6
Hisex	NM	10.4	19.0	78.0 ^{cde}	3.2 ^{de}	2.09	0.94	.
White	NA20	5.7	2.7	10.4 ^f	1.8 ^{def}	0.04	0.43	7
	NA25	5.3	2.7	10.5 ^f	3.3 ^{de}	0.06	0.40	7
Bovans	NM	10.0	20.5	77.5 ^{cde}	0.5 ^{ef}	2.25	1.01	.
Robust	NA20	5.5	2.9	10.8 ^f	0.4 ^f	0.10	0.46	7
	NA25	5.2	2.6	10.2 ^f	2.2 ^{def}	0.03	0.41	7
All	NM	10.4 ^X	20.4 ^Y	79.4	1.6	2.23 ^Y	1.03 ^X	.
Strains	NA20	5.5 ^Y	2.7 ^Z	10.6	2.1	0.05 ^Z	0.42 ^Y	7
	NA25	5.0 ^Z	2.8 ^Z	10.7	1.9	0.04 ^Z	0.38 ^Z	7

X, Y, Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

a,b,c,d - Different letters denote significant strain * molt program interactions (P<.01).

TABLE 22. EFFECT OF WHITE EGG STRAIN AND POPULATION ON BODY WEIGHT OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder	Population ¹	17 Wk Body Wt	66 Wk Body Wt	1st Cycle Wt Gain	Lowest Body Wt	Molt Wt Loss	70 Wk Body Wt
(Strain)		(kg)	(kg)	(%)	(kg)	(%)	(kg)
Hy-Line	6	1.17	1.73	48.4	1.42	17.8	1.50
W-36	8	1.15	1.70	48.7	1.41	17.2	1.47
	Average	1.16 ^{CD}	1.71 ^{CD}	48.6 ^{ABC}	1.41 ^{CD}	17.5	1.49 ^{BCD}
Hy-Line	6	1.24	1.83	47.9	1.50	18.1	1.57
W-98	8	1.24	1.87	51.9	1.54	18.0	1.63
	Average	1.24 ^A	1.85 ^A	49.9 ^{ABC}	1.52 ^A	18.1	1.60 ^A
Hy-Line	6	1.24	1.84	49.8	1.52	17.8	1.59
CV-22	8	1.17	1.84	62.3	1.48	19.9	1.60
	Average	1.20 ^{ABC}	1.84 ^A	56.0 ^A	1.50 ^{AB}	18.9	1.60 ^A
Shaver	6	1.12	1.65	47.7	1.34	19.2	1.48
White	8	1.13	1.62	43.8	1.29	20.5	1.44
	Average	1.13 ^D	1.64 ^E	45.7 ^{BC}	1.31 ^E	19.8	1.46 ^D
DeKalb	6	1.22	1.78	46.9	1.44	19.1	1.53
TX	8	1.21	1.78	47.4	1.44	19.2	1.54
	Average	1.21 ^{AB}	1.78 ^{ABC}	47.2 ^{ABC}	1.44 ^{BC}	19.2	1.54 ^{BC}
Lohmann	6	1.23	1.81	47.7	1.45	19.6	1.55
LSL-Lite	8	1.24	1.79	44.8	1.43	20.5	1.55
	Average	1.23 ^A	1.80 ^{AB}	46.2 ^{BC}	1.44 ^{BC}	20.0	1.55 ^{AB}
H&N	6	1.19	1.69	42.2	1.36	19.4	1.45
Nick Chick	8	1.19	1.75	46.8	1.42	18.9	1.50
	Average	1.19 ^{ABC}	1.72 ^{CD}	44.5 ^{BC}	1.39 ^{CD}	19.2	1.48 ^{CD}
Bovans	6	1.13	1.69	50.5	1.38	18.4	1.54
White	8	1.13	1.69	49.2	1.36	19.3	1.49
	Average	1.13 ^D	1.69 ^{DE}	49.8 ^{ABC}	1.37 ^{DE}	18.9	1.52 ^{BCD}
Hisex	6	1.16	1.77	52.7	1.38	22.0	1.52
White	8	1.18	1.75	48.7	1.41	19.3	1.51
	Average	1.17 ^{BCD}	1.76 ^{BCD}	50.7 ^{AB}	1.39 ^{CD}	20.6	1.52 ^{BCD}
Bovans	6	1.19	1.71	43.9	1.43	16.4	1.52
Robust	8	1.20	1.66	39.3	1.39	16.3	1.50
	Average	1.19 ^{ABC}	1.69 ^{DE}	41.6 ^C	1.41 ^{CD}	16.4	1.51 ^{BCD}
All	6	1.19	1.75	47.8	1.42	18.8	1.52
Strains	8	1.18	1.75	48.3	1.42	18.9	1.52

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 23. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT ON BODY WEIGHT OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder	Molt Program	17 Wk Body Wt	66 Wk Body Wt	1st Cycle Wt Gain	Lowest Body Wt	Molt Wt Loss	70 Wk Body Wt
(Strain)		(kg)	(kg)	(%)	(kg)	(%)	(kg)
Hy-Line W-36	NM	1.15	1.72	49.6	1.64	4.5 ^{gh}	1.69
	NA20	1.16	1.70	45.7	1.29	23.6 ^{ef}	1.43
	NA25	1.15	1.72	50.5	1.30	24.4 ^{def}	1.34
Hy-Line W-98	NM	1.22	1.89	54.7	1.81	4.3 ^{gh}	1.84
	NA20	1.23	1.82	47.8	1.38	24.3 ^{def}	1.50
	NA25	1.25	1.85	47.2	1.37	25.6 ^{bcdef}	1.47
Hy-Line CV-22	NM	1.12	1.90	76.1	1.79	5.7 ^{gh}	1.85
	NA20	1.23	1.80	46.0	1.35	24.9 ^{def}	1.49
	NA25	1.26	1.83	46.0	1.36	26.1 ^{bcde}	1.46
Shaver White	NM	1.12	1.66	48.4	1.60	3.8 ^{gh}	1.64
	NA20	1.16	1.62	40.8	1.19	26.8 ^{bcde}	1.38
	NA25	1.10	1.62	48.0	1.15	28.9 ^{abc}	1.35
DeKalb TX	NM	1.20	1.79	49.5	1.70	5.2 ^{gh}	1.75
	NA20	1.21	1.77	47.1	1.31	25.6 ^{bcdef}	1.44
	NA25	1.23	1.78	44.9	1.31	26.7 ^{bcde}	1.42
Lohmann LSL-Lite	NM	1.23	1.82	49.3	1.76	3.6 ^{gh}	1.79
	NA20	1.23	1.79	45.6	1.33	25.9 ^{bcdef}	1.43
	NA25	1.25	1.79	43.8	1.24	30.6 ^a	1.42
H&N Nick Chick	NM	1.18	1.72	45.7	1.68	2.1 ^h	1.72
	NA20	1.19	1.76	48.1	1.29	26.5 ^{bcde}	1.40
	NA25	1.20	1.68	39.6	1.19	28.9 ^{abc}	1.32
Bovans White	NM	1.14	1.72	50.5	1.69	1.7 ^h	1.72
	NA20	1.11	1.68	53.7	1.22	27.7 ^{abcd}	1.45
	NA25	1.15	1.66	45.3	1.21	27.2 ^{abcde}	1.38
Hisex White	NM	1.15	1.74	52.6	1.62	7.2 ^g	1.67
	NA20	1.17	1.76	50.7	1.31	25.1 ^{cdef}	1.49
	NA25	1.19	1.78	48.7	1.25	29.5 ^{ab}	1.39
Bovans Robust	NM	1.17	1.69	45.1	1.65	2.8 ^h	1.67
	NA20	1.21	1.68	39.8	1.31	22.0 ^f	1.42
	NA25	1.21	1.69	39.8	1.28	24.3 ^{def}	1.42
All Strains	NM	1.17	1.77	52.2 ^X	1.69 ^X	4.1	1.73 ^X
	NA20	1.19	1.74	46.5 ^Y	1.30 ^Y	25.2	1.44 ^Y
	NA25	1.20	1.74	45.4 ^Y	1.27 ^Y	27.2	1.40 ^Z

X, Y, Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.
a, b, c, d, e, f, g, h- Different letters denote significant strain * molt program interactions (P<.01).

TABLE 24. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder	Population ¹	Feed Consumption	Eggs Per Bird Housed	Egg Production	Mortality	Egg Income	Feed Costs	Days to 0% Production
(Strain)		(kg/100hens/d)		(HD%)	(%)	(\$/hen)	(\$/hen)	(Days)
ISA	6	7.4	9.3	36.2	2.9	0.89	0.63	8
Brown	8	7.0	9.4	36.4	1.9	0.87	0.61	9
	Average	7.2	9.4	36.3	2.4	0.88	0.62	8
Hy-Line	6	7.1	9.1	35.4	1.6	0.82	0.62	9
Brown	8	7.0	9.5	35.2	0.2	0.93	0.64	9
	Average	7.0	9.3	35.3	0.9	0.88	0.63	9
Hy-Line	6	7.5	10.0	38.9	3.1	0.93	0.63	10
Silver Brown	8	6.6	9.2	35.6	1.0	0.79	0.59	8
	Average	7.0	9.6	37.3	2.0	0.86	0.61	9
Bovans	6	7.6	9.9	37.0	2.3	0.92	0.68	8
Brown	8	7.6	9.2	35.1	2.7	0.84	0.66	8
	Average	7.6	9.5	36.1	2.5	0.88	0.67	8
Hisex	6	7.0	9.2	35.3	2.1	0.86	0.64	8
Brown	8	7.6	8.9	35.1	1.6	0.75	0.67	8
	Average	7.3	9.1	35.2	1.8	0.81	0.65	8
DeKalb	6	7.4	9.6	36.9	1.4	0.88	0.66	9
Amber Link	8	7.4	9.3	36.3	1.4	0.81	0.65	8
	Average	7.4	9.4	36.6	1.4	0.85	0.66	8
All	6	7.3	9.5	36.6	2.3	0.88	0.64	9
Strains	8	7.2	9.2	35.6	1.4	0.83	0.64	8

¹All strains were housed at a constant density of: 413 cm² equals 64 in². There are no significant differences among these means.

TABLE 25. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder (Strain)	Molt Program ¹	Feed Consumption (kg/100hens/d)	Eggs		Mortality (%)	Egg Income (\$/hen)	Feed Costs (\$/hen)	Days to 0% Production (Days)
			Per Bird Housed	Egg Production (HD%)				
ISA Brown	NM	10.2	21.6 ^b	83.2 ^{ab}	1.9	2.44	0.98	.
	NA20	6.0	3.1 ^e	12.3 ^e	1.7	0.07	0.46	9
	NA25	5.3	3.3 ^{de}	13.3 ^e	3.5	0.14	0.43	8
Hy-Line Brown	NM	10.5	21.0 ^{bc}	80.1 ^{bc}	1.4	2.32	1.01	.
	NA20	5.3	3.5 ^{de}	13.1 ^e	0.9	0.20	0.44	9
	NA25	5.3	3.5 ^{de}	12.8 ^e	0.5	0.12	0.44	8
Hy-Line Silver Brown	NM	9.9	20.4 ^c	78.8 ^c	2.6	2.25	0.95	.
	NA20	5.6	4.6 ^d	18.0 ^d	1.4	0.21	0.42	9
	NA25	5.6	3.8 ^{de}	15.1 ^{de}	2.1	0.12	0.46	9
Bovans Brown	NM	10.7	21.9 ^b	82.4 ^{abc}	0.9	2.44	1.07	.
	NA20	5.9	3.7 ^{de}	13.9 ^e	2.5	0.14	0.47	8
	NA25	6.1	3.0 ^e	11.8 ^e	4.1	0.07	0.47	8
Hisex Brown	NM	11.1	20.9 ^{bc}	81.3 ^{abc}	1.1	2.28	1.07	.
	NA20	5.5	3.3 ^{de}	12.6 ^e	2.1	0.05	0.45	9
	NA25	5.4	3.0 ^e	11.8 ^e	2.3	0.09	0.45	7
DeKalb Amber Link	NM	11.2	21.7 ^b	84.7 ^a	2.4	2.32	1.09	.
	NA20	5.3	3.3 ^{de}	12.6 ^e	0.8	0.14	0.43	8
	NA25	5.7	3.2 ^e	12.4 ^e	1.1	0.08	0.45	8
All Strains	NM	10.6 ^Y	21.3	81.7	1.7	2.34 ^Y	1.03 ^Y	.
	NA20	5.6 ^Z	3.6	13.7	1.6	0.13 ^Z	0.44 ^Z	9
	NA25	5.6 ^Z	3.3	12.9	2.3	0.10 ^Z	0.45 ^Z	8

Y, Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

a, b, c, d, e - Different letters denote significant strain * molt program interactions (P<.01).

TABLE 26. EFFECT OF BROWN EGG STRAIN AND POPULATION ON BODY WEIGHT OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder	Population ¹	17 Wk Body Wt	66 Wk Body Wt	1st Cycle Wt Gain	Lowest Body Wt	Molt Wt Loss	70 Wk Body Wt
(Strain)		(kg)	(kg)	(%)	(kg)	(%)	(kg)
ISA Brown	6	1.45	2.02	40.3	1.68	16.6	1.75
	8	1.48	1.96	33.7	1.67	14.8	1.78
	Average	1.46 ^A	1.99 ^{ABC}	37.0 ^B	1.68 ^{BC}	15.7	1.77 ^C
Hy-Line Brown	6	1.34	1.96	47.1	1.71	13.2	1.77
	8	1.36	1.96	45.0	1.67	15.0	1.74
	Average	1.35 ^B	1.96 ^{BC}	46.1 ^A	1.69 ^{ABC}	14.1	1.76 ^C
Hy-Line Silver Brown	6	1.45	2.07	43.4	1.75	15.2	1.88
	8	1.41	2.08	47.8	1.78	14.5	1.89
	Average	1.43 ^A	2.07 ^A	45.6 ^A	1.76 ^A	14.9	1.89 ^A
Bovans Brown	6	1.46	2.04	40.3	1.72	15.3	1.79
	8	1.46	1.99	37.2	1.65	17.2	1.78
	Average	1.46 ^A	2.01 ^{ABC}	38.7 ^{AB}	1.68 ^{BC}	16.3	1.79 ^{BC}
Hisex Brown	6	1.46	1.96	34.5	1.63	17.0	1.71
	8	1.43	1.95	36.3	1.65	15.3	1.76
	Average	1.44 ^A	1.95 ^C	35.4 ^B	1.64 ^C	16.1	1.74 ^C
DeKalb Amber Link	6	1.45	2.02	40.5	1.71	15.1	1.83
	8	1.47	2.03	38.4	1.73	14.8	1.82
	Average	1.46 ^A	2.03 ^{AB}	39.4 ^{AB}	1.72 ^{AB}	15.0	1.83 ^{AB}
All Strains	6	1.43	2.01	41.0	1.70	15.4	1.79
	8	1.43	2.00	39.7	1.69	15.3	1.80

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 27. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT ON BODY WEIGHT OF HENS IN THE 37th NCLP&MT (462-490 DAYS)

Breeder	Molt Program	17 Wk Body Wt (kg)	66 Wk Body Wt (kg)	1st Cycle Wt Gain (%)	Lowest Body Wt (kg)	Molt Wt Loss (%)	70 Wk Body Wt (kg)
ISA	NM	1.45	1.97	36.6	1.86	5.7	1.89
Brown	NA20	1.47	2.02	38.9	1.61	20.3	1.73
	NA25	1.46	1.98	35.6	1.56	21.1	1.67
Hy-Line	NM	1.36	1.99	46.4	1.92	3.4	1.95
Brown	NA20	1.36	1.95	44.3	1.55	20.5	1.68
	NA25	1.32	1.94	47.5	1.58	18.5	1.63
Hy-Line	NM	1.39	2.04	46.5	1.94	4.8	2.01
Silver Brown	NA20	1.44	2.13	47.8	1.68	21.0	1.84
	NA25	1.45	2.06	42.5	1.67	18.9	1.81
Bovans	NM	1.41	2.00	41.9	1.89	5.3	1.91
Brown	NA20	1.48	2.02	36.8	1.60	20.8	1.75
	NA25	1.48	2.03	37.5	1.57	22.8	1.70
Hisex	NM	1.45	2.00	38.2	1.89	5.0	1.92
Brown	NA20	1.44	1.91	32.9	1.50	21.7	1.70
	NA25	1.44	1.94	35.2	1.52	21.7	1.59
DeKalb	NM	1.43	1.99	39.4	1.94	2.2	1.98
Amber Link	NA20	1.48	2.04	37.8	1.61	21.1	1.76
	NA25	1.48	2.06	41.1	1.61	21.6	1.75
All Strains	NM	1.41	2.00	41.5	1.91 ^Y	4.4 ^Z	1.94 ^X
	NA20	1.44	2.01	39.7	1.59 ^Z	20.9 ^Y	1.74 ^Y
	NA25	1.44	2.00	39.9	1.59 ^Z	20.8 ^Y	1.69 ^Z

X, Y, Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 28. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder (Strain)	Population ¹	Feed	Feed	Eggs	Egg	Egg	Mortality
		Consumption (kg/100/hens/d)	Conversion (g egg/g feed)	Per Bird Housed	Production (HD%)	Mass (g/HD)	
Hy-Line	6	11.4	0.41	162.9	70.2	47.8	3.6
W-36	8	10.4	0.45	161.7	69.1	47.3	4.3
	Average	10.9 ^C	0.43	162.3 ^A	69.7 ^D	47.6 ^D	3.9 ^C
Hy-Line	6	12.4	0.40	163.7	70.8	50.1	4.2
W-98	8	11.7	0.41	156.4	68.8	48.0	7.2
	Average	12.0 ^{AB}	0.41	160.1 ^{AB}	69.8 ^D	49.1 ^{CD}	5.7 ^{BC}
Hy-Line	6	12.1	0.42	149.7	71.6	51.1	5.1
CV-22	8	11.5	0.43	147.0	70.1	49.4	6.7
	Average	11.8 ^{AB}	0.42	148.3 ^{BC}	70.8 ^{CD}	50.2 ^{BC}	5.9 ^{BC}
Shaver	6	12.1	0.40	150.7	74.9	49.6	9.4
White	8	11.4	0.42	146.0	72.6	48.2	9.7
	Average	11.8 ^{AB}	0.41	148.4 ^{BC}	73.8 ^{BC}	48.9 ^{CD}	9.6 ^{AB}
Dekalb	6	12.1	0.41	158.7	72.5	49.0	7.2
TX	8	11.6	0.40	143.5	69.6	46.7	7.9
	Average	11.8 ^{AB}	0.41	151.1 ^{ABC}	71.1 ^{CD}	47.9 ^{CD}	7.5 ^{ABC}
Lohmann	6	12.2	0.43	165.8	78.2	52.9	4.4
LSL-Lite	8	12.3	0.43	151.0	76.3	52.8	11.4
	Average	12.3 ^A	0.43	158.4 ^{AB}	77.3 ^A	52.9 ^A	7.9 ^{ABC}
H&N	6	12.2	0.43	159.2	79.1	52.8	7.9
Nick Chick	8	11.6	0.44	157.5	77.8	51.4	8.1
	Average	11.9 ^{AB}	0.43	158.4 ^{AB}	78.5 ^A	52.1 ^{AB}	8.0 ^{ABC}
Bovans	6	13.0	0.41	139.5	77.1	53.6	12.2
White	8	11.8	0.43	148.0	75.7	51.3	11.1
	Average	12.4 ^A	0.42	143.7 ^C	76.4 ^{AB}	52.4 ^{AB}	11.7 ^A
Hisex	6	12.3	0.41	153.9	75.1	51.0	8.7
White	8	11.7	0.42	144.1	72.6	49.5	10.1
	Average	12.0 ^{AB}	0.42	149.0 ^{BC}	73.8 ^{BC}	50.2 ^{BC}	9.4 ^{AB}
Bovans	6	11.9	0.42	154.9	71.6	52.0	6.2
Robust	8	11.2	0.43	151.7	68.4	48.6	7.7
	Average	11.5 ^{BC}	0.42	153.3 ^{ABC}	70.0 ^D	50.3 ^{ABC}	6.9 ^{BC}
All	6	12.2 ^Y	0.41	155.9	74.1 ^Y	51.0 ^Y	6.9
Strains	8	11.5 ^Z	0.42	150.7	72.1 ^Z	49.3 ^Z	8.4

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 29. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder (Strain)	Population ¹	Egg	Pee				Extra
		Weight (g)	Wee (%)	Small (%)	Medium (%)	Large (%)	Large (%)
Hy-Line	6	66.7	0.0	0.0	0.1	3.2	96.6
W-36	8	67.1	0.0	0.0	0.1	3.4	95.9
	Average	66.9 ^B	0.0	0.0	0.1 ^{BC}	3.3 ^{CDE}	96.2 ^{BC}
Hy-Line	6	69.9	0.0	0.0	0.0	0.5	99.4
W-98	8	69.5	0.0	0.0	0.0	1.2	98.7
	Average	69.7 ^A	0.0	0.0	0.0 ^C	0.9 ^E	99.1 ^A
Hy-Line	6	69.6	0.0	0.0	0.0	1.8	98.1
CV-22	8	69.8	0.0	0.0	0.1	1.1	98.7
	Average	69.7 ^A	0.0	0.0	0.0 ^C	1.4 ^{DE}	98.4 ^{AB}
Shaver	6	64.0	0.0	0.0	0.3	11.9	87.8
White	8	64.6	0.0	0.0	0.2	9.1	90.7
	Average	64.3 ^C	0.0	0.0	0.3 ^{ABC}	10.5 ^A	89.2 ^E
Dekalb	6	67.1	0.0	0.0	0.1	5.0	94.9
TX	8	66.6	0.1	0.0	0.5	4.6	94.9
	Average	66.9 ^B	0.0	0.0	0.3 ^{ABC}	4.8 ^{BC}	94.9 ^{CD}
Lohmann	6	67.4	0.0	0.1	0.2	2.6	97.1
LSL-Lite	8	67.8	0.0	0.0	0.0	2.8	97.1
	Average	67.6 ^B	0.0	0.0	0.1 ^{BC}	2.7 ^{CDE}	97.1 ^{ABC}
H&N	6	65.7	0.0	0.0	0.2	6.5	93.1
Nick Chick	8	65.1	0.0	0.1	0.9	7.0	91.9
	Average	65.4 ^C	0.0	0.0	0.5 ^A	6.7 ^B	92.5 ^D
Bovans	6	68.5	0.0	0.0	0.0	2.7	97.2
White	8	67.1	0.0	0.0	0.0	5.2	94.7
	Average	67.8 ^B	0.0	0.0	0.0 ^C	3.9 ^{CD}	95.9 ^{BC}
Hisex	6	67.2	0.0	0.0	0.2	3.9	95.8
White	8	67.2	0.0	0.2	0.7	2.8	96.0
	Average	67.2 ^B	0.0	0.1	0.4 ^{AB}	3.4 ^{CDE}	95.9 ^{BC}
Bovans	6	69.6	0.0	0.0	0.0	1.4	98.5
Robust	8	68.8	0.0	0.0	0.1	1.2	98.6
	Average	69.2 ^A	0.0	0.0	0.0 ^C	1.3 ^E	98.6 ^{AB}
All	6	67.6	0.0	0.0	0.1	3.9	95.9
Strains	8	67.4	0.0	0.0	0.3	3.9	95.7

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 30. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder (Strain)	Population ¹	Grade	Grade	Cracks	Loss	Egg	Feed
		A	B			Income	Costs
		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line	6	90.8	5.8	3.1	0.2	15.86	8.85
W-36	8	85.8	4.1	3.8	0.0	14.69	8.10
	Average	88.3	5.0	3.5	0.1	15.28	8.47 ^B
Hy-Line	6	85.4	6.7	3.9	0.2	15.17	9.58
W-98	8	75.5	7.8	4.9	0.3	12.90	8.89
	Average	80.4	7.3	4.4	0.2	14.04	9.24 ^A
Hy-Line	6	89.1	6.9	3.9	0.1	14.44	8.37
CV-22	8	88.1	7.3	4.6	0.0	14.15	8.12
	Average	88.6	7.1	4.2	0.1	14.30	8.25 ^{BC}
Shaver	6	87.6	8.3	4.0	0.0	14.39	8.06
White	8	90.3	4.6	4.7	0.4	14.24	7.65
	Average	89.0	6.4	4.3	0.2	14.32	7.86 ^{BC}
Dekalb	6	87.6	8.4	3.8	0.1	15.17	8.79
TX	8	90.0	6.4	3.6	0.0	13.95	8.03
	Average	88.8	7.4	3.7	0.1	14.56	8.41 ^B
Lohmann	6	88.3	9.0	2.5	0.0	15.92	8.65
LSL-Lite	8	89.6	7.7	2.8	0.0	14.62	8.05
	Average	88.9	8.4	2.7	0.0	15.27	8.35 ^{BC}
H&N	6	88.8	6.9	3.8	0.2	15.30	8.19
Nick Chick	8	84.6	6.3	3.3	0.1	14.28	7.89
	Average	86.7	6.6	3.6	0.1	14.79	8.04 ^{BC}
Bovans	6	80.9	10.2	3.5	0.2	12.35	7.81
White	8	85.8	7.1	3.1	0.2	13.80	7.74
	Average	83.4	8.7	3.3	0.2	13.07	7.78 ^C
Hisex	6	83.5	6.9	4.6	0.1	13.97	8.35
White	8	81.2	7.9	4.2	0.1	12.74	7.78
	Average	82.4	7.4	4.4	0.1	13.35	8.06 ^{BC}
Bovans	6	81.2	8.2	4.4	0.1	13.82	8.62
Robust	8	86.5	4.6	2.9	0.3	14.05	8.32
	Average	83.8	6.4	3.7	0.2	13.93	8.47 ^B
All	6	86.3	7.7 ^Y	3.7	0.1	14.64	8.53 ^Y
Strains	8	85.8	6.4 ^Z	3.8	0.1	13.94	8.06 ^Z

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 31. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder (Strain)	Molt Program	Feed Consumption (kg/100 hens/d)	Feed Conversion (g egg/g feed)	Eggs	Egg Production (HD%)	Egg Mass (g/HD)	Mortality (%)
				Per Bird Housed			
Hy-Line W-36	NM	11.1	0.37	141.0	62.3	41.3	7.3
	NA20	10.6	0.46	176.0	73.6	49.2	1.0
	NA25	11.0	0.45	169.8	73.1	52.3	3.6
Hy-Line W-98	NM	12.3	0.34	137.5	59.9	41.8	10.3
	NA20	11.9	0.44	172.8	74.5	52.8	2.2
	NA25	11.9	0.44	169.9	74.9	52.6	4.5
Hy-Line CV-22	NM	12.1	0.35	126.0	61.1	42.2	11.0
	NA20	11.5	0.46	161.9	75.4	52.8	4.2
	NA25	11.8	0.45	157.1	76.0	55.6	2.6
Shaver White	NM	13.0	0.32	131.0	65.4	41.9	14.4
	NA20	10.9	0.47	156.7	78.0	51.8	6.3
	NA25	11.4	0.44	157.4	77.9	53.0	8.0
Dekalb TX	NM	12.6	0.34	136.0	65.0	42.7	10.7
	NA20	11.6	0.44	162.0	75.8	51.2	5.5
	NA25	11.3	0.44	155.4	72.5	49.7	6.4
Lohmann LSL-Lite	NM	12.8	0.37	135.7	68.5	46.2	12.4
	NA20	12.2	0.46	167.7	83.4	56.9	5.4
	NA25	11.8	0.46	171.8	79.8	55.5	5.8
H&N Nick Chick	NM	12.3	0.37	138.2	70.8	45.4	9.9
	NA20	11.5	0.47	172.1	82.5	54.6	5.7
	NA25	11.9	0.46	164.8	82.1	56.4	8.4
Bovans White	NM	12.8	0.37	127.5	69.2	46.8	15.7
	NA20	12.7	0.44	155.4	81.2	56.1	7.0
	NA25	11.8	0.45	148.3	78.8	54.4	12.3
Hisex White	NM	12.1	0.36	121.1	64.2	42.7	13.2
	NA20	12.0	0.45	167.6	79.0	53.7	6.4
	NA25	11.9	0.45	158.3	78.3	54.3	8.5
Bovans Robust	NM	11.6	0.38	142.1	63.6	43.6	7.9
	NA20	11.6	0.44	169.8	72.8	52.4	3.8
	NA25	11.5	0.45	147.9	73.6	55.1	9.1
All Strains	NM	12.3 ^Y	0.36 ^Z	133.6 ^Z	65.0 ^Z	43.4 ^Z	11.3 ^Y
	NA20	11.7 ^Z	0.45 ^Y	166.2 ^Y	77.6 ^Y	53.2 ^Y	4.8 ^Z
	NA25	11.6 ^Z	0.45 ^Y	160.1 ^Y	76.7 ^Y	53.9 ^Y	6.9 ^Z

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 32. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
Hy-Line	NM	66.4	0.0	0.0	0.2	4.5	94.3
W-36	NA20	66.6	0.0	0.0	0.1	3.6	96.3
	NA25	67.7	0.0	0.0	0.0	1.9	98.1
Hy-Line	NM	69.6	0.0	0.0	0.0	0.9	99.0
W-98	NA20	70.1	0.0	0.0	0.0	1.4	98.6
	NA25	69.4	0.0	0.0	0.0	0.3	99.6
Hy-Line	NM	69.2	0.0	0.0	0.1	1.6	98.0
CV-22	NA20	69.2	0.0	0.0	0.0	1.5	98.5
	NA25	70.5	0.0	0.0	0.0	1.3	98.7
Shaver	NM	64.1	0.0	0.0	0.5	13.8	85.7
White	NA20	64.7	0.0	0.0	0.3	10.3	89.3
	NA25	64.2	0.0	0.0	0.0	7.3	92.7
Dekalb	NM	65.9	0.0	0.0	0.8	7.2	92.0
TX	NA20	66.8	0.1	0.0	0.0	4.3	95.6
	NA25	67.8	0.0	0.0	0.0	2.8	97.1
Lohmann	NM	67.4	0.0	0.1	0.1	3.3	96.5
LSL-Lite	NA20	67.4	0.0	0.0	0.0	2.4	97.6
	NA25	68.0	0.0	0.0	0.3	2.4	97.2
H&N	NM	64.2	0.0	0.1	0.4	8.7	90.4
Nick Chick	NA20	65.9	0.0	0.0	0.4	6.0	93.6
	NA25	66.1	0.0	0.0	0.8	5.5	93.5
Bovans	NM	67.7	0.0	0.0	0.0	5.1	94.9
White	NA20	68.1	0.1	0.0	0.0	2.0	97.6
	NA25	67.6	0.0	0.0	0.0	4.6	95.4
Hisex	NM	66.5	0.0	0.1	0.1	5.2	94.2
White	NA20	67.3	0.0	0.0	0.5	3.5	96.0
	NA25	67.8	0.0	0.2	0.7	1.4	97.6
Bovans	NM	68.6	0.0	0.1	0.0	1.2	98.7
Robust	NA20	68.9	0.0	0.0	0.0	2.7	97.3
	NA25	70.2	0.0	0.0	0.1	0.1	99.6
All	NM	67.0 ^Z	0.0	0.0	0.2	5.2 ^Y	94.4 ^Z
Strains	NA20	67.5 ^{YZ}	0.0	0.0	0.1	3.8 ^{YZ}	96.0 ^Y
	NA25	67.9 ^Y	0.0	0.0	0.2	2.8 ^Z	97.0 ^Y

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 33. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line	NM	87.8	6.5 ^{defgh}	5.6	0.2	13.37	7.63
W-36	NA20	82.3	5.4 ^{efgh}	2.3	0.0	15.42	8.84
	NA25	94.8	3.0 ^h	2.5	0.2	17.03	8.94
Hy-Line	NM	79.6	13.3 ^{abc}	6.7	0.5	12.38	8.62
W-98	NA20	76.0	4.7 ^{fgh}	3.3	0.2	14.28	9.65
	NA25	85.8	3.8 ^{fgh}	3.2	0.1	15.44	9.44
Hy-Line	NM	83.8	11.6 ^{abc}	4.6	0.0	11.66	7.53
CV-22	NA20	90.4	5.1 ^{fgh}	4.3	0.2	15.76	8.65
	NA25	91.7	4.6 ^{fgh}	3.7	0.0	15.47	8.57
Shaver	NM	81.2	11.1 ^{abcd}	7.1	0.6	11.95	7.88
White	NA20	93.5	3.4 ^{gh}	2.9	0.1	15.51	7.63
	NA25	92.2	4.8 ^{fgh}	3.0	0.0	15.49	8.06
Dekalb	NM	79.7	15.2 ^a	5.0	0.2	12.20	7.96
TX	NA20	93.2	4.0 ^{fgh}	2.8	0.0	16.05	8.66
	NA25	93.6	3.1 ^{gh}	3.3	0.0	15.44	8.62
Lohmann	NM	80.8	15.2 ^a	4.1	0.0	12.28	7.55
LSL-Lite	NA20	92.1	5.7 ^{efgh}	2.0	0.1	16.47	8.59
	NA25	94.0	4.2 ^{fgh}	1.8	0.0	17.06	8.90
H&N	NM	84.4	10.1 ^{bcde}	5.3	0.2	12.87	7.26
Nick Chick	NA20	92.4	4.2 ^{fgh}	3.5	0.2	16.99	8.42
	NA25	83.3	5.5 ^{efgh}	1.9	0.0	14.51	8.44
Bovans	NM	82.7	13.8 ^{ab}	3.4	0.1	11.70	7.14
White	NA20	91.3	5.2 ^{fgh}	3.3	0.2	15.15	8.43
	NA25	76.1	7.0 ^{defg}	3.1	0.4	12.37	7.76
Hisex	NM	82.4	11.6 ^{abc}	5.7	0.3	11.09	6.86
White	NA20	82.6	6.2 ^{efgh}	3.7	0.0	14.99	8.85
	NA25	82.0	4.4 ^{fgh}	3.7	0.0	13.98	8.48
Bovans	NM	86.1	8.4 ^{cdef}	5.2	0.3	13.45	7.81
Robust	NA20	83.6	4.8 ^{fgh}	1.7	0.3	15.32	9.46
	NA25	81.8	6.0 ^{efgh}	4.1	0.1	13.03	8.14
All	NM	82.8	11.7	5.3 ^Y	0.2	12.30 ^Z	7.62 ^Z
Strains	NA20	87.7	4.9	3.0 ^Z	0.1	15.59 ^Y	8.72 ^Y
	NA25	87.5	4.6	3.0 ^Z	0.1	14.98 ^Y	8.53 ^Y

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

a,b,c,d,e,f,g,h - Different letters denote significant strain*molt program interactions (P<.01).

TABLE 34. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Population ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)		(kg/100 hens/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
ISA	6	12.3	0.41	157.0	75.3	51.4	6.8
Brown	8	11.5	0.41	151.1	71.3	48.7	9.4
	Average	11.9 ^{ABC}	0.41 ^{AB}	154.1	73.3 ^A	50.0 ^{ABC}	8.1
Hy-Line	6	11.9	0.40	148.7	68.2	47.6	6.6
Brown	8	11.0	0.44	159.7	69.5	47.7	4.2
	Average	11.5 ^C	0.42 ^{AB}	154.2	68.8 ^B	47.7 ^{CD}	5.4
Hy-Line	6	12.4	0.40	150.9	73.5	48.5	5.5
Silver Brown	8	10.8	0.41	155.1	67.8	44.7	2.7
	Average	11.6 ^{BC}	0.40 ^{AB}	153.0	70.6 ^{AB}	46.6 ^D	4.1
Bovans	6	12.6	0.42	163.6	75.3	52.9	6.8
Brown	8	11.8	0.44	155.0	73.2	51.6	11.2
	Average	12.2 ^{AB}	0.43 ^A	159.3	74.2 ^A	52.3 ^A	9.0
Hisex	6	12.7	0.40	149.5	73.4	51.5	9.8
Brown	8	12.3	0.41	148.8	71.8	50.4	10.7
	Average	12.5 ^A	0.40 ^{AB}	149.2	72.6 ^A	51.0 ^{AB}	10.2
Dekalb	6	12.8	0.39	157.4	75.8	50.6	9.4
Amber Link	8	12.0	0.40	153.0	70.9	47.2	8.5
	Average	12.4 ^A	0.39 ^B	155.2	73.4 ^A	48.9 ^{BCD}	9.0
All	6	12.4 ^Y	0.40 ^Z	154.5	73.6 ^Y	50.4 ^Y	7.5
Strains	8	11.5 ^Z	0.42 ^Y	153.8	70.7 ^Z	48.4 ^Z	7.8

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 35. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Population ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
ISA	6	66.7	0.0	0.0	0.4	5.2	94.4
Brown	8	66.6	0.0	0.0	0.4	5.6	93.9
	Average	66.7	0.0	0.0	0.4	5.4 ^B	94.2 ^B
Hy-Line	6	69.0	0.0	0.1	0.4	2.9	96.4
Brown	8	68.7	0.0	0.1	0.3	2.4	97.1
	Average	68.8	0.0	0.1	0.4	2.7 ^C	96.7 ^{AB}
Hy-Line	6	66.0	0.0	0.0	0.4	5.9	93.7
Silver Brown	8	65.6	0.0	0.0	0.8	9.8	89.3
	Average	65.8	0.0	0.0	0.6	7.8 ^A	91.5 ^B
Bovans	6	69.2	0.0	0.0	0.3	2.4	97.1
Brown	8	69.7	0.0	0.0	0.3	2.9	96.7
	Average	69.5	0.0	0.0	0.3	2.7 ^C	96.9 ^A
Hisex	6	68.8	0.0	0.0	0.6	2.3	97.1
Brown	8	68.6	0.0	0.0	0.2	3.7	96.0
	Average	68.7	0.0	0.0	0.4	3.0 ^C	96.6 ^{AB}
Dekalb	6	66.6	0.0	0.0	1.0	4.9	93.9
Amber Link	8	66.4	0.0	0.0	0.7	4.2	94.9
	Average	66.5	0.0	0.0	0.9	4.5 ^{BC}	94.4 ^{AB}
All	6	67.7	0.0	0.0	0.5	3.9	95.4
Strains	8	67.6	0.0	0.0	0.5	4.8	94.7

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 36. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	6	90.4	8.1	1.6	0.2	15.22	8.58
Brown	8	82.1	7.7	4.0	0.2	13.48	8.23
	Average	86.3	7.9	2.8	0.2	14.35	8.40
Hy-Line	6	88.2	8.3	3.1	0.2	14.25	8.71
Brown	8	89.7	7.1	3.1	0.3	15.42	8.60
	Average	89.0	7.7	3.1	0.2	14.83	8.66
Hy-Line	6	84.5	6.3	2.1	0.1	13.68	8.56
Silver Brown	8	90.1	6.3	2.9	0.0	14.96	8.41
	Average	87.3	6.3	2.5	0.0	14.32	8.48
Bovans	6	88.4	7.9	3.6	0.2	15.65	9.21
Brown	8	88.1	7.9	3.6	0.2	14.82	8.50
	Average	88.2	7.9	3.6	0.2	15.24	8.86
Hisex	6	76.0	8.6	3.1	0.0	12.35	8.66
Brown	8	82.3	7.8	4.5	0.2	13.38	8.50
	Average	79.1	8.2	3.8	0.1	12.86	8.58
Dekalb	6	83.4	7.3	1.7	0.4	14.01	8.99
Amber Link	8	85.1	6.8	3.8	0.2	13.94	8.72
	Average	84.2	7.1	2.8	0.3	13.98	8.86
All	6	85.1	7.8	2.5 ^Z	0.2	14.19	8.79
Strains	8	86.2	7.3	3.6 ^Y	0.2	14.34	8.49

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 37. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)		(kg/100 hens/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
ISA Brown	NM	12.0	0.38	144.2	69.2	45.8	14.3
	NA20	11.7	0.44	164.5	75.1	52.3	2.4
	NA25	11.9	0.42	153.6	75.5	51.9	7.6
Hy-Line Brown	NM	11.8	0.38	138.7	63.2	43.6	5.8
	NA20	11.0	0.45	162.3	72.7	49.6	6.4
	NA25	11.6	0.43	161.5	70.6	49.8	4.0
Hy-Line Silver Brown	NM	11.8	0.38	146.4	67.9	44.4	6.6
	NA20	11.7	0.41	158.9	72.5	47.8	3.3
	NA25	11.2	0.42	153.7	71.4	47.6	2.3
Bovans Brown	NM	12.3	0.39	155.1	69.4	47.4	11.3
	NA20	12.2	0.44	170.8	77.3	54.5	6.2
	NA25	12.1	0.45	152.0	76.0	54.9	9.5
Hisex Brown	NM	13.4	0.33	126.4	64.2	44.3	15.3
	NA20	11.7	0.46	171.5	78.1	54.6	4.9
	NA25	12.3	0.42	149.6	75.6	54.0	10.4
Dekalb Amber Link	NM	12.8	0.36	142.0	68.3	45.5	11.1
	NA20	12.1	0.42	166.3	76.8	51.0	8.2
	NA25	12.3	0.41	157.3	75.0	50.1	7.6
All Strains	NM	12.4 ^Y	0.37 ^Z	142.1 ^Z	7.0 ^Z	45.2 ^Z	10.7 ^Y
	NA20	11.7 ^Z	0.44 ^Y	165.7 ^X	75.4 ^Y	51.6 ^Y	5.2 ^Z
	NA25	11.9 ^{YZ}	0.42 ^Y	154.6 ^Y	74.0 ^Y	51.4 ^Y	6.9 ^Z

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 38. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
ISA	NM	66.4 ^{efg}	0.0	0.0 ^b	0.8	5.6	93.6
Brown	NA20	67.6 ^{cdef}	0.0	0.0 ^b	0.1	3.8	96.1
	NA25	66.0 ^{fg}	0.0	0.0 ^b	0.4	6.8	92.8
Hy-Line	NM	69.1 ^{abc}	0.0	0.0 ^b	0.0	1.4	98.3
Brown	NA20	67.7 ^{cdef}	0.0	0.3 ^a	0.7	4.0	94.7
	NA25	69.7 ^{ab}	0.0	0.0 ^b	0.3	2.5	97.2
Hy-Line	NM	65.5 ^g	0.0	0.0 ^b	0.3	7.0	92.6
Silver Brown	NA20	65.9 ^{fg}	0.0	0.0 ^b	0.6	7.0	92.4
	NA25	66.2 ^{efg}	0.0	0.0 ^b	1.0	9.5	89.6
Bovans	NM	68.5 ^{bcd}	0.0	0.0 ^b	0.6	4.1	95.1
Brown	NA20	69.3 ^{abc}	0.0	0.0 ^b	0.1	2.2	97.6
	NA25	70.7 ^a	0.0	0.0 ^b	0.3	1.7	98.1
Hisex	NM	69.1 ^{abc}	0.0	0.0 ^b	0.0	2.7	97.2
Brown	NA20	68.0 ^{bcde}	0.0	0.0 ^b	0.7	3.4	95.8
	NA25	68.9 ^{bc}	0.0	0.0 ^b	0.4	2.9	96.7
Dekalb	NM	66.8 ^{defg}	0.0	0.0 ^b	0.0	3.8	95.7
Amber Link	NA20	66.2 ^{efg}	0.0	0.0 ^b	1.1	5.0	93.9
	NA25	66.5 ^{efg}	0.0	0.0 ^b	1.5	4.8	93.7
All	NM	67.6	0.0	0.0	0.3	4.1	95.4
Strains	NA20	67.4	0.0	0.1	0.5	4.2	95.1
	NA25	68.0	0.0	0.0	0.6	4.7	94.7

a,b,c,d,e,f,g - Different letters denote significant strain*molt program interactions (P<.01).

TABLE 39. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	NM	82.7	13.3	3.8	0.3	13.23	7.70
Brown	NA20	82.3	6.2	3.1	0.0	14.62	9.01
	NA25	93.8	4.2	1.4	0.3	15.20	8.50
Hy-Line	NM	82.1	12.7	4.6	0.6	12.71	7.98
Brown	NA20	91.8	4.8	3.2	0.1	15.85	8.64
	NA25	92.9	5.6	1.5	0.0	15.94	9.35
Hy-Line	NM	87.5	9.6	2.9	0.1	13.89	7.93
Silver Brown	NA20	93.6	5.2	2.3	0.0	15.77	9.00
	NA25	80.9	4.2	2.2	0.0	13.30	8.52
Bovans	NM	82.6	12.2	5.0	0.3	14.21	8.53
Brown	NA20	91.6	6.1	2.2	0.1	16.69	9.49
	NA25	90.6	5.3	3.6	0.3	14.80	8.55
Hisex	NM	82.3	12.8	4.9	0.0	11.62	8.13
Brown	NA20	83.5	4.7	2.9	0.1	15.25	9.00
	NA25	71.6	7.2	3.6	0.1	11.72	8.62
Dekalb	NM	84.9	10.8	3.6	0.8	13.16	8.24
Amber Link	NA20	75.5	4.9	3.0	0.0	13.36	9.25
	NA25	92.3	5.6	1.8	0.0	15.41	9.07
All	NM	83.7	11.9 ^Y	4.1 ^Y	0.3	13.14 ^Z	8.09 ^Z
Strains	NA20	86.4	5.3 ^Z	2.8 ^Z	0.1	15.26 ^Y	9.07 ^Y
	NA25	87.0	5.4 ^Z	2.4 ^Z	0.1	14.40 ^{YZ}	8.77 ^Y

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 40. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-MOLTED

Breeder (Strain)	Population ¹	Feed	Feed	Eggs Per Bird	Egg	Egg	Mortality (%)
		Consumption (kg/100 hens/d)	Conversion (g egg/g feed)	Housed	Production (HD%)	Mass (g/HD)	
Hy-Line	6	11.3	0.36	143.6	62.3	41.1	5.0
W-36	8	10.9	0.38	138.5	62.3	41.5	9.6
	Average	11.1	0.37	141.0	62.3	41.3	7.3
Hy-Line	6	12.3	0.35	141.6	61.6	42.7	8.2
W-98	8	12.3	0.33	133.3	58.3	40.9	12.5
	Average	12.3	0.34	137.5	59.9	41.8	10.3
Hy-Line	6	12.7	0.34	122.2	61.8	42.6	10.2
CV-22	8	11.5	0.37	129.8	60.3	41.9	11.8
	Average	12.1	0.35	126.0	61.1	42.2	11.0
Shaver White	6	13.4	0.32	140.6	67.2	43.3	13.7
	8	12.6	0.32	121.4	63.7	40.5	15.0
	Average	13.0	0.32	131.0	65.4	41.9	14.4
Dekalb TX	6	12.7	0.35	142.4	66.4	43.9	11.9
	8	12.5	0.34	129.5	63.5	41.5	9.6
	Average	12.6	0.34	136.0	65.0	42.7	10.7
Lohmann LSL-Lite	6	12.1	0.38	146.8	68.0	46.0	4.3
	8	13.4	0.35	124.7	68.9	46.4	20.6
	Average	12.8	0.37	135.7	68.5	46.2	12.4
H&N Nick Chick	6	12.4	0.37	143.1	70.7	45.5	7.8
	8	12.2	0.37	133.3	71.0	45.3	12.0
	Average	12.3	0.37	138.2	70.8	45.4	9.9
Bovans White	6	13.3	0.36	126.5	69.5	47.1	16.3
	8	12.2	0.38	128.6	69.0	46.4	15.2
	Average	12.8	0.37	127.5	69.2	46.8	15.7
Hisex White	6	12.3	0.36	121.5	65.7	43.7	16.0
	8	11.9	0.36	119.6	62.9	41.8	11.2
	Average	12.1	0.36	120.6	64.3	42.7	13.6
Bovans Robust	6	11.6	0.39	147.2	65.8	45.2	5.1
	8	11.5	0.37	137.1	61.4	41.9	10.7
	Average	11.6	0.38	142.1	63.6	43.6	7.9
All Strains	6	12.4	0.36	137.6	65.9	44.1	9.8
	8	12.1	0.36	129.6	64.1	42.8	12.8

¹All strains were housed at a constant density of: 413 cm² equals 64 in².
There are no significant differences among these means.

TABLE 41. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-MOLTED

Breeder (Strain)	Population ¹	Egg	Pee				Extra
		Weight (g)	Wee (%)	Small (%)	Medium (%)	Large (%)	Large (%)
Hy-Line	6	65.9	0.0	0.0	0.4	5.8	93.7
W-36	8	66.8	0.0	0.0	0.0	3.2	95.0
	Average	66.4 ^{BCD}	0.0	0.0	0.2	4.5 ^{BCD}	94.3 ^{ABC}
Hy-Line	6	69.2	0.0	0.0	0.0	0.7	99.3
W-98	8	70.1	0.0	0.0	0.0	1.1	98.7
	Average	69.6 ^A	0.0	0.0	0.0	0.9 ^D	99.0 ^A
Hy-Line	6	68.9	0.0	0.0	0.0	2.1	97.5
CV-22	8	69.5	0.0	0.0	0.3	1.0	98.4
	Average	69.2 ^A	0.0	0.0	0.1	1.6 ^D	98.0 ^A
Shaver	6	64.6	0.0	0.0	0.6	12.9	86.5
White	8	63.6	0.0	0.0	0.5	14.7	84.8
	Average	64.1 ^E	0.0	0.0	0.5	13.8 ^A	85.7 ^D
Dekalb	6	66.3	0.0	0.0	0.2	6.0	93.9
TX	8	65.4	0.0	0.0	1.5	8.4	90.1
	Average	65.9 ^{CDE}	0.0	0.0	0.8	7.2 ^{BC}	92.0 ^{BC}
Lohmann	6	67.5	0.0	0.2	0.2	2.3	97.5
LSL-Lite	8	67.3	0.0	0.0	0.0	4.4	95.5
	Average	67.4 ^{ABC}	0.0	0.1	0.1	3.3 ^{CD}	96.5 ^{AB}
H&N	6	64.5	0.0	0.0	0.2	7.9	91.5
Nick Chick	8	63.8	0.0	0.2	0.7	9.5	89.3
	Average	64.2 ^{DE}	0.0	0.1	0.4	8.7 ^{AB}	90.4 ^{CD}
Bovans	6	68.0	0.0	0.0	0.0	4.9	95.1
White	8	67.3	0.0	0.0	0.0	5.4	94.7
	Average	67.7 ^{ABC}	0.0	0.0	0.0	5.1 ^{BCD}	94.9 ^{ABC}
Hisex	6	66.4	0.0	0.0	0.0	5.7	94.3
White	8	66.6	0.0	0.2	0.3	4.8	94.0
	Average	66.5 ^{BC}	0.0	0.1	0.2	5.2 ^{BCD}	94.2 ^{ABC}
Bovans	6	68.9	0.0	0.2	0.0	0.9	99.0
Robust	8	68.2	0.0	0.0	0.0	1.5	98.5
	Average	68.6 ^{AB}	0.0	0.1	0.0	1.2 ^D	98.7 ^A
All	6	67.0	0.0	0.0	0.1	4.9	94.8
Strains	8	66.9	0.0	0.0	0.3	5.4	93.9

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 42. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-MOLTED

Breeder (Strain)	Population ¹	Grade	Grade	Cracks	Loss	Egg	Feed
		A	B			Income	Costs
		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line	6	86.7	6.9	6.1	0.3	13.61	7.91
W-36	8	88.8	6.2	5.1	0.0	13.13	7.36
	Average	87.8	6.5	5.6	0.2	13.37	7.63 ^{ABC}
Hy-Line	6	79.7	12.8	7.3	0.4	12.79	8.62
W-98	8	79.5	13.8	6.1	0.6	11.97	8.62
	Average	79.6	13.3	6.7	0.5	12.38	8.62 ^A
Hy-Line	6	87.5	8.7	3.8	0.0	11.59	7.54
CV-22	8	80.1	14.5	5.5	0.0	11.72	7.52
	Average	83.8	11.6	4.6	0.0	11.66	7.53 ^{ABC}
Shaver	6	78.2	14.9	7.0	0.0	12.58	8.50
White	8	84.3	7.3	7.2	1.2	11.32	7.27
	Average	81.2	11.1	7.1	0.6	11.95	7.88 ^{ABC}
Dekalb	6	75.4	18.7	5.6	0.3	12.37	8.17
TX	8	83.9	11.8	4.4	0.0	12.03	7.75
	Average	79.7	15.2	5.0	0.2	12.20	7.96 ^{AB}
Lohmann	6	78.7	17.2	4.2	0.0	13.10	7.89
LSL-Lite	8	82.9	13.1	4.1	0.0	11.46	7.22
	Average	80.8	15.2	4.1	0.0	12.28	7.55 ^{ABC}
H&N	6	81.1	12.0	6.5	0.4	13.02	7.60
Nick Chick	8	87.8	8.2	4.1	0.0	12.72	6.92
	Average	84.4	10.1	5.3	0.2	12.87	7.26 ^{BC}
Bovans	6	82.6	14.8	2.7	0.0	11.56	7.29
White	8	82.8	12.7	4.2	0.3	11.84	6.99
	Average	82.7	13.8	3.4	0.1	11.70	7.14 ^{BC}
Hisex	6	84.0	10.3	5.3	0.4	11.30	6.80
White	8	81.2	12.6	6.0	0.2	10.83	6.86
	Average	82.6	11.5	5.7	0.3	11.06	6.83 ^C
Bovans	6	80.8	11.6	7.6	0.0	13.51	7.81
Robust	8	91.4	5.3	2.9	0.5	13.40	7.80
	Average	86.1	8.4	5.2	0.3	13.45	7.81 ^{ABC}
All	6	81.5	12.8	5.6	0.2	12.54	7.81
Strains	8	84.2	10.5	4.9	0.3	12.04	7.43

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 43. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-MOLTED

Breeder (Strain)	Population ¹	Feed	Feed	Eggs Per Bird	Egg	Egg	Mortality
		Consumption (kg/100 hens/d)	Conversion (g egg/g feed)	Housed	Production (HD%)	Mass (g/HD)	
ISA	6	12.8	0.38	154.9	73.5	47.9	7.5
Brown	8	11.3	0.39	133.5	65.0	43.8	21.0
	Average	12.0 ^B	0.38	144.2	69.2	45.8	14.3
Hy-Line	6	12.4	0.35	131.4	62.0	42.7	8.3
Brown	8	11.2	0.40	146.7	64.5	44.5	2.9
	Average	11.8 ^B	0.38	139.0	63.2	43.6	5.6
Hy-Line	6	12.9	0.36	141.5	69.0	45.3	11.5
Silver Brown	8	10.8	0.41	151.4	66.8	43.6	1.6
	Average	11.8 ^B	0.38	146.4	67.9	44.4	6.6
Bovans	6	12.7	0.38	162.0	70.1	47.7	6.3
Brown	8	11.9	0.40	148.3	68.6	47.2	16.3
	Average	12.3 ^{AB}	0.39	155.1	69.4	47.4	11.3
Hisex	6	13.3	0.33	133.9	63.8	44.0	12.5
Brown	8	13.4	0.34	119.0	64.5	44.7	18.1
	Average	13.4 ^A	0.33	126.4	64.2	44.3	15.3
Dekalb	6	13.3	0.35	144.7	69.3	46.5	11.6
Amber Link	8	12.3	0.36	139.3	67.3	44.5	10.6
	Average	12.8 ^{AB}	0.36	142.0	68.3	45.5	11.1
All	6	12.9 ^Y	0.36	144.7	68.0	45.7	9.6
Strains	8	11.8 ^Z	0.38	139.7	66.1	44.7	11.8

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 44. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-MOLTED

Breeder (Strain)	Population ¹	Egg	Pee			Extra	
		Weight (g)	Wee (%)	Small (%)	Medium (%)	Large (%)	Large (%)
ISA Brown	6	65.2	0.0	0.0	0.5	5.2	94.3
	8	67.7	0.0	0.0	1.0	6.1	92.9
	Average	66.4 ^C	0.0	0.0	0.8	5.6 ^{AB}	93.6 ^B
Hy-Line Brown	6	69.0	0.0	0.0	0.0	1.3	98.4
	8	69.2	0.0	0.0	0.1	1.6	98.2
	Average	69.1 ^A	0.0	0.0	0.1	1.5 ^C	98.3 ^A
Hy-Line Silver Brown	6	65.7	0.0	0.0	0.0	5.0	94.8
	8	65.3	0.0	0.0	0.7	9.0	90.4
	Average	65.5 ^C	0.0	0.0	0.3	7.0 ^A	92.6 ^B
Bovans Brown	6	68.2	0.0	0.0	1.0	4.4	94.2
	8	68.7	0.0	0.0	0.2	3.9	96.0
	Average	68.5 ^{AB}	0.0	0.0	0.6	4.1 ^{ABC}	95.1 ^{AB}
Hisex Brown	6	69.1	0.0	0.0	0.0	2.3	97.6
	8	69.2	0.0	0.0	0.0	3.1	96.8
	Average	69.1 ^A	0.0	0.0	0.0	2.7 ^{BC}	97.2 ^A
Dekalb Amber Link	6	67.3	0.0	0.0	0.0	3.5	96.1
	8	66.3	0.0	0.0	0.0	4.1	95.3
	Average	66.8 ^{BC}	0.0	0.0	0.0	3.8 ^{ABC}	95.7 ^{AB}
All Strains	6	67.4	0.0	0.0	0.3	3.6	95.9
	8	67.7	0.0	0.0	0.3	4.6	94.9

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 45. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-MOLTED

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	6	85.3	13.1	1.7 ^{bc}	0.0	14.44	8.25
Brown	8	80.2	13.4	5.9 ^{ab}	0.5	12.03	7.15
	Average	82.7	13.3	3.8	0.3	13.23	7.70
Hy-Line	6	80.6	13.5	5.4 ^{ab}	0.5	11.92	8.02
Brown	8	83.9	11.9	3.6 ^{abc}	0.6	13.58	7.97
	Average	82.3	12.7	4.5	0.6	12.75	7.99
Hy-Line	6	88.8	9.5	1.6 ^{bc}	0.2	13.52	8.21
Silver Brown	8	86.1	9.7	4.3 ^{abc}	0.0	14.26	7.65
	Average	87.5	9.6	2.9	0.1	13.89	7.93
Bovans	6	81.5	11.9	6.4 ^a	0.2	14.74	9.04
Brown	8	83.6	12.5	3.6 ^{abc}	0.3	13.67	8.02
	Average	82.6	12.2	5.0	0.3	14.21	8.53
Hisex	6	84.0	13.2	2.8 ^{abc}	0.0	12.42	8.57
Brown	8	80.6	12.4	6.9 ^a	0.0	10.82	7.68
	Average	82.3	12.8	4.9	0.0	11.62	8.13
Dekalb	6	84.7	13.0	1.2 ^c	1.3	13.31	8.57
Amber Link	8	85.1	8.6	6.0 ^{ab}	0.4	13.01	7.92
	Average	84.9	10.8	3.6	0.8	13.16	8.24
All	6	84.1	12.4	3.2	0.4	13.39	8.44 ^Y
Strains	8	83.3	11.4	5.0	0.3	12.89	7.73 ^Z

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

a,b,c - Different letters denote significant strain*population interactions (P<.01).

TABLE 46. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 20% BODY WEIGHT LOSS

Breeder (Strain)	Population ¹	Feed	Feed	Eggs	Egg	Egg	Mortality (%)
		Consumption (kg/100 hens/d)	Conversion (g egg/g feed)	Per Bird Housed	Production (HD%)	Mass (g/HD)	
Hy-Line	6	11.3	0.44	172.9	73.9	49.3	1.7
W-36	8	9.9	0.49	180.1	73.4	49.2	0.0
	Average	10.6 ^D	0.47	176.5	73.7 ^F	49.2 ^D	0.9
Hy-Line	6	12.2	0.43	174.5	75.6	53.8	0.4
W-98	8	11.7	0.44	172.3	73.5	51.9	4.3
	Average	12.0 ^{ABC}	0.44	173.4	74.5 ^{EF}	52.9 ^{ABCD}	2.4
Hy-Line	6	11.9	0.45	165.4	76.4	53.5	3.1
CV-22	8	11.1	0.47	158.4	74.5	52.2	5.3
	Average	11.5 ^{BCD}	0.46	161.9	75.4 ^{DEF}	52.8 ^{BCD}	4.2
Shaver	6	10.9	0.47	161.7	78.8	52.2	4.8
White	8	11.0	0.47	150.4	77.5	51.7	8.5
	Average	11.0 ^{CD}	0.47	156.1	78.1 ^{CDE}	51.9 ^{CD}	6.7
Dekalb	6	11.9	0.44	175.5	77.8	53.3	4.1
TX	8	11.3	0.44	149.4	73.8	49.3	6.8
	Average	11.6 ^{BCD}	0.44	162.5	75.8 ^{DEF}	51.3 ^{CD}	5.4
Lohmann	6	12.5	0.45	164.6	83.6	56.6	5.4
LSL-Lite	8	11.8	0.48	173.6	83.6	57.5	4.5
	Average	12.2 ^{AB}	0.47	169.1	83.6 ^A	57.0 ^A	4.9
H&N	6	11.8	0.47	166.9	83.0	55.2	8.1
Nick Chick	8	11.2	0.48	176.3	81.9	53.9	3.7
	Average	11.5 ^{BCD}	0.47	171.6	82.4 ^{AB}	54.6 ^{ABC}	5.9
Bovans	6	13.3	0.44	158.9	84.5	58.8	3.2
White	8	12.1	0.44	154.5	78.7	53.9	9.4
	Average	12.7 ^A	0.44	156.7	81.6 ^{ABC}	56.3 ^{AB}	6.3
Hisex	6	12.5	0.43	176.2	79.9	53.9	6.3
White	8	11.6	0.46	159.0	78.1	53.5	6.5
	Average	12.0 ^{ABC}	0.45	167.6	79.0 ^{BCD}	53.7 ^{ABC}	6.4
Bovans	6	12.3	0.43	175.1	75.0	54.5	3.9
Robust	8	10.9	0.44	165.1	70.7	50.3	3.9
	Average	11.6 ^{ABCD}	0.43	170.1	72.8 ^F	52.4 ^{BCD}	3.9
All	6	12.1 ^Y	0.44	169.2	78.9 ^Y	54.1	4.1
Strains	8	11.3 ^Z	0.46	163.9	76.6 ^Z	52.3	5.3

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E,F - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 47. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 20% BODY WEIGHT LOSS

Breeder (Strain)	Population ¹	Egg	Pee				Extra
		Weight (g)	Wee (%)	Small (%)	Medium (%)	Large (%)	Large (%)
Hy-Line W-36	6	66.4	0.0	0.0	0.0	2.5	97.5
	8	66.8	0.0	0.0	0.2	5.0	94.9
	Average	66.6 ^{DEF}	0.0	0.0	0.1 ^{AB}	3.8 ^B	96.2 ^A
Hy-Line W-98	6	70.2	0.0	0.0	0.0	1.0	99.0
	8	70.0	0.0	0.0	0.0	2.0	98.0
	Average	70.1 ^A	0.0	0.0	0.0 ^B	1.5 ^B	98.5 ^A
Hy-Line CV-22	6	69.5	0.0	0.1	0.0	1.8	98.2
	8	68.9	0.0	0.0	0.0	1.2	98.8
	Average	69.2 ^{AB}	0.0	0.0	0.0 ^B	1.5 ^B	98.5 ^A
Shaver White	6	64.1	0.0	0.0	0.5	13.4	86.1
	8	65.3	0.0	0.0	0.2	6.5	93.3
	Average	64.7 ^F	0.0	0.0	0.3 ^{AB}	9.9 ^A	89.7 ^B
Dekalb TX	6	67.3	0.0	0.0	0.0	4.0	96.0
	8	66.5	0.2	0.0	0.0	4.5	95.3
	Average	66.9 ^{DE}	0.1	0.0	0.0 ^B	4.3 ^B	95.7 ^A
Lohmann LSL-Lite	6	67.3	0.0	0.0	0.0	2.3	97.7
	8	67.5	0.0	0.0	0.0	2.6	97.4
	Average	67.4 ^{BCDE}	0.0	0.0	0.0 ^B	2.4 ^B	97.6 ^A
H&N Nick Chick	6	66.3	0.0	0.0	0.2	5.9	93.9
	8	65.5	0.0	0.0	0.6	6.1	93.3
	Average	65.9 ^{EF}	0.0	0.0	0.4 ^{AB}	6.0 ^{AB}	93.6 ^{AB}
Bovans White	6	68.4	0.0	0.0	0.0	1.7	97.9
	8	67.7	0.1	0.0	0.0	2.6	97.1
	Average	68.1 ^{BCD}	0.1	0.0	0.0 ^B	2.2 ^B	97.5 ^A
Hisex White	6	67.1	0.0	0.0	0.2	4.2	95.6
	8	67.4	0.0	0.0	0.7	2.7	96.3
	Average	67.3 ^{CDE}	0.0	0.0	0.5 ^A	3.5 ^B	96.0 ^A
Bovans Robust	6	69.3	0.0	0.0	0.0	3.5	96.5
	8	68.4	0.0	0.0	0.0	2.0	98.0
	Average	68.9 ^{ABC}	0.0	0.0	0.0 ^B	2.8 ^B	97.3 ^A
All Strains	6	67.6	0.0	0.0	0.1	4.0	95.8
	8	67.4	0.0	0.0	0.2	3.5	96.2

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E,F - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 48. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 20% BODY WEIGHT LOSS

Breeder (Strain)	Population ¹	Grade	Grade	Cracks	Loss	Egg	Feed
		A	B			Income	Costs
		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line W-36	6	90.6	7.5	1.9	0.0	16.80	9.21
	8	72.9	2.8	2.7	0.0	13.89	8.50
	Average	81.8	5.2	2.3	0.0	15.34	8.85 ^{ABC}
Hy-Line W-98	6	83.2	3.2	1.9	0.2	15.42	9.80
	8	68.3	6.5	5.1	0.1	13.24	9.63
	Average	75.7	4.9	3.5	0.1	14.33	9.71 ^A
Hy-Line CV-22	6	90.6	5.4	3.7	0.4	16.10	8.95
	8	90.1	4.8	5.0	0.1	15.42	8.35
	Average	90.4	5.1	4.3	0.2	15.76	8.65 ^{BC}
Shaver White	6	93.9	3.8	2.3	0.0	16.04	7.85
	8	92.8	3.7	3.4	0.2	14.84	7.44
	Average	93.3	3.7	2.8	0.1	15.44	7.64 ^D
Dekalb TX	6	92.6	4.7	2.7	0.0	17.32	9.36
	8	93.8	3.3	2.9	0.0	14.85	8.00
	Average	93.2	4.0	2.8	0.0	16.09	8.68 ^{BC}
Lohmann LSL-Lite	6	92.1	5.9	1.9	0.1	16.14	8.63
	8	92.3	5.6	2.1	0.0	17.08	8.63
	Average	92.2	5.8	2.0	0.1	16.61	8.63 ^{BCD}
H&N Nick Chick	6	92.4	4.4	3.0	0.2	16.45	8.30
	8	91.8	4.1	4.0	0.1	17.32	8.49
	Average	92.1	4.2	3.5	0.2	16.88	8.39 ^{CD}
Bovans White	6	91.7	4.6	3.5	0.2	15.56	8.64
	8	91.4	5.2	3.2	0.2	15.06	8.27
	Average	91.6	4.9	3.4	0.2	15.31	8.45 ^{BCD}
Hisex White	6	75.4	6.1	3.6	0.0	14.62	9.45
	8	89.9	6.3	3.8	0.0	15.35	8.24
	Average	82.6	6.2	3.7	0.0	14.99	8.85 ^{ABC}
Bovans Robust	6	92.4	5.4	1.8	0.5	17.21	10.02
	8	76.9	4.0	1.6	0.2	13.79	8.96
	Average	84.6	4.7	1.7	0.3	15.50	9.49 ^{AB}
All Strains	6	89.5	5.1	2.6	0.2	16.17	9.02 ^Y
	8	86.0	4.6	3.4	0.1	15.08	8.45 ^Z

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E,F - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 49. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 20% BODY WEIGHT LOSS

Breeder (Strain)	Population ¹	Feed	Feed	Eggs Per Bird	Egg	Egg	Mortality
		Consumption (kg/100 hens/d)	Conversion (g egg/g feed)	Housed	Production (HD%)	Mass (g/HD)	
ISA	6	11.8	0.44	167.5	76.1	53.5	2.3
Brown	8	11.5	0.43	161.4	74.2	51.1	2.4
	Average	11.7	0.44	164.5	75.1	52.3 ^{AB}	2.4
Hy-Line	6	11.2	0.44	160.2	72.4	50.0	6.0
Brown	8	10.8	0.46	164.3	73.0	49.2	6.7
	Average	11.0	0.45	162.3	72.7	49.6 ^B	6.4
Hy-Line	6	12.4	0.41	159.5	76.5	50.6	2.2
Silver Brown	8	11.0	0.41	158.2	67.8	44.4	6.6
	Average	11.7	0.41	158.8	72.1	47.5 ^B	4.4
Bovans	6	12.4	0.43	180.5	77.4	54.4	3.6
Brown	8	11.9	0.45	161.2	77.2	54.5	8.7
	Average	12.2	0.44	170.8	77.3	54.5 ^{AB}	6.2
Hisex	6	11.9	0.45	175.4	78.8	54.8	2.5
Brown	8	11.5	0.46	167.3	77.4	54.5	7.7
	Average	11.7	0.46	171.4	78.1	54.7 ^A	5.1
Dekalb	6	12.6	0.43	168.6	81.2	54.1	10.2
Amber Link	8	11.7	0.41	163.9	72.3	48.0	6.1
	Average	12.1	0.42	166.3	76.8	51.0 ^{AB}	8.2
All	6	12.0 ^Y	0.43	168.6	77.0 ^Y	52.9 ^Y	4.5
Strains	8	11.4 ^Z	0.44	162.7	73.6 ^Z	50.3 ^Z	6.4

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 50. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 20% BODY WEIGHT LOSS

Breeder	Population ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
ISA	6	68.2	0.0	0.0	0.1	3.4	96.4
Brown	8	67.1	0.0	0.0	0.0	4.2	95.7
	Average	67.6 ^{AB}	0.0	0.0	0.1	3.8	96.1
Hy-Line	6	68.1	0.0	0.4	1.2	4.8	93.5
Brown	8	67.3	0.0	0.3	0.3	3.3	96.0
	Average	67.7 ^{AB}	0.0	0.3	0.7	4.0	94.7
Hy-Line	6	66.2	0.0	0.0	0.2	5.2	94.6
Silver Brown	8	65.5	0.0	0.0	1.0	8.6	90.1
	Average	65.9 ^B	0.0	0.0	0.6	6.9	92.4
Bovans	6	69.0	0.0	0.0	0.0	2.3	97.7
Brown	8	69.5	0.0	0.0	0.2	2.2	97.5
	Average	69.3 ^A	0.0	0.0	0.1	2.2	97.6
Hisex	6	68.2	0.0	0.0	1.2	2.5	96.2
Brown	8	67.9	0.0	0.0	0.0	4.2	95.5
	Average	68.0 ^{AB}	0.0	0.0	0.6	3.4	95.9
Dekalb	6	66.2	0.0	0.0	0.5	5.2	94.4
Amber Link	8	66.1	0.0	0.0	1.8	4.8	93.4
	Average	66.2 ^B	0.0	0.0	1.1	5.0	93.9
All	6	67.7	0.0	0.1	0.5	3.9	95.5
Strains	8	67.2	0.0	0.1	0.5	4.5	94.7

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 51. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 20% BODY WEIGHT LOSS

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	6	92.4	5.8	1.8	0.0	16.47	9.19
Brown	8	72.1	6.6	4.4	0.0	12.76	8.83
	Average	82.3	6.2	3.1	0.0	14.62	9.01
Hy-Line	6	93.2	4.5	2.4	0.0	15.77	8.66
Brown	8	90.5	5.2	4.0	0.2	15.94	8.63
	Average	91.8	4.8	3.2	0.1	15.85	8.64
Hy-Line	6	93.0	4.8	2.3	0.0	15.74	9.05
Silver Brown	8	92.6	5.4	2.0	0.0	15.51	9.03
	Average	92.8	5.1	2.1	0.0	15.62	9.04
Bovans	6	90.8	6.8	2.4	0.0	17.57	10.18
Brown	8	92.4	5.5	2.0	0.2	15.81	8.79
	Average	91.6	6.1	2.2	0.1	16.69	9.49
Hisex	6	76.5	4.1	2.7	0.0	14.33	9.27
Brown	8	91.5	5.3	3.0	0.2	16.35	8.70
	Average	84.0	4.7	2.8	0.1	15.34	8.98
Dekalb	6	73.7	3.7	1.9	0.0	13.24	9.21
Amber Link	8	77.3	6.1	4.0	0.0	13.47	9.30
	Average	75.5	4.9	3.0	0.0	13.36	9.25
All	6	86.6	4.9	2.2	0.0	15.52	9.26
Strains	8	86.1	5.7	3.3	0.1	14.97	8.88

¹All strains were housed at a constant density of: 413 cm² equals 64 in². There are no significant differences among these means.

TABLE 52. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 25% BODY WEIGHT LOSS

Breeder	Population ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)		(kg/100 hens/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Hy-Line	6	11.7	0.43	171.8	74.3	53.0	3.9
W-36	8	10.4	0.47	168.6	72.1	51.6	3.2
	Average	11.0	0.45	170.2 ^A	73.2 ^E	52.3 ^{BC}	3.6 ^B
Hy-Line	6	12.8	0.42	176.2	75.3	53.9	4.2
W-98	8	11.1	0.46	164.5	74.4	51.3	4.9
	Average	11.9	0.44	170.3 ^A	74.9 ^{CDE}	52.6 ^{ABC}	4.5 ^B
Hy-Line	6	11.8	0.46	159.7	76.6	57.1	2.2
CV-22	8	11.8	0.45	154.6	75.4	54.1	2.9
	Average	11.8	0.45	157.1 ^{AB}	76.0 ^{BCDE}	55.6 ^{AB}	2.6 ^B
Shaver	6	12.1	0.42	148.5	79.1	53.4	10.4
White	8	10.8	0.46	164.4	76.8	52.5	6.1
	Average	11.4	0.44	156.4 ^{AB}	77.9 ^{ABCD}	52.9 ^{ABC}	8.2 ^{AB}
Dekalb	6	11.6	0.43	157.6	73.3	50.0	6.1
TX	8	11.0	0.45	154.1	71.7	49.8	6.7
	Average	11.3	0.44	155.8 ^{AB}	72.5 ^E	49.9 ^C	6.4 ^{AB}
Lohmann	6	12.0	0.47	186.5	82.9	56.1	2.6
LSL-Lite	8	11.6	0.46	158.5	77.2	55.0	8.9
	Average	11.8	0.46	172.5 ^A	80.0 ^{AB}	55.5 ^{AB}	5.8 ^B
H&N	6	12.3	0.45	169.3	83.8	57.7	7.3
Nick Chick	8	11.5	0.46	161.3	80.5	55.2	9.1
	Average	11.9	0.46	165.3 ^{AB}	82.2 ^A	56.4 ^A	8.2 ^{AB}
Bovans	6	12.5	0.43	135.1	77.6	55.0	16.5
White	8	11.2	0.47	159.3	79.4	53.6	9.0
	Average	11.8	0.45	147.2 ^B	78.5 ^{ABC}	54.3 ^{AB}	12.7 ^A
Hisex	6	12.2	0.45	162.7	79.8	55.5	4.6
White	8	11.6	0.45	153.6	76.7	53.0	13.4
	Average	11.9	0.45	158.1 ^{AB}	78.2 ^{ABC}	54.2 ^{ABC}	9.0 ^{AB}
Bovans	6	11.7	0.45	143.4	74.2	56.5	9.5
Robust	8	11.2	0.46	152.4	72.9	53.7	8.8
	Average	11.5	0.45	147.9 ^B	73.6 ^{DE}	55.1 ^{AB}	9.1 ^{AB}
All	6	12.1 ^Y	0.44	161.1	77.7	54.8	6.7
Strains	8	11.2 ^Z	0.46	159.1	75.7	53.0	7.3

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 53. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 25% BODY WEIGHT LOSS

Breeder	Population ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
Hy-Line	6	67.8	0.0	0.0	0.0	1.7	98.4
W-36	8	67.7	0.0	0.0	0.1	2.0	97.9
	Average	67.7 ^{CD}	0.0	0.0	0.0	1.8 ^{CD}	98.1 ^{AB}
Hy-Line	6	70.5	0.0	0.0	0.0	0.0	99.8
W-98	8	68.6	0.0	0.0	0.0	0.6	99.4
	Average	69.5 ^{AB}	0.0	0.0	0.0	0.3 ^D	99.6 ^A
Hy-Line	6	70.1	0.0	0.0	0.0	1.5	98.5
CV-22	8	70.9	0.0	0.0	0.0	1.0	99.0
	Average	70.5 ^A	0.0	0.0	0.0	1.3 ^{CD}	98.7 ^{AB}
Shaver	6	63.5	0.0	0.0	0.0	8.5	91.5
White	8	64.9	0.0	0.0	0.0	6.1	93.9
	Average	64.2 ^E	0.0	0.0	0.0	7.3 ^A	92.7 ^D
Dekalb	6	67.6	0.0	0.0	0.0	4.7	95.2
TX	8	68.2	0.0	0.0	0.0	0.5	99.5
	Average	67.9 ^{BCD}	0.0	0.0	0.0	2.6 ^{BCD}	97.4 ^{ABC}
Lohmann	6	67.5	0.0	0.1	0.4	3.1	96.4
LSL-Lite	8	68.4	0.0	0.0	0.2	1.8	98.0
	Average	68.0 ^{BC}	0.0	0.1	0.3	2.5 ^{BCD}	97.2 ^{ABC}
H&N	6	66.2	0.0	0.0	0.2	5.6	94.0
Nick Chick	8	65.9	0.0	0.0	1.5	5.5	93.1
	Average	66.1 ^D	0.0	0.0	0.8	5.6 ^{AB}	93.5 ^{CD}
Bovans	6	68.9	0.0	0.0	0.0	1.9	98.1
White	8	66.5	0.0	0.0	0.0	6.9	93.1
	Average	67.7 ^{CD}	0.0	0.0	0.0	4.4 ^{ABC}	95.6 ^{BCD}
Hisex	6	68.1	0.0	0.0	0.4	1.8	97.6
White	8	67.4	0.0	0.3	1.1	1.1	97.6
	Average	67.8 ^{BCD}	0.0	0.2	0.7	1.4 ^{CD}	97.6 ^{AB}
Bovans	6	70.7	0.0	0.0	0.0	0.0	100.0
Robust	8	69.7	0.0	0.0	0.2	0.2	99.3
	Average	70.2 ^A	0.0	0.0	0.1	0.1 ^D	99.6 ^A
All	6	68.1	0.0	0.0	0.1	2.9	96.9
Strains	8	67.8	0.0	0.0	0.3	2.6	97.1

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 54. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 25% BODY WEIGHT LOSS

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line	6	95.1	2.9	1.7	0.3	17.19	9.40
W-36	8	93.9	2.9	3.2	0.1	16.79	8.50
	Average	94.5	2.9	2.4	0.2	16.99	8.95 ^{AB}
Hy-Line	6	92.8	4.5	2.8	0.0	17.41	10.45
W-98	8	79.1	3.4	3.6	0.1	13.66	8.55
	Average	85.9	3.9	3.2	0.1	15.53	9.50 ^A
Hy-Line	6	90.4	5.7	3.9	0.0	15.59	8.60
CV-22	8	93.0	3.5	3.5	0.0	15.35	8.54
	Average	91.7	4.6	3.7	0.0	15.47	8.57 ^{ABC}
Shaver	6	90.6	6.8	2.6	0.0	14.40	7.86
White	8	93.7	2.9	3.4	0.0	16.36	8.17
	Average	92.2	4.8	3.0	0.0	15.38	8.02 ^{BC}
Dekalb	6	94.6	2.3	3.1	0.0	15.76	8.83
TX	8	92.1	4.3	3.7	0.0	15.18	8.46
	Average	93.4	3.3	3.4	0.0	15.47	8.64 ^{ABC}
Lohmann	6	94.3	4.2	1.5	0.0	18.56	9.44
LSL-Lite	8	93.7	4.2	2.1	0.0	15.71	8.40
	Average	94.0	4.2	1.8	0.0	17.14	8.92 ^{AB}
H&N	6	93.6	4.3	2.2	0.0	16.75	8.75
Nick Chick	8	74.5	6.5	1.7	0.0	12.63	8.17
	Average	84.1	5.4	1.9	0.0	14.69	8.46 ^{ABC}
Bovans	6	69.6	10.6	4.0	0.4	10.26	7.57
White	8	81.4	4.0	2.4	0.3	14.11	7.88
	Average	75.5	7.3	3.2	0.4	12.19	7.73 ^C
Hisex	6	91.5	3.9	4.6	0.0	15.92	8.73
White	8	70.6	5.0	2.6	0.0	11.69	8.23
	Average	81.1	4.4	3.6	0.0	13.81	8.48 ^{ABC}
Bovans	6	72.1	7.6	4.1	0.0	11.15	8.09
Robust	8	91.5	4.3	4.1	0.1	14.90	8.20
	Average	81.8	6.0	4.1	0.1	13.03	8.15 ^{BC}
All	6	88.5	5.3	3.1	0.1	15.30	8.77
Strains	8	86.3	4.1	3.0	0.1	14.64	8.31

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 55. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 25% BODY WEIGHT LOSS

Breeder (Strain)	Population ¹	Feed	Feed	Eggs	Egg	Egg	Mortality
		Consumption (kg/100 hens/d)	Conversion (g egg/g feed)	Per Bird Housed	Production (HD%)	Mass (g/HD)	
ISA	6	12.3	0.42	149.7	76.7	53.0	9.7
Brown	8	11.6	0.42	158.0	74.6	51.1	5.0
	Average	11.9	0.42	153.9	75.6	52.0 ^{ABC}	7.3
Hy-Line	6	12.3	0.40	157.2	70.4	50.2	4.0
Brown	8	11.0	0.45	165.9	70.9	49.4	4.1
	Average	11.6	0.43	161.5	70.6	49.8 ^{BC}	4.0
Hy-Line	6	11.8	0.42	151.8	74.9	49.8	2.6
Silver Brown	8	10.9	0.42	151.0	67.6	45.6	4.2
	Average	11.3	0.42	151.4	71.3	47.7 ^C	3.4
Bovans	6	12.6	0.44	147.7	78.6	56.9	10.5
Brown	8	11.6	0.45	154.9	73.9	53.2	9.2
	Average	12.1	0.45	151.3	76.2	55.1 ^A	9.8
Hisex	6	12.8	0.42	137.8	77.7	55.9	15.0
Brown	8	11.9	0.43	158.8	73.8	52.5	7.0
	Average	12.4	0.42	148.3	75.7	54.2 ^{AB}	11.0
Dekalb	6	12.6	0.40	159.1	76.4	50.9	6.3
Amber Link	8	12.0	0.41	156.2	73.4	49.1	8.5
	Average	12.3	0.41	157.7	74.9	50.0 ^{BC}	7.4
All	6	12.4 ^Y	0.42	150.6	75.8	52.8	8.0
Strains	8	11.5 ^Z	0.43	157.5	72.3	50.1	6.3

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 56. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 25% BODY WEIGHT LOSS

Breeder	Population ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
ISA	6	66.6	0.0	0.0	0.6	6.9	92.6
Brown	8	65.3	0.0	0.0	0.3	6.6	93.1
	Average	66.0 ^C	0.0	0.0	0.4	6.7 ^{AB}	92.9 ^{AB}
Hy-Line	6	69.8	0.0	0.0	0.0	2.9	97.1
Brown	8	69.5	0.0	0.0	0.6	2.1	97.3
	Average	69.7 ^{AB}	0.0	0.0	0.3	2.5 ^{BC}	97.2 ^A
Hy-Line	6	66.2	0.0	0.0	1.0	7.4	91.7
Silver Brown	8	66.3	0.0	0.0	0.9	11.8	87.4
	Average	66.2 ^C	0.0	0.0	0.9	9.6 ^A	89.5 ^B
Bovans	6	70.6	0.0	0.0	0.0	0.5	99.5
Brown	8	70.8	0.0	0.0	0.4	2.5	97.1
	Average	70.7 ^A	0.0	0.0	0.2	1.5 ^C	98.3 ^A
Hisex	6	69.1	0.0	0.0	0.4	1.9	97.6
Brown	8	68.8	0.0	0.0	0.3	3.8	95.9
	Average	68.9 ^B	0.0	0.0	0.3	2.9 ^{BC}	96.7 ^A
Dekalb	6	66.3	0.0	0.0	2.8	5.9	91.3
Amber Link	8	66.7	0.0	0.0	0.3	3.8	95.7
	Average	66.5 ^C	0.0	0.0	1.6	4.8 ^{ABC}	93.5 ^{AB}
All	6	68.1	0.0	0.0	0.8	4.2	95.0
Strains	8	67.9	0.0	0.0	0.5	5.1	94.4

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 57. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (491-771 DAYS), NON-ANOREXIC WITH A TARGETED 25% BODY WEIGHT LOSS

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	6	93.1	5.3	1.1	0.6	14.74	8.33
Brown	8	95.4	3.1	1.5	0.0	15.84	8.70
	Average	94.2	4.2	1.3	0.3	15.29	8.51
Hy-Line	6	92.2	7.0	0.8	0.0	15.43	9.62
Brown	8	93.7	4.2	2.3	0.0	16.46	9.08
	Average	92.9	5.6	1.5	0.0	15.94	9.35
Hy-Line	6	71.7	4.8	2.4	0.0	11.78	8.42
Silver Brown	8	94.9	3.4	1.7	0.0	15.10	8.56
	Average	83.3	4.1	2.1	0.0	13.44	8.49
Bovans	6	93.2	4.5	1.8	0.5	14.60	8.36
Brown	8	89.4	5.7	4.7	0.2	15.01	8.65
	Average	91.3	5.1	3.3	0.3	14.80	8.50
Hisex	6	67.5	8.8	4.1	0.0	10.23	8.06
Brown	8	75.9	5.8	3.5	0.2	13.10	9.05
	Average	71.7	7.3	3.8	0.1	11.66	8.55
Dekalb	6	92.1	5.9	1.9	0.0	15.52	9.25
Amber Link	8	92.9	5.2	1.9	0.0	15.42	8.94
	Average	92.5	5.6	1.9	0.0	15.47	9.10
All	6	85.0	6.1	2.0	0.2	13.72	8.67
Strains	8	90.4	4.6	2.6	0.1	15.15	8.83

¹All strains were housed at a constant density of: 413 cm² equals 64 in². There are no significant differences among these means.

TABLE 58. EFFECT OF WHITE EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Population ¹	Feed	Feed	Eggs	Egg	Egg	Mortality
		Consumption	Conversion	Per Bird Housed	Production	Mass	
		(kg/100 hens/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Hy-Line	6	10.1	0.47	448.2	75.2	48.6	6.6
W-36	8	9.7	0.49	443.3	74.8	48.8	7.4
	Average	9.9 ^E	0.48	445.8 ^{AB}	75.0 ^E	48.7 ^F	7.0 ^E
Hy-Line	6	10.9	0.46	449.3	75.6	50.6	8.4
W-98	8	10.7	0.46	439.0	74.4	49.7	11.6
	Average	10.8 ^{ABCD}	0.46	444.2 ^{AB}	75.0 ^E	50.1 ^{CDE}	10.0 ^{DE}
Hy-Line	6	10.9	0.46	430.5	76.3	51.1	14.1
CV-22	8	10.6	0.47	431.7	75.8	50.7	17.0
	Average	10.7 ^{BCD}	0.47	431.1 ^B	76.1 ^{DE}	50.9 ^{BCD}	15.6 ^{BCD}
Shaver	6	10.6	0.46	437.0	78.3	49.2	17.8
White	8	10.2	0.47	425.6	77.7	49.0	20.9
	Average	10.4 ^D	0.47	431.3 ^B	78.0 ^{BC}	49.1 ^{EF}	19.3 ^{AB}
Dekalb	6	10.9	0.46	450.6	78.0	50.5	11.8
TX	8	10.5	0.46	429.3	76.0	48.8	17.3
	Average	10.7 ^{BCD}	0.46	440.0 ^{AB}	77.0 ^{CD}	49.7 ^{DEF}	14.5 ^{BCD}
Lohmann	6	11.1	0.47	458.1	81.2	52.5	13.3
LSL-Lite	8	11.2	0.46	447.7	80.8	52.9	21.9
	Average	11.2 ^A	0.46	452.9 ^A	81.0 ^A	52.7 ^A	17.6 ^{BC}
H&N	6	10.8	0.47	454.2	81.6	51.3	17.0
Nick Chick	8	10.7	0.48	449.5	81.7	52.1	18.8
	Average	10.7 ^{BCD}	0.47	451.9 ^A	81.6 ^A	51.7 ^{AB}	17.9 ^{BC}
Bovans	6	11.4	0.47	432.4	80.9	53.4	26.1
White	8	10.8	0.48	441.2	80.6	52.1	24.0
	Average	11.1 ^{AB}	0.47	436.8 ^{AB}	80.7 ^A	52.7 ^A	25.0 ^A
Hisex	6	11.0	0.46	447.6	79.7	51.8	17.3
White	8	10.8	0.46	435.7	78.4	50.9	21.7
	Average	10.9 ^{ABC}	0.46	441.7 ^{AB}	79.0 ^B	51.3 ^{ABC}	19.5 ^{AB}
Bovans	6	10.6	0.47	440.7	76.4	51.5	11.7
Robust	8	10.4	0.47	437.8	75.5	50.3	13.5
	Average	10.5 ^{CD}	0.47	439.3 ^{AB}	76.0 ^{DE}	50.9 ^{BCD}	12.6 ^{CDE}
All	6	10.8 ^Y	0.47	444.9	78.3	51.0	14.4 ^Z
Strains	8	10.6 ^Z	0.47	438.1	77.6	50.5	17.4 ^Y

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E,F - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 59. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Population ¹	Egg	Pee				Extra
		Weight (g)	Wee (%)	Small (%)	Medium (%)	Large (%)	Large (%)
Hy-Line	6	62.5	0.0	0.9	8.5	17.6	72.9
W-36	8	62.8	0.0	0.8	7.8	16.3	74.9
	Average	62.6 ^B	0.0	0.8 ^{ABC}	8.1 ^{ABC}	17.0 ^C	73.9 ^B
Hy-Line	6	65.3	0.0	0.3	5.4	10.8	83.5
W-98	8	65.3	0.0	0.3	5.9	10.5	83.2
	Average	65.3 ^A	0.0	0.3 ^D	5.6 ^{EF}	10.7 ^D	83.4 ^A
Hy-Line	6	65.3	0.0	0.2	4.8	11.8	83.2
CV-22	8	65.4	0.0	0.2	5.0	11.4	83.3
	Average	65.4 ^A	0.0	0.2 ^D	4.9 ^F	11.6 ^D	83.2 ^A
Shaver	6	60.6	0.0	1.5	8.4	25.5	64.6
White	8	61.1	0.0	0.9	8.4	24.9	65.8
	Average	60.9 ^C	0.0	1.2 ^A	8.4 ^{AB}	25.2 ^A	65.2 ^C
Dekalb	6	63.1	0.0	0.7	7.5	16.7	75.0
TX	8	62.5	0.0	0.8	7.3	18.1	73.7
	Average	62.8 ^B	0.0	0.8 ^{ABC}	7.4 ^{ABCD}	17.4 ^C	74.4 ^B
Lohmann	6	63.1	0.0	0.8	6.5	17.4	75.2
LSL-Lite	8	63.3	0.0	0.7	6.9	15.7	76.6
	Average	63.2 ^B	0.0	0.8 ^{ABC}	6.7 ^{CDE}	16.6 ^C	75.9 ^B
H&N	6	61.3	0.0	0.9	9.3	22.9	66.8
Nick Chick	8	61.4	0.0	1.1	8.2	21.5	69.1
	Average	61.4 ^C	0.0	1.0 ^{AB}	8.7 ^A	22.2 ^B	67.9 ^C
Bovans	6	63.7	0.0	0.5	6.5	17.4	75.5
White	8	62.7	0.0	0.8	7.5	18.7	73.0
	Average	63.2 ^B	0.0	0.6 ^{BCD}	7.0 ^{BCDE}	18.0 ^C	74.2 ^B
Hisex	6	63.1	0.0	1.2	6.6	15.7	76.4
White	8	63.2	0.0	1.2	6.8	15.6	76.3
	Average	63.2 ^B	0.0	1.2 ^A	6.7 ^{CDE}	15.7 ^C	76.3 ^B
Bovans	6	64.8	0.0	0.6	6.3	12.3	80.8
Robust	8	64.4	0.0	0.4	6.0	13.0	80.6
	Average	64.6 ^A	0.0	0.5 ^{CD}	6.1 ^{DEF}	12.6 ^D	80.7 ^A
All	6	63.3	0.0	0.8	7.0	16.8	75.4
Strains	8	63.2	0.0	0.7	7.0	16.6	75.6

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E,F - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 60. EFFECT OF WHITE EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line	6	91.5	5.0	3.2	0.2	47.40	21.70
W-36	8	87.4	4.5	3.5	0.2	44.77	20.78
	Average	89.5	4.7 ^B	3.4 ^{ABC}	0.2	46.09	21.24 ^{CD}
Hy-Line	6	85.6	7.3	4.0	0.1	45.83	23.20
W-98	8	79.9	7.2	4.3	0.2	42.16	22.75
	Average	82.8	7.3 ^A	4.1 ^{AB}	0.2	44.00	22.97 ^A
Hy-Line	6	89.7	6.3	3.8	0.1	45.80	21.95
CV-22	8	89.1	5.8	5.1	0.1	45.90	21.64
	Average	89.4	6.1 ^{AB}	4.4 ^A	0.1	45.85	21.80 ^{BC}
Shaver	6	89.4	6.7	3.8	0.0	45.74	21.02
White	8	90.2	5.5	4.0	0.3	44.87	20.08
	Average	89.8	6.1 ^{AB}	3.9 ^{AB}	0.2	45.31	20.55 ^D
Dekalb	6	89.1	7.1	3.6	0.2	47.26	22.56
TX	8	90.1	6.3	3.5	0.2	45.35	21.40
	Average	89.6	6.7 ^A	3.5 ^{ABC}	0.2	46.30	21.98 ^{BC}
Lohmann	6	90.4	6.9	2.6	0.1	48.53	22.50
LSL-Lite	8	90.7	6.5	2.7	0.1	47.59	22.37
	Average	90.5	6.7 ^A	2.7 ^C	0.1	48.06	22.44 ^{AB}
H&N	6	90.0	6.0	3.7	0.1	47.79	21.56
Nick Chick	8	87.2	5.8	2.8	0.1	45.39	21.19
	Average	88.6	5.9 ^{AB}	3.3 ^{BC}	0.1	46.59	21.38 ^{CD}
Bovans	6	84.8	7.9	3.2	0.3	43.56	21.52
White	8	88.0	6.4	2.7	0.2	45.68	21.23
	Average	86.4	7.2 ^A	3.0 ^{BC}	0.2	44.62	21.37 ^{CD}
Hisex	6	85.9	6.5	3.6	0.2	45.34	22.12
White	8	84.0	7.3	4.0	0.2	43.64	21.58
	Average	84.9	6.9 ^A	3.8 ^{ABC}	0.2	44.49	21.85 ^{BC}
Bovans	6	84.4	7.1	4.0	0.2	44.47	22.14
Robust	8	86.7	5.5	3.7	0.2	44.91	21.78
	Average	85.5	6.3 ^A	3.9 ^{AB}	0.2	44.69	21.96 ^{BC}
All	6	88.1	6.7	3.5	0.2	46.17	22.03 ^Y
Strains	8	87.3	6.1	3.6	0.2	45.03	21.48 ^Z

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 61. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Molt Program	Feed Consumption (kg/100 hens/d)	Feed Conversion (g egg/g feed)	Eggs		Mortality (%)	
				Per Bird Housed	Egg Production (HD%)		
Hy-Line W-36	NM	10.1	0.46	438.6	74.0	46.0	10.4
	NA20	9.7	0.50	453.4	75.6	49.2	3.3
	NA25	10.0	0.49	445.3	75.4	50.8	7.4
Hy-Line W-98	NM	11.1	0.44	431.7	73.0	47.5	14.0
	NA20	10.6	0.47	449.8	75.5	51.4	6.7
	NA25	10.6	0.47	451.1	76.6	51.5	9.3
Hy-Line CV-22	NM	11.0	0.44	421.8	74.0	48.2	20.0
	NA20	10.5	0.48	434.3	76.7	51.4	12.6
	NA25	10.7	0.48	437.1	77.6	53.1	14.2
Shaver White	NM	11.2	0.43	425.7	76.4	46.7	23.7
	NA20	9.9	0.49	433.4	78.6	49.9	17.7
	NA25	10.1	0.48	434.9	78.9	50.7	16.6
Dekalb TX	NM	11.2	0.43	440.4	76.5	47.7	17.1
	NA20	10.5	0.47	442.3	78.1	50.7	13.7
	NA25	10.4	0.47	437.2	76.4	50.6	12.8
Lohmann LSL-Lite	NM	11.4	0.45	442.7	79.8	50.7	21.7
	NA20	11.1	0.47	453.3	82.5	53.9	18.3
	NA25	10.9	0.47	462.8	80.8	53.5	12.8
H&N Nick Chick	NM	11.1	0.44	446.2	80.6	48.8	20.9
	NA20	10.6	0.49	455.9	82.0	52.7	15.5
	NA25	10.6	0.49	453.5	82.4	53.5	17.3
Bovans White	NM	11.5	0.45	430.2	80.1	50.7	28.9
	NA20	11.1	0.48	444.7	81.5	54.2	21.0
	NA25	10.6	0.49	435.5	80.6	53.2	25.2
Hisex White	NM	10.9	0.45	422.4	76.8	48.3	26.0
	NA20	10.9	0.47	457.8	80.3	52.7	14.8
	NA25	10.9	0.47	444.7	80.1	53.1	17.7
Bovans Robust	NM	10.6	0.46	440.6	75.2	48.4	13.2
	NA20	10.5	0.47	454.9	76.4	51.3	7.0
	NA25	10.5	0.48	422.3	76.2	53.0	17.7
All Strains	NM	11.0 ^Y	0.45 ^Z	434.0 ^Z	76.6 ^Z	48.3 ^Z	19.6 ^Y
	NA20	10.5 ^Z	0.48 ^Y	448.0 ^Y	78.7 ^Y	51.8 ^Y	13.1 ^Z
	NA25	10.5 ^Z	0.48 ^Y	442.4 ^{YZ}	78.5 ^Y	52.3 ^Y	15.1 ^Z

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 62. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
Hy-Line W-36	NM	62.4	0.0	0.8	7.9	17.6	73.2
	NA20	62.3	0.0	0.9	9.2	17.7	72.3
	NA25	63.2	0.0	0.8	7.3	15.6	76.3
Hy-Line W-98	NM	65.4	0.0	0.3	5.4	10.5	83.9
	NA20	65.5	0.0	0.3	5.8	10.7	83.2
	NA25	65.1	0.0	0.3	5.8	10.9	83.0
Hy-Line CV-22	NM	65.6	0.0	0.1	4.2	11.0	84.5
	NA20	64.8	0.0	0.3	5.7	12.2	81.7
	NA25	65.6	0.0	0.2	4.7	11.6	83.4
Shaver White	NM	61.0	0.0	1.3	8.0	25.5	65.2
	NA20	60.9	0.0	1.1	8.3	25.6	64.9
	NA25	60.6	0.0	1.3	8.7	24.5	65.5
Dekalb TX	NM	62.4	0.0	0.9	7.4	19.7	72.0
	NA20	62.6	0.0	0.9	7.7	18.3	73.0
	NA25	63.4	0.0	0.6	7.0	14.2	78.3
Lohmann LSL-Lite	NM	63.5	0.0	0.5	5.6	16.2	77.6
	NA20	62.9	0.0	0.9	6.8	16.9	75.3
	NA25	63.2	0.0	0.9	7.7	16.5	74.9
H&N Nick Chick	NM	60.6	0.0	1.3	9.3	23.4	65.6
	NA20	61.8	0.1	0.6	8.9	20.0	70.4
	NA25	61.6	0.0	1.1	7.9	23.2	67.7
Bovans White	NM	63.4	0.0	0.6	6.4	18.4	74.5
	NA20	63.5	0.0	0.6	6.5	17.1	75.7
	NA25	62.9	0.0	0.7	8.1	18.6	72.5
Hisex White	NM	62.9	0.0	1.4	7.2	16.6	74.7
	NA20	63.3	0.0	1.0	6.6	16.5	75.9
	NA25	63.4	0.0	1.2	6.3	13.9	78.5
Bovans Robust	NM	64.4	0.0	0.3	6.2	11.9	81.6
	NA20	64.2	0.0	0.4	7.3	14.5	77.7
	NA25	65.3	0.0	0.7	4.9	11.6	82.8
All Strains	NM	63.2	0.0	0.7	6.8	17.1	75.3
	NA20	63.2	0.0	0.7	7.3	16.9	75.0
	NA25	63.4	0.0	0.8	6.8	16.1	76.3

There are no significant differences among these means.

TABLE 63. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder	Molt Program	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Hy-Line W-36	NM	90.2	5.6	4.0	0.2	46.42	20.82
	NA20	84.9	4.9	3.0	0.1	44.20	21.29
	NA25	93.4	3.7	3.0	0.2	47.64	21.61
Hy-Line W-98	NM	84.6	10.3	4.8	0.3	44.68	22.77
	NA20	79.0	5.2	4.1	0.1	42.26	23.18
	NA25	84.7	6.3	3.6	0.1	45.05	22.97
Hy-Line CV-22	NM	87.2	8.7	4.0	0.1	44.59	21.63
	NA20	90.0	4.7	5.2	0.2	46.12	21.66
	NA25	91.0	4.9	4.1	0.0	46.83	22.09
Shaver White	NM	86.5	8.4	4.7	0.3	44.21	21.26
	NA20	91.3	4.4	4.0	0.2	45.79	20.01
	NA25	91.6	5.5	2.9	0.0	45.92	20.38
Dekalb TX	NM	85.9	10.1	3.8	0.2	45.68	22.21
	NA20	90.8	6.0	3.0	0.1	46.66	21.79
	NA25	92.0	3.9	3.8	0.2	46.57	21.94
Lohmann LSL-Lite	NM	87.6	9.3	3.1	0.1	46.75	21.83
	NA20	91.7	5.6	2.6	0.1	48.22	22.44
	NA25	92.3	5.3	2.3	0.1	49.23	23.05
H&N Nick Chick	NM	89.1	7.5	3.3	0.1	46.78	21.26
	NA20	91.8	4.9	3.4	0.1	48.34	21.56
	NA25	85.0	5.2	3.1	0.1	44.65	21.31
Bovans White	NM	87.4	9.4	2.9	0.3	45.14	21.39
	NA20	91.0	5.6	3.1	0.2	47.09	21.87
	NA25	80.7	6.5	2.9	0.2	41.64	20.86
Hisex White	NM	86.6	8.9	4.2	0.3	43.93	20.67
	NA20	84.1	6.1	4.0	0.2	45.48	22.68
	NA25	84.2	5.6	3.2	0.1	44.06	22.21
Bovans Robust	NM	89.0	6.9	3.8	0.3	46.70	21.57
	NA20	84.1	5.6	3.6	0.2	45.34	23.04
	NA25	83.5	6.4	4.2	0.1	42.02	21.27
All Strains	NM	87.4	8.5 ^Y	3.9	0.2	45.49	21.54
	NA20	87.9	5.3 ^Z	3.6	0.1	45.95	21.95
	NA25	87.8	5.3 ^Z	3.3	0.1	45.36	21.77

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 64. EFFECT OF BROWN EGG STRAIN AND POPULATION ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Population ¹	Feed	Feed	Eggs Per Bird	Egg	Egg	Mortality (%)
		Consumption (kg/100 hens/d)	Conversion (g egg/g feed)	Housed	Production (HD%)	Mass (g/HD)	
ISA	6	10.9	0.46	446.7	79.0	51.5	15.3
Brown	8	10.5	0.47	439.1	77.0	50.1	18.6
	Average	10.7 ^{BC}	0.47 ^A	442.9	78.0 ^{AB}	50.8 ^B	16.9 ^A
Hy-Line	6	10.6	0.46	436.2	74.9	49.6	12.8
Brown	8	10.2	0.48	443.6	74.9	49.0	7.2
	Average	10.4 ^C	0.47 ^A	439.9	74.9 ^C	49.3 ^{BC}	10.0 ^B
Hy-Line	6	11.0	0.44	441.2	78.0	48.7	15.5
Silver Brown	8	10.2	0.45	443.7	75.4	47.2	9.3
	Average	10.6 ^{BC}	0.45 ^{BC}	442.4	76.7 ^{BC}	48.0 ^C	12.4 ^{AB}
Bovans	6	11.3	0.47	457.7	79.4	53.8	12.5
Brown	8	10.8	0.48	445.3	78.2	52.7	19.1
	Average	11.1 ^{AB}	0.47 ^A	451.5	78.8 ^A	53.2 ^A	15.8 ^{AB}
Hisex	6	11.2	0.46	445.7	78.9	52.9	17.4
Brown	8	11.2	0.46	444.5	78.5	52.4	18.5
	Average	11.2 ^A	0.46 ^{AB}	445.1	78.7 ^{AB}	52.6 ^A	18.0 ^A
Dekalb	6	11.4	0.44	448.7	79.5	51.2	16.5
Amber Link	8	11.0	0.44	442.3	77.1	49.5	16.5
	Average	11.2 ^A	0.44 ^C	445.5	78.3 ^{AB}	50.3 ^B	16.5 ^A
All	6	11.1 ^Y	0.46	446.0	78.3 ^Y	51.3 ^Y	15.0
Strains	8	10.7 ^Z	0.46	443.1	76.9 ^Z	50.1 ^Z	14.9

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

Y,Z - Different letters denote significant differences (P<.01), comparisons made among population average values.

TABLE 65. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder	Population ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g)	(%)	(%)	(%)	(%)	(%)
ISA	6	63.4	0.0	0.7	5.9	15.3	78.0
Brown	8	63.2	0.0	0.8	5.9	16.0	77.3
	Average	63.3 ^C	0.0	0.7 ^{AB}	5.9 ^{BC}	15.7 ^B	77.6 ^B
Hy-Line	6	65.0	0.0	0.3	3.8	13.3	82.6
Brown	8	64.6	0.0	0.3	4.6	12.5	82.3
	Average	64.8 ^B	0.0	0.3 ^C	4.2 ^D	12.9 ^{BCD}	82.5 ^A
Hy-Line	6	61.7	0.0	0.6	8.3	21.9	69.2
Silver Brown	8	61.7	0.0	0.5	8.4	23.4	67.5
	Average	61.7 ^D	0.0	0.5 ^{BC}	8.4 ^A	22.7 ^A	68.3 ^C
Bovans	6	66.0	0.0	0.4	4.2	9.4	85.8
Brown	8	65.9	0.0	0.2	4.1	10.3	85.3
	Average	66.0 ^A	0.0	0.3 ^C	4.2 ^D	9.9 ^D	85.6 ^A
Hisex	6	65.1	0.0	0.2	4.2	11.5	84.0
Brown	8	64.9	0.0	0.4	5.0	12.8	81.7
	Average	65.0 ^B	0.0	0.3 ^C	4.6 ^{CD}	12.1 ^{CD}	82.9 ^A
Dekalb	6	63.1	0.0	1.1	6.5	14.6	77.8
Amber Link	8	62.8	0.0	0.9	6.6	15.5	76.8
	Average	63.0 ^C	0.0	1.0 ^A	6.6 ^B	15.1 ^{BC}	77.3 ^B
All	6	64.1	0.0	0.5	5.5	14.3	79.6
Strains	8	63.9	0.0	0.5	5.8	15.1	78.5

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 66. EFFECT OF BROWN EGG STRAIN AND POPULATION ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT 119-771 DAYS)

Breeder	Population ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
ISA	6	89.6	6.9	3.6	0.2	47.31	22.26
Brown	8	84.5	7.0	4.0	0.2	44.28	21.75
	Average	87.1	6.9 ^{ABC}	3.8	0.2	45.80	22.00 ^C
Hy-Line	6	89.4	6.3	4.1	0.1	46.54	22.16
Brown	8	89.9	6.0	3.7	0.5	47.39	22.03
	Average	89.6	6.2 ^{BC}	3.9	0.3	46.96	22.10 ^{BC}
Hy-Line	6	86.7	5.8	2.4	0.1	44.82	22.33
Silver Brown	8	89.9	5.6	3.9	0.0	46.84	22.02
	Average	88.3	5.7 ^C	3.2	0.0	45.83	22.18 ^{ABC}
Bovans	6	88.1	7.4	4.3	0.1	48.19	23.64
Brown	8	87.9	7.0	4.8	0.1	47.13	22.38
	Average	88.0	7.2 ^{AB}	4.5	0.1	47.66	23.01 ^{AB}
Hisex	6	80.2	7.5	3.4	0.0	43.08	22.84
Brown	8	84.4	7.8	4.0	0.1	45.29	22.78
	Average	82.3	7.7 ^A	3.7	0.1	44.19	22.81 ^{ABC}
Dekalb	6	84.9	6.9	3.0	0.2	44.79	23.25
Amber Link	8	86.2	6.9	3.7	0.2	45.03	22.96
	Average	85.6	6.9 ^{ABC}	3.3	0.2	44.91	23.10 ^A
All	6	86.5	6.8	3.5	0.1	45.79	22.75
Strains	8	87.1	6.7	4.0	0.2	45.99	22.32

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 67. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON PERFORMANCE OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Molt Program	Feed Consumption (kg/100 hens/d)	Feed Conversion (g egg/g feed)	Eggs Per Bird Housed	Egg Production (HD%)	Egg Mass (g/HD)	Mortality (%)
ISA Brown	NM	10.8	0.46	443.9	78.1	49.7	21.7
	NA20	10.7	0.47	448.8	78.0	51.8	11.2
	NA25	10.6	0.46	436.0	77.8	50.8	18.0
Hy-Line Brown	NM	10.7	0.46	437.3	75.0	49.0	12.0
	NA20	10.0	0.48	442.3	75.4	49.3	10.2
	NA25	10.4	0.47	440.1	74.4	49.7	7.7
Hy-Line Silver Brown	NM	10.8	0.45	447.1	77.4	47.7	14.2
	NA20	10.6	0.44	441.9	76.7	47.8	11.6
	NA25	10.4	0.45	438.3	76.1	48.4	11.4
Bovans Brown	NM	11.2	0.46	458.1	78.4	51.3	16.4
	NA20	11.0	0.48	458.3	79.2	53.6	12.0
	NA25	11.0	0.48	438.1	79.0	54.8	19.0
Hisex Brown	NM	11.8	0.44	430.4	77.1	50.6	23.6
	NA20	10.8	0.49	464.1	80.1	53.8	11.2
	NA25	11.0	0.47	440.7	78.8	53.5	19.1
Dekalb Amber Link	NM	11.6	0.43	445.3	78.5	49.8	19.4
	NA20	10.9	0.45	453.7	78.8	50.7	13.8
	NA25	11.1	0.44	437.6	77.5	50.5	16.2
All Strains	NM	11.1 ^Y	0.45 ^Z	443.7	77.4	49.7 ^Z	17.9 ^Y
	NA20	10.7 ^Z	0.47 ^Y	451.5	78.0	51.2 ^Y	11.7 ^Z
	NA25	10.8 ^Z	0.46 ^Y	438.5	77.3	51.3 ^Y	15.3 ^{YZ}

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 68. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Molt Program	Egg Weight (g)	Pee Wee (%)	Small (%)	Medium (%)	Large (%)	Extra Large (%)
ISA Brown	NM	63.6	0.0	0.8	5.7	14.6	79.0
	NA20	63.7	0.0	0.8	5.7	15.7	77.8
	NA25	62.6	0.0	0.6	6.4	16.8	76.1
Hy-Line Brown	NM	65.4	0.0	0.2	3.7	10.7	85.3
	NA20	63.9	0.0	0.4	5.4	15.4	78.7
	NA25	65.2	0.0	0.2	3.5	12.6	83.4
Hy-Line Silver Brown	NM	61.8	0.0	0.5	7.8	22.2	69.4
	NA20	61.4	0.0	0.7	9.0	23.0	67.3
	NA25	62.0	0.0	0.5	8.3	22.9	68.3
Bovans Brown	NM	65.3	0.0	0.5	4.8	10.4	84.1
	NA20	65.8	0.0	0.4	3.8	11.1	84.6
	NA25	66.8	0.0	0.0	3.9	8.1	88.0
Hisex Brown	NM	65.7	0.0	0.3	3.9	10.1	85.5
	NA20	64.4	0.0	0.3	5.5	14.3	79.7
	NA25	65.0	0.0	0.2	4.4	12.0	83.4
Dekalb Amber Link	NM	63.4	0.0	1.3	5.6	13.1	79.8
	NA20	62.5	0.0	0.7	7.3	17.0	74.9
	NA25	63.1	0.0	0.9	6.8	15.1	77.1
All Strains	NM	64.2	0.0	0.6	5.2	13.5	80.5
	NA20	63.6	0.0	0.6	6.1	16.1	77.2
	NA25	64.1	0.0	0.4	5.6	14.6	79.4

There are no significant differences among these means.

TABLE 69. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 37th NCLP&MT (119-771 DAYS)

Breeder (Strain)	Molt Program	Grade A (%)	Grade B (%)	Cracks (%)	Loss (%)	Egg Income (\$/hen)	Feed Costs (\$/hen)
ISA	NM	87.3	9.4	3.0	0.2	46.50	21.51
Brown	NA20	83.3	6.1	4.6	0.1	44.59	22.72
	NA25	90.5	5.3	3.8	0.3	46.30	21.79
Hy-Line	NM	87.1	8.5	4.1	0.3	46.25	21.87
Brown	NA20	90.7	4.6	4.6	0.1	47.34	21.67
	NA25	91.0	5.4	3.0	0.6	47.31	22.75
Hy-Line	NM	90.8	6.7	2.5	0.1	47.55	21.81
Silver Brown	NA20	92.0	5.4	3.3	0.0	47.45	22.51
	NA25	82.1	5.1	3.7	0.0	42.48	22.20
Bovans	NM	86.7	9.3	3.8	0.2	47.80	23.07
Brown	NA20	89.0	6.0	4.9	0.0	48.69	23.44
	NA25	88.5	6.4	4.8	0.2	46.49	22.52
Hisex	NM	86.3	9.7	3.9	0.0	45.34	22.81
Brown	NA20	83.6	6.7	3.3	0.1	45.95	23.00
	NA25	77.0	6.6	3.9	0.1	41.26	22.63
Dekalb	NM	88.0	8.2	3.4	0.5	46.66	22.99
Amber Link	NA20	78.3	6.7	3.2	0.0	41.78	23.23
	NA25	90.4	5.8	3.5	0.1	46.30	23.10
All	NM	87.7	8.6 ^Y	3.5	0.2	46.68	22.34
Strains	NA20	86.2	5.9 ^Z	4.0	0.1	45.97	22.76
	NA25	86.6	5.8 ^Z	3.8	0.2	45.02	22.50

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

TABLE 70. EFFECT OF WHITE EGG STRAIN AND POPULATION ON HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Population ¹	110 Wk Body Wt	2nd Cycle Wt Gain	2nd Cycle Wt Gain	Total Wt Gain	Total Wt Gain
(Strain)		(kg)	(g)	(%)	(g)	(%)
Hy-Line	6	1.83	332.7	23.5	665.3	57.2
W-36	8	1.75	272.4	20.7	598.8	52.8
	Average	1.79 ^D	302.6	22.1	632.1	55.0
Hy-Line	6	2.02	451.5	31.1	786.5	63.8
W-98	8	2.01	377.6	24.8	774.8	63.0
	Average	2.02 ^A	414.6	28.0	780.6	63.4
Hy-Line	6	1.94	346.9	23.9	703.0	57.5
CV-22	8	1.93	330.1	22.2	764.5	70.1
	Average	1.94 ^B	338.5	23.0	733.7	63.8
Shaver	6	1.72	239.0	18.0	594.2	53.3
White	8	1.71	273.2	20.2	581.0	52.1
	Average	1.72 ^E	256.1	19.1	587.6	52.7
Dekalb	6	1.88	352.7	24.5	663.8	55.2
TX	8	1.83	289.4	20.7	625.7	52.1
	Average	1.86 ^C	321.0	22.6	644.8	53.6
Lohmann	6	1.83	285.8	20.6	605.5	49.5
LSL-Lite	8	1.89	343.9	24.1	648.3	52.3
	Average	1.86 ^C	314.8	22.4	626.9	50.9
H&N	6	1.80	348.9	26.8	611.0	51.5
Nick Chick	8	1.77	267.2	19.6	577.5	48.7
	Average	1.79 ^D	308.0	23.2	594.3	50.1
Bovans	6	1.81	268.3	18.9	676.1	61.5
White	8	1.77	279.5	20.3	640.2	56.9
	Average	1.79 ^D	273.9	19.6	658.2	59.2
Hisex	6	1.80	277.8	20.2	634.1	54.9
White	8	1.83	314.4	21.8	651.6	55.5
	Average	1.81 ^{CD}	296.1	21.0	642.8	55.2
Bovans	6	1.81	291.5	20.3	614.2	51.7
Robust	8	1.78	281.4	19.8	581.2	48.7
	Average	1.79 ^D	286.5	20.0	597.7	50.2
All	6	1.84	319.5	22.8	655.4	55.6
Strains	8	1.83	302.9	21.4	644.4	55.2

¹All strains were housed at a constant density of: 413 cm² equals 64 in².

A,B,C,D,E - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 71. EFFECT OF WHITE EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	110 Wk Body Wt	2nd Cycle Wt Gain	2nd Cycle Wt Gain	Total Wt Gain	Total Wt Gain
(Strain)		(kg)	(g)	(%)	(g)	(%)
Hy-Line W-36	NM	1.73	41.3 ^h	2.6 ^f	576.3 ^{efghi}	50.1 ^{defg}
	NA20	1.77	341.4 ^g	24.2 ^{de}	605.9 ^{defghi}	52.1 ^{defg}
	NA25	1.86	525.0 ^{abc}	39.4 ^a	714.0 ^{bcd}	62.8 ^{abcd}
Hy-Line W-98	NM	1.95	101.3 ^h	5.6 ^f	721.3 ^{bcd}	59.2 ^{abcdefg}
	NA20	2.11	612.2 ^a	41.5 ^a	877.2 ^a	71.4 ^{ab}
	NA25	2.00	530.2 ^{abc}	36.8 ^{abc}	743.4 ^{bc}	59.5 ^{abcdefg}
Hy-Line CV-22	NM	1.88	30.0 ^h	1.8 ^f	760.0 ^{ab}	75.2 ^a
	NA20	1.92	437.5 ^{bcd}	29.6 ^{bcd}	694.2 ^{bcd}	56.6 ^{cdefg}
	NA25	2.01	548.0 ^{ab}	37.8 ^{ab}	747.0 ^{bc}	59.6 ^{abcdef}
Shaver White	NM	1.66	13.8 ^h	1.0 ^f	533.8 ^{hi}	47.7 ^{efg}
	NA20	1.76	376.1 ^{fg}	27.9 ^{cde}	600.0 ^{defghi}	52.6 ^{defg}
	NA25	1.73	378.6 ^{efg}	28.3 ^{bcd}	629.0 ^{cdefghi}	57.7 ^{bcd}
Dekalb TX	NM	1.76	10.0 ^h	1.2 ^f	556.3 ^{fghi}	46.7 ^{efg}
	NA20	1.94	497.1 ^{abcde}	34.4 ^{abc}	732.9 ^{bc}	61.5 ^{abcde}
	NA25	1.88	456.0 ^{bcd}	32.3 ^{abcde}	645.1 ^{bcd}	52.7 ^{defg}
Lohmann LSL-Lite	NM	1.77	-22.5 ^h	-1.1 ^f	541.3 ^{ghi}	44.5 ^{fg}
	NA20	1.89	464.7 ^{bcd}	32.9 ^{abcd}	660.9 ^{bcd}	53.8 ^{cdefg}
	NA25	1.93	502.3 ^{abcd}	35.5 ^{abc}	678.5 ^{bcd}	54.4 ^{cdefg}
H&N Nick Chick	NM	1.69	-23.8 ^h	-1.3 ^f	510.0 ⁱ	43.5 ^g
	NA20	1.83	426.2 ^{cdefg}	30.7 ^{bcd}	638.2 ^{bcd}	53.8 ^{cdefg}
	NA25	1.84	521.7 ^{abc}	40.2 ^a	634.6 ^{bcd}	53.0 ^{cdefg}
Bovans White	NM	1.75	30.0 ^h	1.9 ^f	607.5 ^{defghi}	53.6 ^{cdefg}
	NA20	1.84	390.2 ^{defg}	27.1 ^{cde}	733.2 ^{bc}	68.3 ^{abc}
	NA25	1.78	401.5 ^{defg}	29.7 ^{bcd}	633.8 ^{cdefghi}	55.6 ^{cdefg}
Hisex White	NM	1.75	86.7 ^h	5.5 ^f	608.1 ^{defghi}	53.3 ^{cdefg}
	NA20	1.83	339.0 ^g	22.9 ^e	661.0 ^{bcd}	57.0 ^{bcd}
	NA25	1.85	462.6 ^{bcd}	34.6 ^{abc}	659.3 ^{bcd}	55.3 ^{cdefg}
Bovans Robust	NM	1.73	55.0 ^h	3.4 ^f	556.3 ^{fghi}	47.8 ^{defg}
	NA20	1.82	392.4 ^{defg}	27.7 ^{cde}	608.9 ^{defghi}	50.7 ^{defg}
	NA25	1.84	412.0 ^{cdefg}	29.0 ^{bcd}	628.0 ^{cdefghi}	52.1 ^{defg}
All Strains	NM	1.77 ^Z	32.2	2.1	597.1	52.2
	NA20	1.87 ^Y	427.7	29.9	681.2	57.8
	NA25	1.87 ^Y	473.8	34.4	671.3	56.3

Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.
a,b,c,d,e,f,g,h,i - Different letters denote significant strain*molt program interactions (P<.01).

TABLE 72. EFFECT OF BROWN EGG STRAIN AND POPULATION ON HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Population ¹	110 Wk Body Wt	2nd Cycle Wt Gain	2nd Cycle Wt Gain	Total Wt Gain	Total Wt Gain
(Strain)		(kg)	(g)	(%)	(g)	(%)
ISA	6	2.07	318.8	18.8	628.1	43.8
Brown	8	2.07	297.2	17.4	598.5	41.2
	Average	2.07 ^A	308.0 ^{AB}	18.1 ^A	613.3 ^{BC}	42.5 ^{BC}
Hy-Line	6	2.10	331.4	20.2	764.1	57.7
Brown	8	2.05	307.1	18.5	689.8	51.4
	Average	2.07 ^A	319.2 ^A	19.3 ^A	727.0 ^A	54.5 ^A
Hy-Line	6	2.15	263.3	14.5	700.8	48.7
Silver Brown	8	2.07	180.6	10.1	665.2	47.2
	Average	2.11 ^A	222.0 ^{BC}	12.3 ^B	683.0 ^{AB}	48.0 ^{AB}
Bovans	6	2.09	291.7	16.9	631.3	43.7
Brown	8	2.05	269.1	15.7	589.4	40.8
	Average	2.07 ^A	280.4 ^{ABC}	16.3 ^{AB}	610.3 ^{BC}	42.3 ^{BC}
Hisex	6	2.02	311.2	19.2	565.5	39.1
Brown	8	1.97	208.1	12.6	543.1	38.1
	Average	2.00 ^B	259.6 ^{ABC}	15.9 ^{AB}	554.3 ^C	38.6 ^C
Dekalb	6	2.02	183.7	10.8	564.6	40.5
Amber Link	8	2.06	236.5	13.4	586.4	40.4
	Average	2.04 ^{AB}	210.1 ^C	12.1 ^B	575.5 ^C	40.5 ^{BC}
All	6	2.07	283.3	16.7	642.4	45.6
Strains	8	2.05	249.8	14.6	612.1	43.2

¹All strains were housed at a constant density of: 413 cm² equals 64 in².


A,B,C - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 73. EFFECT OF BROWN EGG STRAIN AND SYNCHRONIZED MOLT PROGRAM ON HENS IN THE 37th NCLP&MT (491-771 DAYS)

Breeder	Molt Program	110 Wk Body Wt	2nd Cycle Wt Gain	2nd Cycle Wt Gain	Total Wt Gain	Total Wt Gain
(Strain)		(kg)	(g)	(%)	(g)	(%)
ISA Brown	NM	1.98	88.8	4.9	535.0	37.3
	NA20	2.12	387.0	22.3	646.0	45.1
	NA25	2.12	448.3	27.0	658.9	45.1
Hy-Line Brown	NM	2.03	72.7	4.0	664.7	49.4
	NA20	2.07	393.8	23.5	716.3	53.1
	NA25	2.12	491.3	30.5	800.0	61.0
Hy-Line Silver Brown	NM	2.04	22.5	1.3	643.8	46.6
	NA20	2.17	322.0	17.6	724.8	50.4
	NA25	2.13	321.4	18.0	680.6	47.0
Bovans Brown	NM	1.98	75.0	4.0	568.8	40.8
	NA20	2.12	366.0	21.1	641.0	43.7
	NA25	2.10	400.2	23.7	621.3	42.2
Hisex Brown	NM	1.97	57.0	3.1	528.0	36.8
	NA20	1.99	290.7	17.2	549.5	38.3
	NA25	2.02	431.2	27.3	585.3	40.8
Dekalb Amber Link	NM	2.02	35.0	1.8	588.8	41.7
	NA20	2.07	310.0	17.9	588.3	39.9
	NA25	2.03	285.3	16.5	549.5	39.7
All Strains	NM	2.00 ^Z	58.5 ^Z	3.2 ^Z	588.2	42.1
	NA20	2.09 ^Y	344.9 ^Y	20.0 ^Y	644.3	45.1
	NA25	2.09 ^Y	396.3 ^Y	23.8 ^X	649.3	46.0

X,Y,Z - Different letters denote significant differences (P<.01), comparisons made among molt program average values.

Figure 1. Hy-Line W-36, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 2. Hy-Line W-98, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 3. Hy-Line CV-22, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 4. Shaver White, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 5. Dekalb TX, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 6. Lohmann LSL-Lite, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 7. H & N “Nick Chick”, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 8. Bovans White, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 9. Hisex White, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 10. Bovans Robust, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted


³ NA = non-anorexic molt program

Figure 11. ISA Brown, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)



¹ kg per 100 Hens
² NM = non-molted
³ NA = non-anorexic molt program

Figure 12. Hy-Line Brown, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 13. Hy-Line Silver Brown, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 14. Bovans Brown, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 15. Hisex Brown, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)




¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Figure 16. Dekalb Amber Link, bi-weekly hen-day egg production and period feed consumption¹ by hen population (6 or 8) and molt program (NM² or NA³)



¹ kg per 100 Hens

² NM = non-molted

³ NA = non-anorexic molt program

Table 74. Entries in the 37th NCLP&MT by Breeder, Stock Suppliers, and Categories

Breeder	Stock	Category ¹	Source
Hy-Line International 2583 240 th Street Dallas Center, IA 50063	W-36	I-A	Hy-Line International 4432 Highway 213, Box 309 Mansfield, GA 30255
	W-98	I-A	Hy-Line International 17458 G. Avenue Perry, IA 50220
	Hy-Line Brown	I-A	(Same)
	Hy-Line Silver Brown	I-A	Dallas Center Research Farm 2418 N Ave. Dallas Center, IA 50063
	CV-22	I-A	(Same)
Lohmann Tierzucht Gmbh Am Seedeich 9-11 . P.O.Box 460 D-27454 Cuxhaven, Germany	Lohmann LSL-Lite	I-A	Hy-Line North America 79 Industrial Rd E-town, PA 17022
H&N International 321 Burnett Ave South, Suite 300 Renton, Washington 98055	H&N “Nick Chick”	I-A	Feather Land Farms 32832 E. Peral Road Coberg, OR 97408
Centurion Poultry, Inc. P.O. Box 591 Lexington, Georgia 30648	Bovans White	I-A	CPI-South Central Hatchery 5087 County Road 35 Bremen, AL 35033
	Bovans Robust	II-A	(Same)
	Bovans Brown	I-A	(Same)
Centurion Poultry, Inc. P.O. Box 591 Lexington, Georgia 30648	Hisex White	I-A	(Same)
	Hisex Brown	I-A	(Same)
Centurion Poultry, Inc. P.O. Box 591 Lexington, Georgia 30648	Dekalb TX	I-A	(Same)
	Dekalb Amber Link	II-A	(Same)
Instiut de Selection Animale (A Hendrix Genetic Company) ISA North America 650 Riverbend Drive, Suite C Kitchener, Ontario N2K 3S2 Canada	Shaver White	II-A	McKinley Hatchery P O Box 1900 772 Queen Street St. Mary's, Ontario N4X 1C2 Canada
	ISA Brown	II-A	(Same)

¹ I = Extensive distribution in southeast United States
 II = Little or no distribution in southeast United States
 III = Unavailable for commercial distribution in United States
 A = Entry requested
 C = Entry not requested