

**FINAL REPORT OF THE
THIRTY NINTH NORTH CAROLINA LAYER PERFORMANCE
AND MANAGEMENT TEST¹**

Vol. 39, No. 5
December 2015

The North Carolina Layer Performance and Management Tests are conducted under the auspices of the North Carolina Layer Performance and Management Program, Prestage Department of Poultry Science, Cooperative Extension Service at North Carolina State University and the North Carolina Department of Agriculture and Consumer Services. The flock is maintained at the Piedmont Research Station-Poultry Unit, Salisbury, North Carolina. Mr. Joe Hampton is Piedmont Research Station Superintendent; Mr. Aaron Sellers is Poultry Unit Manager; Dr. Ramon D. Malheiros, Research Associate, is coordinator of data compilation and statistical analysis; and Dr. Kenneth E. Anderson is Project Leader. The purpose of this program is to assist poultry management teams in evaluation of commercial layer stocks and management systems.

The data presented herein represents the analysis of the first production cycle and molt of the 39th North Carolina Layer Performance and Management Test. Performance summary tables are available for each strain, molt, density and the production systems of Enrichable Cage, Enriched Environmental Housing, and Conventional Cage System.

Copies of current and past reports are maintained for public access at:

<https://poultry.ces.ncsu.edu/layer-performance/>

For further information contact:



Dr. Kenneth E. Anderson
Poultry Science Department
North Carolina State University
Box 7608
Raleigh, NC 27695-7608
Phone (919) 515-5527
FAX (919) 515-7070
ken_anderson@ncsu.edu

¹The use of trade names in this publication does not imply endorsement by the North Carolina Cooperative Extension Service of the products named nor criticism of similar ones not mentioned.

**39th NORTH CAROLINA LAYER PERFORMANCE AND
MANAGEMENT TEST
Volume 39, No. 3**

Report on the First Laying Cycle and Molt

Dates of Importance:

Twenty entries were hatched on July 31, 2013. There were twelve commercial white egg strains and eight commercial brown egg strains that are participating in the current test. The chicks were all sexed according to their genetics (vent, feather, or color), vaccinated for Marek's disease, and wing banded for identification before being transferred to the brood/grow houses.

Table 1 shows the source of the laying stock, strain which was entered, and which environments the strains are participating in the test. Table 91 is a list of the breeder, source of eggs, and entry status of each strain. This report will only present the production data from the hens in Houses 5 and 7 representing the production systems are conventional cages (C), enrichable cages (EC), and the enriched environmental housing system (ECS).

Experimental Components of Importance:

The rearing phase for the systems of conventional cage, enrichable cage, and enriched environmental housing system were completed at 16 weeks, after which the pullets were moved to the laying phase during their 17th week of age.

First cycle production records commenced on November 27, 2013 (17 weeks of age), through the molt period which was induced on November 24, 2014. The molt records commenced on November 24, 2014 (69 weeks of age), and ended on December 22, 2014 (73 weeks of age). The second cycle began on December 22, 2014 (73 weeks of age) and ended on September 2, 2015 (109 weeks of age). This report includes production data summarized from 17 to 69 weeks, 69 to 73 weeks, 73 to 109 weeks, and 17 to 109 weeks of age for each production system and density. A table showing the changes in body weights from 17 to 69 weeks of age, the weight loss during the molt period included in the molt period information, and the changes in body weight during the second cycle.

A maximum of approximately 830 and minimum of 300 white and brown egg pullets per strain were placed at the initiation of the layer portion of the test depending on which of the test environments the strain was entered into.

Strain: Samples of fertile eggs were provided from the breeders according to the rules, which govern the conductance of the test. All eggs were set and hatched concurrently as described in the hatch report (39th Hatch/Serology Report Vol. 39, No. 1). However, due to hatch complications, additional chicks had to be acquired and delivered to the station; the added chicks had hatch dates that were within two days of the original hatch. At hatch the chicks were sexed to remove the males. All strains were sexed according to breeder recommendations (feather, color, or vent sexing).

Table 1. 39th North Carolina Layer Performance and Management Test Strain Code Assignments and Participation

Strain No.	Source of Stock	Source Code	Strain	Participation¹
1	Hendrix-genetics	ISA	Bovans White	C, EC, ECS
2	Hendrix-genetics	ISA	Shaver White	C, EC, ECS
3	Hendrix-genetics	ISA	Dekalb White	C, CF, EC, ECS
4	Hendrix-genetics	ISA	Babcock White	C, EC, ECS
5	Hendrix-genetics	ISA	B-400	C, EC, ECS
6	Hy-Line Int.	HL	W-36	C, CF, EC, ECS
7	Hy-Line Int.	HL	CV-26	C, CF
8	Hy-Line Int.	HL	CV-24	C, CF, EC, ECS
9	Hy-Line Int.	HL	CV-22	C, CF, R
10	Lohmann	L	LSL Lite	C, CF, EC, ECS
11	H&N International	L	H&N Nick Chick	C, CF, EC, ECS
12	Novogen	N	White	C, CF, EC, ECS
13	Tetra Americana	TA	TETRA Amber	C, CF, EC, ECS
14	Tetra Americana	TA	TETRA Brown	C, CF, EC, ECS
15	Novogen	N	Brown	C, CF, EC, ECS
16	Lohmann	L	LB-Lite	C, CF, EC, ECS
17	Hy-Line Int.	HL	Silver Brown	C, CF, EC, ECS, R
18	Hy-Line Int.	HL	Brown	C, CF, EC, ECS, R
19	Hendrix-genetics	ISA	ISA Brown	C, CF, EC, ECS
20	Hendrix-genetics	ISA	Bovans Brown	C, CF, EC, ECS

¹ Participation for each strain in the different components of the tests are indicated by the following codes, a strain may have more than one code: Cage=C; Enrichable Cage=EC; Enriched Colony Housing System=ECS; Cage Free = CF; Range = R. If the code is in bold they are included in this report.

Pullet Housing and Management:

The pullets were reared in the environment to which they would be in during the laying phase (39th NCLP&MT Grow Report, Vol.39, No. 2). White egg strains occupied approximately 60% and brown egg strains occupied the other 40% of cage replicates. Individual chicks were identified by strain assignment codes that indicated the cage arrangement, replicate identification numbers, and the strain assignments for brood-grow House 8. Strain codes are maintained by the PI and Poultry Unit Manager for strain and replicate identification and record keeping. Individual birds were identified by a permanent identification tag applied at hatch; at the time they were transferred to the laying house, each hen was retagged with the laying house replicate number, indicating row, level and replicate.

The brood-grow system in House 8 was used to rear the pullets for the conventional cage, enrichable cage, and the enriched environmental housing system. House 8 is an environmentally controlled, windowless brood-grow facility with four rooms, each containing 72 replicates within a Big Dutchman quad-deck cage layout. This system allows for a total of 3,744 pullets per room. This study utilized all four rooms for a total of 11,062 pullets. The white and brown egg strains were randomly assigned to the replicates in a restricted randomized manner with the restrictions being that all strains were equally represented in all rooms, rows, and levels, as described in the grow report (39th NCLP&MT Grow Report, Vol.39, No. 2). Thirteen white-egg or brown-egg chicks were in the same 60.9 x 66.0 cm (24 x 26 in) cage during the entire 16 week rearing period. Rearing density was 310 cm² (48 in²) for both the white and brown-egg layers.

Pullet Management and Nutrition:

Pullets were fed *ad libitum* by hand daily. Feed consumption and body weights were monitored bi-weekly beginning at 2 weeks of age. All mortality was recorded daily, but mortality attributed to the removal of males (sex slips) and accidental deaths from a replicate have been excluded from the 39th NCLP&MT Grow Report.

Layer Housing and Cage Layout Description:

The pullets were moved to the laying facilities, Houses 5 and 7, in accordance with NCSU IACUC approved methods. The strains were randomly assigned to replicate cages with white-egg strains occupying approximately 60% and brown-egg strains the other 40% of the replicates being intermingled throughout the houses such that all strains were approximately equally represented in all rows, levels and cage sizes. Both houses contain a feeder system that allows feed consumption to be determined by replicate. The replicates are equipped with feed hoppers to supply and monitor feed consumption for each individual replicate. The feed is distributed by an automatic feeding system. Laying Hen Cage Facilities reported in this test consist of two houses shown in Table 2.

House 5 is a standard height, windowless, forced ventilated laying house with the enriched and enrichable colony cages that utilize a belted manure handling system. It has five rows of the FDI Condo Layer System; three rows are Enriched Environmental Housing Systems and two rows are Enrichable Cages. As with the other houses, each side of a row has been designated as a bank and each bank is divided into nine 8 ft replicates per level. The replicates contain either four 60.9 cm (24 in) cages or a single 243.8 cm (96 in) cage. The 243.8 cm (96 in) cages were equipped with a nesting area of 60.9 x 22.9 x 30.5 cm or 731.5 cm² (24 x 19 x 12 in or 288 in²) and 2 roosts of 1.9 x 5.1 x 121.9 cm (¾ x 2 x 48 in) positioned 7.6 cm (3 in) off the floor. The cages in both houses are 66.0 cm (26 in) deep. Therefore, when the bird population is held constant at 9 hens per cage in the conventional cages and 36 or 18 hens per cage in the enrichable cages and enriched environmental housing system, the densities 175.3, 175.3 and 353.1 cm² (69, 69, and 139 in²), respectively. House 5 population is 8,262 hens.

House 7 is a standard height, windowless, enclosed force ventilated house. It had four rows of a FDI Tri-Deck Stacked Layer Cage System that utilizes a belted manure handling system. There is 18.3 m (60 ft) of cage row with each side being designated as a bank. Each bank is divided into six 3 m (10 ft) cage row sections consisting of four cages per section with a 60.9 cm (24 in) space between cage sections for feed hoppers and feed recovery. This system provided for 144 experimental units each consisting of four 60.9 x 50.8 cm (24 x 20 in) cages. Two densities were examined- 1) 125.3 cm² (69 in²), which allowed for 7 hens per cage for a total of 28 hens per replicate and 2) 304.8 cm² (120 in²), which allowed for 4 hens per cage for a total of 16 hens per replicate, resulting in a house total of 3,168 hens.

Table 2. Replicate numbers and Hen populations in the Enrichable Cage, Enriched Environmental Housing, and Conventional Cage System

House	Cage Style ¹	Number of Replicates	Hens per replicate	Hen No.	Total Hens
5	EC	104	36	3,744	
5	ECS	79	36	2,844	
5	ECS	76	18	1,368	7,956
7	C	62	28	1,736	
7	C	77	16	1,232	2,968

¹Cage=C; Enrichable Cage=EC; Enriched Environmental Housing System=ECS

FDA Egg Safety Plan Testing

In accordance with the Egg Safety Rule and the NCLP&MT Egg Safety Plan, the conventional cage, enrichable cage, and enriched environmental housing environments were tested between 40 and 44 weeks of age and post-molt between 77 and 79 weeks of age for the presence of *Salmonella enteritidis*. All of the environments at both time intervals were found to be negative for *Salmonella enteritidis*.

Lighting Schedule

The lighting schedule for the hens in controlled environment facilities are outlined in Table 3.

Table 3. Layer House Lighting¹ Schedules

Age	Date	Light Control Houses	
		5	7
		Photo Period ¹	
		(Daylight Hours)	(Daylight Hours)
16-17 weeks	Nov 19, 2013	10.0	10.0
17 Weeks	Nov. 27, 2013	11.0	11.0
18 Weeks	Dec. 4, 2013	11.5	11.5
19 Weeks	Dec. 11, 2013	12.0	12.0
20 Weeks	Dec. 18, 2013	12.5	12.5
21 Weeks	Dec. 24, 2013	13.0	13.0
22 Weeks	Jan. 1, 2014	13.5	13.5
23 Weeks	Jan. 8, 2014	14.0	14.0
24 Weeks	Jan. 15, 2014	14.25	14.25
25 Weeks	Jan. 22, 2014	14.5	14.5
26 Weeks	Jan. 29, 2014	14.75	14.75
27 Weeks	Feb. 5, 2014	15.0	15.0
28 Weeks	Feb. 12, 2014	15.25	15.25
29 Weeks	Feb. 19, 2014	15.5	15.5
30 Weeks	Feb. 26, 2014	15.75	15.75
31 Weeks	March 5, 2014	16.0	16.0
Molt Period			
Through 69 Weeks	Nov. 25, 2014	16.0	9.0
73 Weeks	Dec. 23, 2014	16.0	9.0
Post-Molt 2nd Cycle			
73 Weeks	Dec. 23, 2014	16.0	15.5
74 weeks	Dec. 30, 2014	16.0	16.0
74 Weeks through end of test (109 wk)	Dec. 30, 2014 to Sept. 1, 2015	16.0	16.0

¹Lighting intensity for Houses 5 and 7 was 0.5 to 0.7 ft candle at the second tier

Test Design:

The arrangement for the laying test involved a completely randomized design and the main effects were set up in a factorial arrangement. The main effects within Houses 5 and 7 were strain, density, and production system. Following are general descriptions of the main effects:

Strain: Strains were provided from the breeders according to the rules, which govern the conductance of the test. Fertile eggs were set and hatched concurrently as described in the hatch report (39th Hatch/Serology Report Vol. 39, No. 1). Additional chicks had to be acquired and delivered to the station to provide adequate bird numbers; the added chicks had hatch dates that were within two days of the original hatch.

Density - In Houses 5 and 7, all individual replicates within each section contained one strain of layers. The cage density in House 5 was dictated by the cage size 243.8 cm (96 in) or 60.9 cm (24 in) and populations of 9, 36, 18 hens per cage (Table 4). In House 7, all cages were 60.9 cm (24 in); density was dictated by either 7 or 5 hens per cage.

Table 4. Population and Density Allocations in Enrichable Cage, Enriched Environmental Housing, and Conventional Cage System

House	Hens per Cage	Cage Size Width Depth	Floor Space per Bird	Feeder Space per Bird	Water Nipples per Cage
5	36 ¹	243.8 cm x 66.0 cm	445.2 cm ² (69 in ²)	6.9 cm (2.7 in)	6
5	18 ²	243.8 cm x 66.0 cm	890.3 cm ² (138 in ²)	13.5 cm (5.3 in)	6
5	9	60.9 cm x 66.0 cm	445.2 cm ² (69 in ²)	6.9 cm (2.7 in)	2
7	7	60.9 cm x 50.8 cm	445.2 cm ² (69 in ²)	8.7 cm (3.4 in)	2
7	4	60.9 cm x 50.8 cm	774.2 cm ² (120 in ²)	15.2 cm (6.0 in)	2

¹Nest area was 51.6 cm²(20.3 in²)/hen, Scratch area 51.6 cm²(20.3 in²)/hen and the roost space was 6.8 cm (2.7 in)/hen

²Nest area was 103.2 cm²(40.6 in²)/hen, Scratch area 103.2 cm²(40.6 in²)/hen and the roost space was 13.5 cm (5.3 in)/hen

Layer Management (Molting):

The non-molted hens were in House 5. The full fed control replicates were maintained according to the standard layer management program. The molt was conducted utilizing all conventionally caged hens in House 7. The non-anorexic (NA) molt program hens were fed a low protein, low energy diet with supplemental Ca for maintenance.

Procedural steps:

Day (-)7	Sample of birds will be weighed to determine the pre-molt weight. Target weight loss (20% body weight) will be calculated using the pre-molt weight.
Day 0	NA program started with the remaining layer feed being removed and replaced with the NA molt diet and daylight hours reduced. Controlled light housing, reduce the day length to nine hours. Remove morbid birds <u>before</u> commencement of molt program.
Day +7	Body weights taken on two replicates from each strain and density in Houses 5 and 7.
Day +9	Body weights taken on two replicates from each strain and density in Houses 5 and 7. From the daily weight loss the day post initiation (Day 0) when the hens would be predicted to reach 20% weight loss. This is verified by body weight.
Day +28	Body weights were taken then the birds were fed layer diet and light stimulated.

The NA program was designed to keep hens out of production and provide balanced nutrition for body maintenance only. The diet is bulky, such that a full trailer load will only weigh two-thirds of a normal full load. The birds in the replicates being molted were weighted on days 7 and 9 to predict body weight loss. They were then weighed every other day until target weight was reached, at which time that replicate and sister replicates were provided the resting diet until the

end of the molting period. The induced molt was started at 69 weeks of age. The standard weight loss curve developed was utilized to manage the NA molt program.

Specific monitored criteria for all of the molt programs include the following: The birds were to attain approximately 20% body weight loss $\pm 3\%$. Maintain house temperature at $80 \pm 5^\circ \text{F}$, but the birds should not pant. House temperature management reacted to ambient environmental temperatures and weight loss rates.

The post-molt production period light schedule (Table 2) was the guide by which the lights were adjusted following the molt. Actual house conditions and the flock's reaction to the NCSU Non-Anorexic Molting Program may affect how the light stimulation was actually given. The hens ceased egg production by Day 6-10 of the molt program. However, some of the brown-egg strains never achieved zero egg production. The hens were allowed to consume all of the molt feed provided between feedings. The molting ration was designed to keep hens out of production, and to provide for skeletal and muscle maintenance. Livability was excellent with this program.

Layer Nutrition:

Laying hen diets are identified as Diets D, E, F, G, H, I, M, N, and O, which consisted of a pre-lay diet and a series of layer diets formulated to assure a daily protein, mineral, and amino acid intake (Table 5). Feed was offered ad libitum in accordance with the guidelines that all birds should receive acceptable nutrient intake at all times, depending on the bird's age and production rate, as shown in the Laying House Feeding Program (Table 6).

The diets provided during the molt consisted of a low protein/energy diet and a resting diet, described in the following Molt Diet Tables.

Table 5. Minimum Daily Intake of Nutrients per Bird at Various Stages of Production in the 39th NCLP&MT

Production Stage	Pre-Peak > 87%	87-80%	80-70%	<70%
White Egg Layers				
Protein ¹ (g/day)	19	18	17	16
Calcium (g/day)	4.0	4.1	4.2	4.3
Lysine (mg/day)	820	780	730	690
TSAA (mg)day)	700	670	630	590
Brown Egg Layers				
Protein ¹ (g/day)	20	19	18	17
Calcium (g/day)	4.0	4.0	4.1	4.2
Lysine (mg/day)	830	820	780	730
TSAA (mg)day)	710	700	670	630

¹ If the egg production is higher than predicted values, protein intake should be increased by 1%

Note: House temperatures dictate the body maintenance demand of the hen if the house temperature is 75 to 80°F feed protein content should be increased accordingly to compensate for metabolic heat needed to maintain a homeostatic body temperature. If the house temperature is at or above 85°F, no adjustment is needed.

Table 6 : NCLP&MT Laying House Feeding Program

Rate of Production	Consumption Per (kg/100 Birds/Day)	Diet Fed	
		White Egg Strains	Brown Egg Strains
Weeks 15-17	< 9.52	D	D
Pre-Peak and > 90%	< 9.52 - 10.43	D	E
	10.43 - 12.20	E	F
	12.25 ->13.11	F	G
90-80%	10.43 - 11.29	F	G
	11.34 - 12.20	G	H
	12.25 ->13.11	H	I
70-80%	10.43 - 11.29	H	I
	11.34 - 12.20	I	M
	12.25 ->13.11	M	N
< 70%	10.43 - 11.29	M	N
	11.34 - 12.20	N	O
	12.25 ->13.11	O	O

Note: Low house temperatures and egg production higher than breeder guides for any given hen age will require an adjustment to the dietary phase feeding program to ensure the hens are in a positive nutrient status.

Table 7. 39th NCLP&MT Laying Periods Feed Formulations¹ D through H

Ingredients	D	E	F	G	H
Corn	1054.24	1166.03	1202.7	1240.88	1285.39
Soybean meal	637.29	564.55	533.71	506.44	473.06
Fat (Lard)	10.00	10.00			15.68
D.L. Methionine	3.41	2.92	2.31	2.04	1.80
Soybean oil	61.8	25.90	36.29	25.06	
Ground Limestone	124.15	122.36	121.69	110.55	111.82
Coarse Limestone	70.00	70.00	70.00	75.00	75.00
Bi-Carbonate	2.00	2.00	2.00	3.00	2.00
Phosphate Mono/D	21.93	21.50	17.93	26.03	23.89
Salt	6.96	6.41	5.88	5.00	5.48
Vit. premix	1.00	1.00	1.00	1.00	1.00
Min. premix	1.00	1.00	1.00	1.00	1.00
HyD ₃ Broiler (62.5 mg/lb)			0.50		
Prop Acid 50% Dry	1.00	1.00	1.00	1.00	1.00
T-Premix	1.00	1.00	1.00	1.00	1.00
.06% Selenium Premix	1.00	1.00	1.00	1.00	1.00
Choline Cl 60%	1.62	1.94	1.59	1.00	0.87
Avizyme	1.00	1.00			
Ronozyme P-CT 540%	0.40	0.40	0.40		
Total	2000.00	2000.00	2000.00	2000.00	2000.00
Calculated Analysis					
Protein %	19.43	18.10	17.50	17.00	16.37
ME kcal/kg	2926.0	2904.0	2882	2860.0	2843.0
Calcium %	4.10	4.05	4.00	3.95	3.95
A. Phos. %	0.45	0.44	0.40	0.38	0.35
Lysine %	1.10	1.00	0.96	0.91	0.87
TSAA %	0.80	0.74	0.69	0.66	0.63

¹Feeds were manufactured by Southern States

Table 8. 39th NCLP&MT Laying Periods Feed Formulations I through O

Ingredients	I	M	N	O
Corn	1330.70	1315.29	1303.73	1290.76
Soybean meal	440.37	417.79	378.54	337.65
Wheat Midds		39.27	89.80	145.56
D.L. Methionine	1.56	1.24	1.14	0.78
Lysine 78.8%	2.23	0.10		
Ground Limestone	115.69	119.22	123.59	124.94
Coarse Limestone	75.00	75.00	75.00	75.00
Bi-Carbonate	2.00	2.00	2.00	2.00
Phosphate Mono/D	21.74	19.89	16.49	14.00
Salt	5.20	5.10	4.71	4.31
Vit. premix	1.00	1.00	1.00	1.00
Min. premix	1.00	1.00	1.00	1.00
Prop Acid 50% Dry	1.00	1.00	1.00	1.00
T-Premix	1.00	1.00	1.00	1.00
.06% Selenium Premix	1.00	1.00	1.00	1.00
Choline Cl 60%	0.52	0.10		
Total	2000.00	2000.00	2000.00	2000.00
Calculated Analysis				
Protein %	15.87	15.49	14.93	14.37
ME kcal/kg	2821.9	2800.0	2777.8	2755.8
Calcium %	4.00	4.05	4.10	4.10
A. Phos. %	0.33	0.31	.28	0.26
Lysine %	0.91	0.80	0.75	0.71
TSAA %	0.60	0.58	0.56	0.53

¹Feeds were manufactured by Southern States

Table 9. 39th NCLP&MT Laying Periods Feed Formulations Molt and Resting Diets

Ingredient	Molt Diets	
	<u>Low ME</u>	<u>Resting</u>
Corn	702.50	1427.70
Soybean Hulls	1164.77	226.00
Soybean Meal 48%		117.00
Wheat Midds	18.26	186.50
Coarse Limestone	17.78	16.50
Phosphate Mono/D	69.84	4.00
Salt	9.16	5.00
Methionine	2.69	1.30
Vit. premix	1.00	1.00
Min. premix	1.00	1.00
T - Premix	1.00	1.00
Fat	9.99	10.00
MYC-OUT 65	1.00	2.00
0.06% Sel Premix	1.00	1.00
Total	2000	2000
Calculated Analysis		
Protein %	9.92	11.75
Me kcal/kg	1650	2859
Calcium %	1.33	3.80
T. Phos %	0.88	0.44
Lysine %	0.42	0.55
TSAA %	0.35	0.49

Data Collection Schedule and Procedures:

Age at 50% Production (Maturity): The first day at which the birds in the individual replicates achieved 50% production.

Egg Production: All eggs that had the potential of being marketed were credited toward the test replicates egg production, regardless of the shell condition at the time of collection. All eggs were collected and recorded daily. Egg production was summarized at twenty-eight day intervals and was calculated and reported on a Hen-Housed and Hen-Day basis.

There were unexpected behaviors in the enriched environmental housing system related to broody behavior and laying of floor eggs, which may have influenced the overall performance of the hens in those replicates. In order to mitigate these behavioral issues, the frequency of egg collection was increased to minimize number of eggs the hens could interact with. There was egg breakage within these replicates; the broken egg numbers could not be captured.

Egg Weight: At twenty-eight day intervals, all eggs produced in the previous 24-hour period were weighed and sorted by size (see egg size distribution). Percentages of eggs within each size category, average egg weight (g), and egg mass (g) were calculated and reported and used to calculate egg income.

Egg Quality: At twenty-eight day intervals, all eggs produced within the previous 24-hours were examined by candling light and graded according to current USDA standards for egg quality. Eggs were graded in the pilot processing facility and handled as they would be in a commercial off-line facility. From 17 to 21 weeks of age (period 1), statistical estimates were made for those replicates where quality information was missing due to late onset of maturity from sister replicates. Broody behavior was displayed in many replicates in all of the housing environments, which resulted in hens sitting in the nests longer and attempting to pull eggs back into the nesting areas from egg trays.

Egg Size Distribution: At twenty-eight day intervals, all eggs produced within the previous 24 hours were weighed and sorted according to current USDA standards for egg size. From 17 to 21 weeks of age (period 1), statistical estimates were made for those replicates where size distributions were missing due to late onset of maturity from sister replicates.

Egg Income: Egg income was calculated using current production year calendar and applying a 3 year average egg price on egg production and quality evaluation.

Feed Consumption and Conversion: All feed offered for consumption was recorded for each replicate. At twenty-eight day intervals, feed not consumed was weighed back and feed consumption was calculated. Daily feed intake (kg per 100 hens per day) was calculated and reported for each strain.

The layer diets were reformulated to meet the flock nutrient needs based upon data from previous test reports. Based on the nutrient analysis conducted on each load of feed, the protein and Ca levels were in accordance with the calculated analysis. However, in the extensive environments where the hen activity levels were greater due to the availability of space to move within the environments, there appears to be a change in the partitioning of nutrients. Even though

the diets were formulated to meet the nutritional needs of the hens for optimal performance, there were pauses in the early stages of the production cycle indicating a need for additional nutrients in the diet. This was more pronounced in this test with the reduced daily feed consumption rates and increased bird floor space allowances for the environments to meet recommended density standards.

Feed Costs: Feed costs were based on the actual current feed prices for each feed delivery which were calculated and summarized for the complete production cycle.

Body weights: Birds were weighed and weights recorded at housing (17 weeks), end of 1st cycle (69 weeks), start of the 2nd cycle (73 weeks), and at the end of the 2nd cycle (109 weeks of age). Body weight gain for the 1st cycle was calculated and reported for each strain. In the Molt period, lowest body weight, percent weight loss, and 73 week body weights were taken or calculated and reported for each strain. The weight gain during the 2nd cycle was calculated along with the overall weight gain.

Mortality: All mortalities were recorded daily. Obvious accidents were not included in reported mortalities.

Statistical Analyses and Separation of Means:

All data were subjected to ANOVA utilizing the GLM procedure of JMP11 (SAS, 2014), with main effects of strain, density, and production system used herein. Period was accounted for in the model within each of the production systems. Separate analyses were conducted for white and brown-egg strains, the densities within production systems and between the enrichable and enriched colony housing system. Within each production system the Strain and Strain x Density/Housing System interactions were tested for significance. The LSMeans differences from the GLM Procedure were separated via the Tukey HSD option. Comparisons of overall production systems of Density or Housing System were tested for significance and their LS Means from the GLM Procedure were separated via the Student's t option. Significant differences ($P < 0.01$) within white and brown-egg strains are noted by differing letters among columns of means.

DESCRIPTION OF DATA TABLE STATISTICS

First cycle performance of white and brown-egg strains in the three production systems are reported from 119-483 days of age (17-69 weeks), the molt from 483-511 days of age (69-73 weeks), second cycle from 511-763 days of age (73-109 weeks), and 119-763 days of age (17-109 weeks) for comparative purposes. Conventional cage systems and densities are shown for the first cycle in Table 13 to 18, molt period performance Tables 19 to 24, second cycle in Tables 25 to 30, overall performance in Tables 31 to 36, and weight loss data summary of the white and brown egg strains are shown on Tables 37 and 38. The comparison of the Enrichable and Enriched Environmental Housing System are shown for the first cycle in Table 39 to 44, molt period performance Tables 45 to 50, second cycle in Tables 51 to 56, overall performance in Tables 57 to 62, and weight loss data summary of the white and brown egg strains are shown on Tables 63 and 64.

The comparison of the Enrichable and Enriched Environmental Housing System comparing den-

sities are shown for the first cycle in Table 65 to 70, molt period performance Tables 71 to 76, second cycle in Tables 77 to 82, overall performance in Tables 83 to 88, and weight loss data summary of the white and brown egg strains are shown on Tables 89 and 90.

Breeder (Strain): Short identification codes developed for strain and breeder of the stock are shown in Table 1 and 85.

Hen Housed Eggs per Bird: The total number of eggs produced divided by the number of birds housed at 119 days.

Hen Day Egg Production: The average daily number of eggs produced per 100 hens per day. Hen Day Production graphs for each strain, density and environment are shown in Figures 1 through 37.

Egg Mass: The average daily production of egg mass in grams per hen day.

Mortality: The percentage of birds which died between 119 through 483 days of age (1st Cycle) and 483 through 511 (Molt) which are reported separately. The hens in the Enrichable Cages and Enriched Environmental Housing System were not molted but the period 483 to 511 days are reported separately for comparative purposes.

Feed Consumption: The kilograms of feed consumed daily per 100 hens. Feed consumption graphs for each strain, density and environment are shown in Figures 1 through 37.

Feed Conversion: The grams of egg produced per gram of feed consumed.

Egg Weight: The average egg weight (g) for each period sampled. Weight of all eggs collected from previous 24-hours divided by the number of eggs collected.

Egg Income: The income from egg production per hen housed for the first cycle was calculated using the 3 year regional average egg prices November 27, 2011 to December 25, 2014. The second cycle egg prices used the September 3, 2012 to September 3, 2015 regional average egg prices.

Table 10. Three Year Regional Average Egg Prices

Grade	Size	\$\$/Dozen 1 st Cycle	\$\$/Dozen 2 nd Cycle
A	Extra Large	1.4445	1.6188
A	Large	1.4179	1.5869
A	Medium	1.1385	1.3004
A	Small	0.9408	1.0247
A ¹	Pee Wee	0.4612	0.5123
B ²	All	0.7367	0.8410

Table 10. Three Year Regional Average Egg Prices

Grade	Size	\$\$/Dozen 1 st Cycle	\$\$/Dozen 2 nd Cycle
Cracks ²	All	0.7367	0.8410

¹Prices are estimates based upon the formula provided by D.D. Bell (Small x 0.5)

²Prices are estimates based upon the formula provided by D.D. Bell (Large x 0.53)

The regional weighted average prices for small lots was obtained from the, USDA Grade A and Grade A, white eggs in cartons, from nearby retail outlets of eggs based in North Carolina (USDA-AMS, RA_PY001). The egg price used for the eggs from House 7 where the hens were molted were valued using the B grade price. The value of Pee Wee Eggs was based upon the formula of “Pee Wee value = (Price of Small eggs x 0.5) and under grade eggs (Grade B and Cracks) was based upon the formula “under grade value = (Price of Large eggs x 0.53). These calculated value formulas were provided by D. D. Bell, (1990), personal communications.

Grade Information: The average grade of all eggs sampled according to USDA grading standards over all sampling periods. Grades are established by personnel trained in USDA grading standards.

Egg Size Distribution: The following table shows the size classifications used for establishing the USDA egg size grading. There has been blending of egg size in this test with the weight cut-off between medium and large being 23.5. This maximizes the number of USDA large eggs just as would occur in a commercial plant. The proportion of the eggs falling into the following size categories are reported in the tables.

Table 11. USDA Egg Weights Used To Establish The Egg Size Distribution Weighted for Large Eggs.

Size Category	Ounces/Dozen
Pee Wee	< 18
Small	18 – 21
Medium	21 - 23.5
Large	23.5 – 27
Extra Large	> 27

Feed Cost: The calculated feed cost per hen housed at 119 days, using the kilogram/diet consumed and the average price of each diet per ton.

Table 12. The Average Contract Feed Price For Feed Purchases During The First Cycle.

<u>Diets</u>	Price Per Ton 1 st cycle	Price Per Ton 2 nd cycle
D	380.40	
E	380.34	353.13
F	363.29	340.55
G	342.90	327.71
H	326.60	320.15
I		317.64
M		309.25
N		297.00
O		278.90
Molt Diet LP/LE	277.00	
Resting	270.00	

Metric Conversions:

1 lb. = 453.6 g

1 lb. = .4536 kg

1 oz. = 28.35 g

1 g = .03527 oz.

1 kg = 2.204 lb.

1 g = 1000 mg

1 kg = 1000 g

TABLE 13. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
Bovans White	69	10.18 ^{efghi}	0.501 ^{bcdef}	310.55 ^{abcd}	85.33 ^{bcdefg}	51.79 ^{fghi}	2.70	144.25 ^{abcdef}
	120	11.02 ^{abcd}	0.490 ^{defg}	315.45 ^{efg}	86.64 ^{abcde}	53.80 ^{bcdef}	4.70	142.75 ^{abcdef}
	Average	10.60 ^{AB}	0.495 ^{BC}	313.00 ^{AB}	85.99 ^{CDE}	52.79 ^{DE}	3.70	143.50 ^{BCDE}
Shaver White	69	9.60 ^{ij}	0.543 ^a	316.70 ^{abc}	86.93 ^{abcd}	52.88 ^{efghi}	4.48	140.50 ^f
	120	10.10 ^{fghij}	0.539 ^{ab}	325.00 ^a	89.32 ^a	54.22 ^{bcde}	1.58	140.75 ^{ef}
	Average	9.83 ^{DE}	0.541 ^A	320.85 ^A	88.14 ^{AB}	53.55 ^{CD}	3.02	140.63 ^F
Dekalb White	69	10.33 ^{defghi}	0.503 ^{abcdef}	314.97 ^{abcd}	86.50 ^{abcde}	52.79 ^{efghi}	3.17	142.67 ^{abcdef}
	120	10.96 ^{abcde}	0.492 ^{defg}	321.12 ^{ab}	88.23 ^{ab}	54.79 ^{abcde}	3.12	141.75 ^{cdef}
	Average	10.64 ^{AB}	0.497 ^{BC}	318.04 ^A	87.37 ^{ABC}	53.79 ^{CD}	3.14	142.21 ^{CDEF}
Babcock White	69	9.98 ^{fghij}	0.546 ^a	320.60 ^{abc}	88.12 ^{-abc}	55.36 ^{abc}	1.20	141.67 ^{bcdef}
	120	11.28 ^{ab}	0.495 ^{cdefg}	324.72 ^a	89.22 ^a	56.45 ^a	1.58	140.75 ^{ef}
	Average	10.63 ^{AB}	0.521 ^{AB}	322.66 ^A	88.67 ^A	55.90 ^A	1.39	141.21 ^{EF}
ISA B-400	69	10.11 ^{efghij}	0.529 ^{abcd}	314.27 ^{abcd}	86.19 ^{abcdef}	53.89 ^{bcdef}	3.97	143.67 ^{abcdef}
	120	11.05 ^{abcd}	0.493 ^{defg}	314.90 ^{abcd}	86.45 ^{abcde}	54.70 ^{abcde}	1.58	144.50 ^{abcdef}
	Average	10.58 ^{AB}	0.511 ^{BC}	314.58 ^{AB}	86.32 ^{BCD}	54.30 ^{BC}	2.77	144.08 ^{ABCD}
Hy-Line W-36	69	9.28 ^j	0.525 ^{abcde}	296.37 ^{bcd}	81.29 ^{hij}	49.46 ^{kl}	3.20	146.33 ^{ab}
	120	10.72 ^{bcdef}	0.458 ^g	300.02 ^{abcd}	82.29 ^{ghij}	49.28 ^{kl}	2.08	146.25 ^a
	Average	10.00 ^{CDE}	0.492 ^C	298.19 ^{BC}	81.79 ^F	49.37 ^G	2.64	146.29 ^A
Hy-Line CV-26	69	9.24 ^j	0.502 ^{bcdefg}	288.27 ^d	79.07 ^j	46.88 ^m	3.57	145.00 ^{abcde}
	120	9.92 ^{ghij}	0.479 ^{fg}	293.70 ^{cd}	80.69 ^{ij}	47.72 ^{lm}	0	144.50 ^{abcdef}
	Average	9.58 ^E	0.490 ^C	290.98 ^C	79.88 ^F	47.30 ^H	1.78	144.75 ^{ABC}
Hy-Line CV-24	69	9.78 ^{hij}	0.512 ^{abcdef}	302.33 ^{abcd}	82.85 ^{fghi}	50.67 ^{ijk}	7.57	146.33 ^{ab}
	120	10.67 ^{bcdefg}	0.496 ^{cdefg}	316.92 ^{abc}	86.85 ^{abcd}	53.35 ^{cdefg}	9.40	142.25 ^{abcdef}
	Average	10.22 ^{BCD}	0.504 ^{BC}	309.63 ^{AB}	84.85 ^{DE}	52.01 ^{EF}	8.48	144.29 ^{ABCD}
Hy-Line CV-22	69	9.69 ^{ij}	0.520 ^{abcdef}	304.03 ^{abcd}	83.35 ^{efghi}	51.01 ^{hijk}	5.17	142.33 ^{abcdef}
	120	10.61 ^{bcdefgh}	0.484 ^{efg}	308.82 ^{abcd}	84.76 ^{defg}	51.45 ^{ghij}	4.70	141.25 ^{def}
	Average	10.15 ^{BCD}	0.502 ^{BC}	306.43 ^{ABC}	84.06 ^E	51.21 ^F	4.93	141.79 ^{DEF}
Lohmann LSL Lite	69	10.05 ^{fghij}	0.526 ^{abcde}	314.50 ^{abcd}	86.25 ^{abcdef}	52.90 ^{cdefghi}	5.93	144.67 ^{abcdef}
	120	11.22 ^{abc}	0.487 ^{defg}	319.85 ^{ab}	87.92 ^{abc}	55.07 ^{abcd}	1.58	145.75 ^{abc}
	Average	10.64 ^{AB}	0.507 ^{BC}	317.18 ^A	87.09 ^{ABC}	54.03 ^{BCD}	3.75	145.21 ^{AB}
H&N Nick Chick	69	9.85 ^{ghij}	0.536 ^{abc}	308.80 ^{abcd}	84.74 ^{cdefgh}	53.32 ^{cdefgh}	4.77	145.33 ^{abcd}
	120	11.15 ^{abcd}	0.497 ^{cdefg}	325.48 ^a	89.35 ^a	56.40 ^a	4.70	144.00 ^{abcdef}
	Average	10.50 ^{ABC}	0.517 ^{ABC}	317.14 ^A	87.04 ^{ABC}	54.86 ^{ABC}	4.73	144.67 ^{ABC}
Novogen White	69	10.40 ^{cdefghi}	0.520 ^{abcdef}	316.03 ^{abcd}	86.52 ^{abcde}	55.05 ^{abcde}	5.97	145.33 ^{abcd}
	120	11.58 ^a	0.482 ^{fg}	326.87 ^{ab}	88.69 ^a	55.75 ^{ab}	6.28	144.25 ^{abcdef}
	Average	10.99 ^A	0.501 ^{BC}	319.95 ^A	87.60 ^{ABC}	55.37 ^{AB}	6.12	144.79 ^{ABC}
All Strains	69	9.87	0.522 ^Z	308.95 ^Z	84.76 ^Z	52.17	4.30	144.00 ^Y
	120	10.85	0.491 ^Y	315.82 ^Y	86.70 ^Y	53.60	3.44	143.23 ^Z

¹All strains were housed such that each strain is equally represented in each density.

ABCDEF - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefghijkml - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 14. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans	69	59.04 ^{efgh}	0	8.18	9.57 ^{ab}	26.21 ^a	55.61 ^{ef}
White	120	61.06 ^{abcd}	0	4.96	8.07 ^{ab}	19.94 ^{abcd}	67.04 ^{abc}
	Average	60.05 ^{BCDE}	0	6.57	8.82 ^{ABC}	23.08	61.33 ^{BCDE}
Shaver	69	59.66 ^{cdefgh}	0	6.71	9.27 ^{ab}	26.09 ^{ab}	57.69 ^{def}
White	120	59.12 ^{defgh}	0	5.13	9.41 ^{ab}	25.17 ^{abc}	60.31 ^{bcde}
	Average	59.39 ^{EF}	0	5.92	9.34 ^{ABC}	25.63	59.00 ^{DE}
Dekalb	69	58.93 ^{defgh}	0	6.09	8.33 ^{ab}	24.03 ^{abcd}	61.46 ^{bcde}
White	120	60.53 ^{abcdefg}	0	5.45	6.46 ^{ab}	20.50 ^{abcd}	67.40 ^{abc}
	Average	59.73 ^{CDE}	0	5.77	7.40 ^{ABC}	22.26	64.43 ^{ABCD}
Babcock	69	61.38 ^{abc}	0	3.70	8.50 ^{ab}	20.48 ^{abcd}	67.08 ^{abcd}
White	120	61.96 ^{ab}	0.63	4.50	8.26 ^{ab}	14.56 ^d	71.83 ^a
	Average	61.67 ^A	0.32	4.10	8.38 ^{ABC}	17.52	69.45 ^A
ISA	69	60.78 ^{abcdefg}	0	7.20	5.19 ^{ab}	19.72 ^{abcd}	67.79 ^{abc}
B-400	120	61.42 ^{abc}	0	7.21	5.08 ^{ab}	15.73 ^{cd}	71.54 ^a
	Average	61.10 ^{ABC}	0	7.21	5.14 ^C	17.72	69.67 ^A
Hy-Line	69	59.95 ^{bcdefgh}	0	6.70	10.68 ^{ab}	22.79 ^{abcd}	59.28 ^{bcdef}
W-36	120	58.66 ^{fgh}	0	7.26	12.35 ^{ab}	24.06 ^{abcd}	55.77 ^{ef}
	Average	59.31 ^{EF}	0	6.98	11.51 ^{AB}	23.43	57.52 ^{EF}
Hy-Line	69	58.50 ^h	0	7.86	13.67 ^a	28.92 ^a	49.10 ^f
CV-26	120	58.12 ^h	0	8.36	11.72 ^{ab}	25.96 ^{ab}	54.02 ^{ef}
	Average	58.31 ^F	0	8.11	12.72 ^A	27.44	51.56 ^F
Hy-Line	69	59.56 ^{cdefgh}	0.84	5.33	9.52 ^{ab}	26.54 ^{ab}	57.61 ^{cdef}
CV-24	120	59.75 ^{cdefgh}	0	7.52	7.10 ^{ab}	25.13 ^{abc}	60.15 ^{bcde}
	Average	59.66 ^{DEF}	0.42	6.42	8.31 ^{ABC}	25.83	58.88 ^{DE}
Hy-Line	69	58.77 ^{cdefgh}	1.28	4.24	6.82 ^{ab}	27.28 ^a	60.26 ^{bcde}
CV-22	120	59.67 ^{cdefgh}	0	5.26	8.26 ^{ab}	25.62 ^{ab}	60.73 ^{bcde}
	Average	59.72 ^{CDEF}	0.64	4.75	7.54 ^{ABC}	26.45	60.49 ^{CDE}
Lohmann	69	60.78 ^{abcdefg}	0	5.23	8.19 ^{ab}	22.92 ^{abcd}	63.41 ^{abcde}
LSL Lite	120	60.81 ^{abcde}	0	4.02	7.98 ^{ab}	19.71 ^{abcd}	67.59 ^{abc}
	Average	60.69 ^{ABCD}	0	4.62	8.09 ^{ABC}	21.32	65.50 ^{ABC}
H&N	69	60.82 ^{abcdef}	0	5.23	7.94 ^{ab}	20.28 ^{abcd}	66.51 ^{abcd}
Nick Chick	120	61.53 ^{abc}	0.96	5.90	3.80 ^b	21.25 ^{abcd}	67.75 ^{ab}
	Average	61.18 ^{AB}	0.48	5.56	5.87 ^{BC}	20.77	67.13 ^{AB}
Novogen	69	62.35 ^a	0	7.29	4.39 ^{ab}	15.56 ^{bcd}	72.79 ^a
White	120	61.55 ^{abc}	0	6.71	5.07 ^{ab}	20.38 ^{abcd}	67.79 ^{ab}
	Average	61.95 ^A	0	7.00	4.73 ^C	17.97	70.29 ^A
All Strains	69	60.35	0.18	6.15	8.50	23.55 ^Y	61.55 ^Z
	120	60.13	0.13	6.02	7.80	21.50 ^Z	64.33 ^Y

¹All strains were housed such that each strain is equally represented in each density

ABCDEF - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefgh - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values

TABLE 15. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans	69	94.42	2.59	2.50 ^{ab}	0.48	35.00 ^{ab}	15.31 ^{bcdefg}
White	120	98.20	0.13	1.53 ^{ab}	0.15	35.93 ^{ab}	16.58 ^{abcde}
	Average	96.31	1.36	2.01 ^{AB}	0.32	35.46 ^{ABCD}	15.94 ^{AB}
Shaver	69	97.61	0	2.10 ^{ab}	0.29	34.79 ^{ab}	14.44 ^{fg}
White	120	96.14	0.15	3.71 ^{ab}	0	37.20 ^a	15.16 ^{bcdefg}
	Average	96.87	0.07	2.90 ^{AB}	0.15	35.99 ^{ABC}	14.80 ^{BC}
Dekalb	69	97.42	0	2.15 ^{ab}	0.43	35.62 ^{ab}	15.55 ^{abcdefg}
White	120	96.20	0.57	2.74 ^{ab}	0.49	36.96 ^a	16.50 ^{abcde}
	Average	96.81	0.28	2.45 ^{AB}	0.46	36.29 ^{AB}	16.03 ^{AB}
Babcock	69	96.87	0.28	2.43 ^{ab}	0.41	37.26 ^a	15.01 ^{bcdefg}
White	120	96.60	0.72	2.42 ^{ab}	0.27	37.44 ^a	16.98 ^{ab}
	Average	96.74	0.50	2.42 ^{AB}	0.34	37.3 ^A	15.99 ^{AB}
ISA	69	96.89	0.25	2.35 ^{ab}	0.51	35.58 ^{ab}	15.23 ^{bcdefg}
B-400	120	96.20	0.28	3.02 ^{ab}	0.49	36.00 ^{ab}	16.62 ^{abcde}
	Average	96.55	0.27	2.69 ^{AB}	0.50	35.79 ^{ABCD}	15.93 ^{AB}
Hy-Line	69	96.07	0	3.19 ^{ab}	0.74	33.10 ^{ab}	13.97 ^{fg}
W-36	120	95.31	0.32	3.77 ^{ab}	0.60	33.55 ^{ab}	16.12 ^{abcdef}
	Average	95.69	0.16	3.48 ^{AB}	0.67	33.32 ^{CD}	15.04 ^{BC}
Hy-Line	69	96.78	0.11	2.48 ^{ab}	0.63	32.45 ^b	13.91 ^g
CV-26	120	98.13	0.46	1.41 ^{ab}	0	33.82 ^{ab}	14.94 ^{cdefg}
	Average	97.46	0.28	1.94 ^{AB}	0.31	33.13 ^D	14.42 ^C
Hy-Line	69	97.52	0.64	1.59 ^{ab}	0.25	33.67 ^{ab}	14.70 ^{defg}
CV-24	120	98.42	0	1.46 ^{ab}	0.12	35.31 ^{ab}	16.06 ^{abcdefg}
	Average	97.97	0.32	1.52 ^B	0.18	34.44 ^{BCD}	15.38 ^{ABC}
Hy-Line	69	95.70	0.14	4.04 ^{ab}	0.12	34.43 ^{ab}	14.58 ^{efg}
CV-22	120	95.25	0	4.60 ^a	0.15	35.53 ^{ab}	15.98 ^{abcdefg}
	Average	95.47	0.07	4.32 ^A	0.13	34.98 ^{ABCD}	15.28 ^{ABC}
Lohmann	69	97.64	0.24	1.92 ^{ab}	0.20	35.46 ^{ab}	15.13 ^{bcdefg}
LSL Lite	120	96.03	0	3.12 ^{ab}	0.85	36.70 ^{ab}	16.90 ^{abc}
	Average	96.84	0.12	2.52 ^{AB}	0.52	36.08 ^{ABC}	16.01 ^{AB}
H&N	69	97.10	0.46	2.31 ^{ab}	0.13	35.40 ^{ab}	14.82 ^{cdefg}
Nick Chick	120	97.25	0.27	1.98 ^{ab}	0.50	37.34 ^a	16.79 ^{abcd}
	Average	97.17	0.36	2.14 ^{AB}	0.32	36.37 ^{AB}	15.80 ^{ABC}
Novogen	69	98.96	0.27	0.77 ^b	0	35.51 ^{ab}	15.64 ^{abcdefg}
White	120	98.08	0.30	1.62 ^{ab}	0	36.78 ^{ab}	17.43 ^a
	Average	98.52	0.28	1.20 ^B	0	36.15 ^{ABC}	16.53 ^A
All	69	96.92	0.42	2.32	0.35	34.85 ^Z	14.86 ^Z
Strains	120	96.82	0.27	2.61	0.30	36.05 ^Y	16.34 ^Y

¹All strains were housed such that each strain is equally represented in each density.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 16. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
TETRA	69	10.48 ^{bcde}	0.466 ^{defg}	301.27 ^{cdef}	82.61 ^{gh}	49.20 ^{ef}	4.76	145.33 ^{ab}
Amber	120	11.09 ^{abcd}	0.460 ^{fg}	316.82 ^{abcd}	86.90 ^{bcd}	51.06 ^{de}	6.25	142.75 ^{abc}
	Average	10.79 ^{AB}	0.463 ^C	309.04	84.76 ^{ABC}	50.13 ^B	5.51	144.04 ^{AB}
TETRA	69	10.32 ^{bcdef}	0.474 ^{bcdefg}	303.15 ^{bcdef}	82.43 ^{fgh}	49.63 ^{ef}	17.86	142.50 ^{abc}
Brown	120	11.30 ^{abc}	0.455 ^g	310.15 ^{abcdef}	85.19 ^{cdefg}	50.94 ^{def}	3.12	140.00 ^{bc}
	Average	10.81 ^{AB}	0.464 ^C	303.65	83.81 ^{BC}	50.28 ^B	10.49	141.25 ^{ABC}
Novogen	69	10.06 ^{ef}	0.508 ^{abc}	305.27 ^{bcdef}	83.62 ^{efgh}	51.46 ^{cde}	7.14	143.00 ^{abc}
Brown	120	11.10 ^{abcd}	0.480 ^{bcdefg}	312.12 ^{abcdef}	85.72 ^{cdef}	53.59 ^{bc}	1.56	141.00 ^{bc}
	Average	10.58 ^{AB}	0.494 ^B	308.69	84.67 ^{ABC}	52.53 ^A	4.35	142.00 ^{ABC}
Lohmann	69	9.81 ^{ef}	0.501 ^{abcd}	297.03 ^{ef}	81.53 ^h	49.54 ^{ef}	3.57	142.00 ^{abc}
LB-Lite	120	11.16 ^{ab}	0.494 ^{abcdef}	326.72 ^a	89.78 ^a	55.32 ^{ab}	4.69	141.25 ^{bc}
	Average	10.48 ^{ABC}	0.497 ^{AB}	311.88	85.66 ^{AB}	52.43 ^A	4.13	141.62 ^{BC}
Hy-Line	69	10.09 ^{ef}	0.479 ^{bcdefg}	304.17 ^{cdef}	83.57 ^{efgh}	48.73 ^f	1.19	143.33 ^{abc}
Silver Brown	120	11.46 ^a	0.458 ^g	322.98 ^{ab}	88.63 ^{ab}	52.59 ^{cd}	3.12	139.00 ^c
	Average	10.77 ^{AB}	0.468 ^C	313.57	86.10 ^A	50.65 ^B	2.16	141.67 ^{BC}
Hy-Line	69	9.53 ^f	0.506 ^{abc}	294.77 ^f	81.00 ^h	48.56 ^f	1.19	140.00 ^{bc}
Brown	120	10.40 ^{cde}	0.506 ^{abc}	313.27 ^{abcde}	86.07 ^{bcde}	52.35 ^{cd}	1.56	139.50 ^c
	Average	9.96 ^C	0.506 ^{AB}	304.02	83.54 ^C	50.46 ^B	1.38	139.75 ^C
ISA	69	9.61 ^{ef}	0.529 ^a	302.63 ^{cdef}	82.99 ^{fgh}	51.48 ^{cde}	1.76	147.00 ^a
Brown	120	10.94 ^{abcd}	0.508 ^{ab}	323.10 ^{ab}	88.78 ^{ab}	55.76 ^a	3.15	143.00 ^{abc}
	Average	10.27 ^{BC}	0.519 ^A	312.87	85.88 ^{AB}	53.62 ^A	3.94	145.00 ^A
Bovans	69	10.36 ^{de}	0.497 ^{abcde}	308.17 ^{bcdef}	84.57 ^{defg}	52.33 ^{cd}	6.25	142.25 ^{abc}
Brown	120	11.59 ^a	0.471 ^{cdefg}	319.48 ^{abc}	87.74 ^{abc}	54.16 ^{ab}	3.12	143.50 ^{abc}
	Average	10.97 ^A	0.484 ^{BC}	313.82	86.15 ^A	53.74 ^A	4.69	142.87 ^{ABC}
All Strains	69	10.03 ^Z	0.495 ^Y	302.06 ^Z	82.79 ^Z	50.12 ^Z	5.84	143.18 ^Y
	120	11.13 ^Y	0.479 ^Z	318.08 ^Y	87.32 ^Y	53.35 ^Y	3.32	141.25 ^Z

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefgh - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 17. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN CONVENTIONAL CAGES

Breeder (Strain)	Density ¹ (in ² /hen)	Egg Weight (g/egg)	Pee Wee (%)	Small (%)	Medium (%)	Large (%)	Extra Large (%)
TETRA Amber	69	58.08 ^{ef}	0	5.00	14.56 ^{ab}	30.49 ^{abcde}	49.74 ^{def}
	120	57.34 ^f	0	4.60	13.12 ^{ab}	34.19 ^{ab}	47.56 ^{ef}
	Average	57.71 ^E	0	4.80	13.84 ^{AB}	32.34 ^{AB}	48.65 ^{CD}
TETRA Brown	69	59.48 ^{abcdef}	0	4.81	9.62 ^{abc}	29.54 ^{abcde}	55.73 ^{bcd}
	120	58.94 ^{cdef}	0	3.85	8.88 ^{abc}	32.19 ^{abcd}	54.85 ^{bcd}
	Average	59.21 ^{CD}	0	4.33	9.25 ^{BC}	30.86 ^{AB}	55.29 ^{BC}
Novogen Brown	69	60.56 ^{abc}	0.74	3.15	11.13 ^{abc}	25.13 ^{bcd}	59.72 ^{abcd}
	120	61.40 ^a	0	2.88	6.87 ^{bc}	20.33 ^e	69.83 ^a
	Average	60.98 ^A	0.37	3.02	9.00 ^{BC}	22.73 ^C	64.77 ^A
Lohmann LB-Lite	69	60.17 ^{abcd}	0	5.10	11.29 ^{abc}	27.44 ^{abcde}	55.99 ^{bcd}
	120	60.24 ^{abd}	2.88	3.67	4.85 ^c	24.27 ^{cde}	63.88 ^{ab}
	Average	60.20 ^{ABC}	1.44	4.39	8.07 ^C	25.85 ^{BC}	59.93 ^{AB}
Hy-Line Silver Brown	69	57.54 ^f	0	4.26	16.77 ^a	36.13 ^a	42.72 ^f
	120	58.62 ^{def}	0	2.90	13.67 ^{ab}	32.40 ^{abc}	50.69 ^{def}
	Average	58.08 ^{DE}	0	3.58	15.22 ^A	34.27 ^A	46.70 ^D
Hy-Line Brown	69	59.02 ^{cdef}	0	2.18	10.36 ^{abc}	35.00 ^{ab}	52.18 ^{cdef}
	120	60.14 ^{abcd}	0	1.98	8.62 ^{bc}	26.86 ^{abcde}	62.52 ^{abc}
	Average	59.58 ^{BC}	0	2.08	9.49 ^{BC}	30.93 ^{AB}	57.35 ^B
ISA Brown	69	60.03 ^{abcde}	2.56	3.85	6.69 ^{bc}	24.31 ^{bcd}	62.54 ^{abc}
	120	61.40 ^a	0	4.96	5.71 ^c	20.75 ^{de}	68.50 ^a
	Average	60.72 ^{AB}	1.28	4.40	6.20 ^C	22.53 ^C	65.52 ^A
Bovans Brown	69	60.64 ^{abc}	0	3.69	11.52 ^{abc}	21.67 ^{de}	63.02 ^{ab}
	120	61.03 ^{ab}	0	3.02	7.04 ^{bc}	20.52 ^{de}	68.46 ^a
	Average	60.84 ^A	0	3.36	9.28 ^{BC}	21.60 ^C	65.74 ^A
All Strains	69	59.89	0.41	4.00	11.49 ^Y	28.71	55.20 ^Z
	120	59.44	0.36	3.48	8.59 ^Z	26.56	60.79 ^Y

¹All strains were housed such that each strain is equally represented in each density.

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 18. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69	96.92	0.30	2.26	0.53	33.88 ^{cd}	15.55 ^{abcde}
	120	97.82	0.14	2.04	0	35.19 ^{abc}	16.47 ^{abc}
	Average	97.37	0.22	2.15	0.26	34.54 ^{AB}	16.01 ^A
TETRA Brown	69	96.07	0.46	2.88	0.59	31.36 ^d	15.32 ^{cde}
	120	97.02	0.59	2.13	0.26	34.34 ^{abc}	16.76 ^{abc}
	Average	96.55	0.52	2.50	0.42	33.35 ^B	16.04 ^A
Novogen Brown	69	97.07	0.71	1.89	0.34	33.67 ^{cd}	14.93 ^{cde}
	120	98.14	0.12	1.47	0.26	36.53 ^{abc}	16.48 ^{abc}
	Average	97.60	0.41	1.68	0.30	35.10 ^{AB}	15.70 ^{AB}
Lohmann LB-Lite	69	97.33	0	2.34	0.32	33.66 ^{cd}	14.54 ^{de}
	120	96.97	0.14	2.12	0.77	37.70 ^a	16.57 ^{abc}
	Average	97.15	0.07	2.23	0.54	35.68 ^A	15.56 ^{AB}
Hy-Line Silver Brown	69	97.47	0.23	2.09	0.20	34.73 ^{abcd}	15.00 ^{cde}
	120	97.39	0.27	2.20	0.13	36.68 ^{abc}	17.00 ^{ab}
	Average	97.43	0.25	2.15	0.16	35.70 ^A	15.99 ^A
Hy-Line Brown	69	96.84	0.21	2.62	0.32	33.87 ^{cd}	14.14 ^e
	120	95.28	0.54	3.87	0.30	35.89 ^{abc}	15.45 ^{bde}
	Average	96.07	0.38	3.24	0.31	34.88 ^{AB}	14.79 ^B
ISA Brown	69	96.41	0.85	2.54	0.19	35.00 ^{abc}	14.24 ^e
	120	97.24	0.42	2.22	0.13	37.36 ^{ab}	16.24 ^{abcd}
	Average	96.82	0.63	2.38	0.16	36.18 ^A	15.24 ^{AB}
Bovans Brown	69	97.62	0.43	1.77	0.18	34.77 ^{abcd}	15.36 ^{cde}
	120	96.51	0.62	2.87	0	37.39 ^{ab}	17.20 ^a
	Average	97.03	0.52	2.32	0.09	36.08 ^A	16.28 ^A
All Strains	69	96.97	0.40	2.30	0.33	33.87 ^Z	14.88 ^Z
	120	97.05	0.36	2.36	0.23	36.51 ^Y	16.52 ^Y

¹All strains were housed such that each strain is equally represented in each density.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 19. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder (Strain)	Density ¹ (in ² /hen)	Feed Consumption (kg/100/hen/d)	Feed Conversion (g egg/g feed)	Eggs		Egg Mass (g/HD)	Mortality (%)
				Per Bird Housed	Egg Production (HD%)		
Bovans White	69	5.16	0.06	3.78 ^{abc}	13.42 ^{abcd}	3.24	0.90
	120	5.94	0.04	4.38 ^{abc}	15.62 ^{abcd}	2.29	0
	Average	5.55 ^{AB}	0.05	4.08 ^{AB}	14.52 ^{AB}	2.76	0.45
Shaver White	69	4.55	0	3.88 ^{abc}	13.84 ^{abcd}	0	0
	120	3.99	0	4.50 ^{ab}	15.95 ^{abc}	0	1.58
	Average	4.27 ^{AB}	0	4.19 ^A	14.89 ^A	0	0.79
Dekalb White	69	5.76	0.13	3.90 ^{abc}	13.87 ^{abcd}	7.34	2.80
	120	5.68	0.05	4.68 ^a	16.74 ^a	2.52	0
	Average	5.72 ^A	0.09	4.29 ^A	15.30 ^A	4.93	1.40
Babcock White	69	4.78	0.04	3.17 ^{abc}	11.33 ^{abcd}	2.08	0
	120	4.75	0	3.20 ^{abc}	11.31 ^{abcd}	0	1.58
	Average	4.76 ^{AB}	0.02	3.18 ^{ABC}	11.32 ^{ABC}	1.04	0.79
ISA B-400	69	4.95	0	3.13 ^{abc}	11.01 ^{abcd}	0	2.40
	120	4.79	0.03	2.82 ^{abc}	9.82 ^{cd}	1.66	1.58
	Average	4.87 ^{AB}	0.02	2.98 ^{ABC}	10.42 ^{BC}	0.83	1.99
Hy-Line W-36	69	63.93	0.05	2.57 ^{bc}	9.05 ^{cd}	1.86	0
	120	3.94	0	2.85 ^{abc}	10.08 ^{abcd}	0	0
	Average	3.94 ^B	0.02	2.71 ^C	9.57 ^C	0.93	0
Hy-Line CV-26	69	4.5	0	2.43 ^c	8.65 ^d	0	1.20
	120	3.92	0.04	3.60 ^{abc}	12.82 ^{abcd}	2.01	0
	Average	4.25 ^{AB}	0.02	3.17 ^{ABC}	10.73 ^{ABC}	1.00	0.60
Hy-Line CV-24	69	4.86	0.05	3.40 ^{abc}	12.01 ^{abcd}	2.61	1.20
	120	4.67	0.08	4.65 ^a	16.50 ^{ab}	4.53	1.58
	Average	4.76 ^{AB}	0.07	4.02 ^{ABC}	14.26 ^{AB}	3.57	1.39
Hy-Line CV-22	69	5.08	0.09	3.00 ^{abc}	10.75 ^{abcd}	4.28	0
	120	4.05	0.06	2.80 ^{abc}	10.03 ^{bcd}	2.32	0
	Average	4.57 ^{AB}	0.07	2.90 ^{BC}	10.39 ^{BC}	3.30	0
Lohmann LSL Lite	69	5.28	0	3.63 ^{abc}	12.87 ^{abcd}	0	1.20
	120	4.31	0.04	4.05 ^{abc}	14.34 ^{abcd}	2.43	1.58
	Average	4.80 ^{AB}	0.02	3.84 ^{ABC}	13.61 ^{ABC}	1.22	1.39
H&N Nick Chick	69	5.14	0	3.13 ^{abc}	11.61 ^{abcd}	0	0
	120	4.99	0	4.22 ^{abc}	15.08 ^{abcd}	0	0
	Average	5.07 ^{AB}	0	3.68 ^{ABC}	13.12 ^{ABC}	0	0
Novogen White	69	5.08	0.05	3.60 ^{abc}	12.62 ^{abcd}	2.83	3.57
	120	4.65	0.04	3.75 ^{abc}	13.43 ^{abcd}	2.54	0
	Average	4.87 ^{AB}	0.05	3.68 ^{ABC}	13.02 ^{ABC}	2.68	1.78
All Strains	69	4.93	0.04	3.30 ^Z	11.72 ^Z	2.02	1.10
	120	4.64	0.03	3.79 ^Y	13.48 ^Y	1.69	0.66

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 20. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans	69	50.00	0	12.50	12.50	25.00 ^{ab}	0
White	120	70.00	0	0	0	0 ^b	25.00
	Average	60.00	0	6.25	6.25	12.50 ^{AB}	12.50
Shaver	69	59.70	0	0	0	0 ^b	0
White	120	60.00	0	0	0	0 ^b	0
	Average	59.80	0	0	0	0 ^B	0
Dekalb	69	53.30	0	0	0	100.00 ^a	0
White	120	55.00	0	0	0	25.00 ^{ab}	0
	Average	54.60	0	0	0	62.50 ^A	0
Babcock	69	50.00	0	16.67	0	16.67 ^{AB}	0
White	120	61.30	0	0	0	0 ^b	0
	Average	55.60	0	8.33	0	8.33 ^{AB}	0
ISA	69	60.80	0	0	0	0 ^b	0
B-400	120	50.00	0	0	0	25.00 ^{ab}	0
	Average	55.40	0	0	0	12.50 ^{AB}	0
Hy-Line	69	62.50	0	0	0	16.67 ^{ab}	16.67
W-36	120	59.00	0	0	0	0 ^b	0
	Average	60.70	0	0	0	8.33 ^{AB}	8.33
Hy-Line	69	59.00	0	0	0	0 ^b	0
CV-26	120	60.00	0	0	0	25.00 ^{ab}	0
	Average	59.50	0	0	0	12.50 ^{AB}	0
Hy-Line	69	60.00	0	0	0	0 ^b	33.33
CV-24	120	50.00	0	25.00	0	0 ^b	25.00
	Average	55.00	0	12.50	0	0 ^B	29.17
Hy-Line	69	55.00	0	0	0	66.67 ^{ab}	0
CV-22	120	70.00	0	0	0	0 ^b	25.00
	Average	62.50	0	0	0	33.33 ^{AB}	12.50
Lohmann	69	59.60	0	0	0	0 ^b	0
LSL Lite	120	60.00	0	0	0	25.00 ^{ab}	0
	Average	59.80	0	0	0	12.50 ^{AB}	0
H&N	69	60.40	0	0	0	0 ^b	0
Nick Chick	120	60.70	0	0	0	0 ^b	0
	Average	60.30	0	0	0	0 ^B	0
Novogen	69	60.00	0	0	0	0 ^b	33.33
White	120	60.00	0	0	0	25.00 ^{ab}	0
	Average	60.00	0	0	0	12.50 ^{AB}	16.17
All Strains	69	57.50	0	2.43	1.04	18.75	6.94
	120	59.60	0	2.08	0	10.42	6.25

¹All strains were housed such that each strain is equally represented in each density

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions

TABLE 21. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans	69	50.00	0	0	0	0.65 ^{ab}	0.43
White	120	25.00	0	0	0	0.75 ^{ab}	0.50
	Average	37.50	0	0	0	0.70 ^{AB}	0.47 ^{AB}
Shaver	69	0	0	0	0	0.67 ^{ab}	0.38
White	120	0	0	0	0	0.77 ^{ab}	0.33
	Average	0	0	0	0	0.72 ^{AB}	0.36 ^{AB}
Dekalb	69	66.67	0	0	0	0.67 ^{ab}	0.48
White	120	25.00	0	0	0	0.81 ^a	0.48
	Average	45.83	0	0	0	0.74 ^A	0.48 ^A
Babcock	69	33.33	0	0	0	0.54 ^{ab}	0.40
White	120	0	0	0	0	0.55 ^{ab}	0.40
	Average	16.17	0	0	0	0.55 ^{ABC}	0.40 ^{AB}
ISA	69	0	0	0	0	0.54 ^{ab}	0.41
B-400	120	25.00	0	0	0	0.48 ^{ab}	0.40
	Average	12.50	0	0	0	0.51 ^{BC}	0.41 ^{AB}
Hy-Line	69	33.33	0	0	0	0.44 ^b	0.33
W-36	120	0	0	0	0	0.46 ^{ab}	0.33
	Average	16.17	0	0	0	0.46 ^C	0.33 ^B
Hy-Line	69	0	0	0	0	0.42 ^b	0.38
CV-26	120	25.00	0	0	0	0.62 ^{ab}	0.33
	Average	12.50	0	0	0	0.52 ^{ABC}	0.36 ^{AB}
Hy-Line	69	33.33	0	0	0	0.58 ^{ab}	0.41
CV-24	120	50.00	0	0	0	0.80 ^a	0.5239
	Average	41.67	0	0	0	0.69 ^{AB}	0.40 ^{AB}
Hy-Line	69	66.67	0	0	0	0.52 ^{ab}	0.43
CV-22	120	25.00	0	0	0	0.48 ^{ab}	0.34
	Average	45.83	0	0	0	0.50 ^{BC}	0.38 ^{AB}
Lohmann	69	0	0	0	0	0.62 ^{ab}	0.44
LSL Lite	120	25.00	0	0	0	0.69 ^{ab}	0.36
	Average	12.50	0	0	0	0.66 ^{ABC}	0.40 ^{AB}
H&N	69	0	0	0	0	0.54 ^{ab}	0.43
Nick Chick	120	0	0	0	0	0.73 ^{ab}	0.42
	Average	0	0	0	0	0.63 ^{ABC}	0.42 ^{AB}
Novogen	69	33.33	0	0	0	0.62 ^{ab}	0.42
White	120	25.00	0	0	0	0.65 ^{ab}	0.39
	Average	29.17	0	0	0	0.63 ^{ABC}	0.41 ^{AB}
All	69	26.39	0	2.77	0	0.57 ^Z	0.41
Strains	120	18.75	0	0	0	0.65 ^Y	0.39

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 22. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA	69	5.77 ^{ab}	0.130	5.00 ^{abcd}	17.92 ^{abcde}	7.49	0
Amber	120	5.60 ^{ab}	0.265	7.55 ^a	26.93 ^a	14.80	0
	Average	5.69 ^{AB}	0.197	6.27 ^A	22.42 ^A	11.14	0
TETRA	69	5.67 ^{ab}	0	4.30 ^{cd}	15.38 ^{bcde}	0	0
Brown	120	6.40 ^a	0.145	6.10 ^{abcd}	21.91 ^{abcde}	8.51	0
	Average	6.04 ^A	0.073	5.20 ^{ABC}	18.65 ^{ABC}	4.25	0
Novogen	69	5.87 ^{ab}	0.227	6.27 ^{abcd}	22.35 ^{abcd}	13.29	0
Brown	120	5.31 ^{ab}	0.257	6.58 ^{abc}	22.95 ^{abc}	13.36	3.15
	Average	5.59 ^{AB}	0.242	6.42 ^A	22.65 ^{AB}	13.32	1.58
Lohmann	69	4.52 ^b	0.110	3.70 ^d	13.20 ^{de}	5.02	3.57
LB-Lite	120	5.39 ^{ab}	0.280	7.18 ^{ab}	25.64 ^{ab}	14.98	0
	Average	4.95 ^B	0.195	5.44 ^{ABC}	19.42 ^{AB}	10.00	1.78
Hy-Line	69	5.52 ^{ab}	0.177	5.20 ^{abcd}	18.56 ^{abcde}	9.73	0
Silver Brown	120	5.08 ^{ab}	0.117	5.80 ^{abcd}	20.71 ^{abcde}	6.16	0
	Average	5.30 ^{AB}	0.147	5.50 ^{AB}	19.63 ^{AB}	7.96	0
Hy-Line	69	5.06 ^{ab}	0.127	4.30 ^{cd}	15.37 ^{cde}	6.64	0
Brown	120	4.74 ^b	0.085	3.52 ^d	12.59 ^e	4.08	0
	Average	4.90 ^B	0.106	3.91 ^{BC}	13.98 ^{BC}	5.38	0
ISA	69	4.74 ^b	0.110	3.70 ^d	13.14 ^{de}	4.86	0
Brown	120	5.44 ^{ab}	0.175	5.90 ^{abcd}	21.12 ^{abcde}	9.45	0
	Average	5.09 ^{AB}	0.142	4.80 ^{ABC}	17.31 ^{ABC}	7.16	0
Bovans	69	5.58 ^{ab}	0.060	3.70 ^d	13.20 ^{de}	3.28	0.90
Brown	120	4.80 ^{ab}	0.097	3.85 ^d	13.78 ^{de}	4.86	0
	Average	5.19 ^{AB}	0.079	3.78 ^C	13.49 ^C	4.07	0.45
All Strains	69	5.34	0.118	4.52 ^Z	16.14 ^Z	6.29	0.56
	120	5.35	0.178	5.81 ^Y	20.70 ^Y	9.52	0.39

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*density interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 23. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA	69	56.20	0	8.33	0	58.33	0
Amber	120	55.00	0	0	29.25	33.50	37.50
	Average	55.60	0	4.17	14.63	45.92	18.75
TETRA	69	59.10	0	0	0	0	0
Brown	120	65.00	0	0	0	25.00	25.00
	Average	62.00	0	0	0	12.50	12.50
Novogen	69	59.30	0	0	16.67	44.33	39.00
Brown	120	58.00	0	0	0	77.25	22.75
	Average	58.60	0	0	8.33	60.79	30.88
Lohmann	69	55.80	0	0	11.00	44.33	11.00
LB-Lite	120	58.30	0	0	8.25	75.00	16.75
	Average	57.00	0	0	9.63	59.67	13.88
Hy-Line	69	52.50	0	8.33	33.33	58.33	0
Silver Brown	120	60.00	0	0	0	41.75	8.25
	Average	56.20	0	4.17	16.67	50.04	4.12
Hy-Line	69	61.70	0	0	0	50.00	16.67
Brown	120	50.00	0	0	0	25.00	25.00
	Average	55.80	0	0	0	37.50	20.83
ISA	69	57.50	0	0	0	33.33	33.33
Brown	120	57.80	0	0	25.00	33.25	16.75
	Average	57.60	0	0	12.50	33.29	25.04
Bovans	69	55.00	0	0	0	50.00	0
Brown	120	60.00	0	0	0	20.75	29.25
	Average	57.50	0	0	0	35.38	14.62
All	69	57.30	0	2.08	7.62	42.33	12.50
Strains	120	58.00	0	0	7.81	41.44	22.66

¹All strains were housed such that each strain is equally represented in each density.

TABLE 24. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69	66.67	0	0	0	0.64	0.48 ^{ab}
	120	100.00	0	0	0	0.85	0.47 ^{ab}
	Average	83.33	0	0	0	0.74	0.48 ^{AB}
TETRA Brown	69	0	0	0	0	0	0.48 ^{ab}
	120	50.00	0	0	0	0.44	0.54 ^a
	Average	25.00	0	0	0	0.22	0.51 ^A
Novogen Brown	69	88.89	11.11	0	0	0.64	0.49 ^{ab}
	120	83.33	16.67	0	0	0.64	0.45 ^{ab}
	Average	86.11	13.89	0	0	0.64	0.47 ^{AB}
Lohmann LB-Lite	69	66.67	0	0	0	0.29	0.38 ^b
	120	100.00	0	0	0	0.84	0.45 ^{ab}
	Average	83.33	0	0	0	0.57	0.42 ^B
Hy-Line Silver Brown	69	83.33	16.67	0	0	0.45	0.46 ^{ab}
	120	50.00	0	0	0	0.34	0.43 ^{ab}
	Average	66.67	8.33	0	0	0.39	0.44 ^{AB}
Hy-Line Brown	69	66.67	0	0	0	0.35	0.43 ^{ab}
	120	25.00	25.00	0	0	0.14	0.40 ^b
	Average	45.83	12.50	0	0	0.25	0.41 ^B
ISA Brown	69	66.67	0	0	0	0.28	0.40 ^b
	120	75.00	0	0	0	0.51	0.46 ^{ab}
	Average	70.83	0	0	0	0.40	0.43 ^{AB}
Bovans Brown	69	50.00	0	0	0	0.26	0.47 ^{ab}
	120	37.50	12.50	0	0	0.28	0.40 ^{ab}
	Average	43.75	6.25	0	0	0.27	0.44 ^{AB}
All Strains	69	61.11	3.47	0	0	0.36	0.45
	120	65.10	6.77	0	0	0.50	0.45

¹All strains were housed such that each strain is equally represented in each density.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

TABLE 25. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans White	69	106.69 ^{abcd}	0.51	208.02	82.84 ^{abc}	54.60 ^{abc}	6.25
	120	110.37 ^{abc}	0.51	207.38	82.77 ^{abc}	56.06 ^{abc}	7.81
	Average	108.53 ^{ABC}	0.51	207.70 ^A	82.81 ^A	55.33 ^A	7.03
Shaver White	69	102.62 ^{bcde}	0.53	203.42	81.79 ^{abc}	53.71 ^{abc}	5.36
	120	103.89 ^{bcde}	0.54	211.72	85.04 ^{ab}	56.05 ^{abc}	3.12
	Average	103.26 ^{CDE}	0.54	207.58 ^{AB}	83.41 ^A	54.88 ^A	4.24
Dekalb White	69	106.76 ^{abcd}	0.53	215.10	85.44 ^{ab}	56.15 ^{abc}	6.35
	120	110.03 ^{abc}	0.48	200.20	80.49 ^{abc}	53.56 ^{abc}	3.12
	Average	108.90 ^{ABC}	0.51	207.65 ^{AB}	82.97 ^A	54.86 ^A	4.74
Babcock White	69	107.96 ^{abc}	0.52	204.93	82.40 ^{abc}	55.41 ^{abc}	3.57
	120	112.38 ^{ab}	0.48	199.65	80.38 ^{abc}	54.30 ^{abc}	6.25
	Average	110.17 ^{ABC}	0.50	202.29 ^{ABC}	81.39 ^{ABC}	54.85 ^A	4.91
ISA B-400	69	102.76 ^{bcde}	0.54	207.83	83.21 ^{abc}	54.34 ^{abc}	5.16
	120	105.88 ^{abcd}	0.50	199.52	80.47 ^{abc}	53.01 ^{abc}	1.56
	Average	104.32 ^{BCDE}	0.52	203.68 ^{ABC}	81.84 ^{AB}	53.68 ^{AB}	3.36
Hy-Line W-36	69	93.85 ^e	0.52	184.53	74.41 ^{bc}	49.14 ^{bc}	6.35
	120	105.46 ^{abcd}	0.48	186.82	75.16 ^{bc}	49.82 ^{bc}	8.33
	Average	99.65 ^{DE}	0.50	185.68 ^{BC}	74.79 ^{BC}	49.48 ^{BC}	7.34
Hy-Line CV-26	69	95.33 ^{de}	0.50	183.23	74.01 ^c	47.71 ^c	2.38
	120	100.19 ^{cde}	0.49	184.00	74.17 ^c	48.21 ^c	3.12
	Average	97.76 ^E	0.50	183.67 ^C	74.09 ^C	47.96 ^C	2.75
Hy-Line CV-24	69	103.75 ^{bcde}	0.54	208.47	82.82 ^{abc}	55.42 ^{abc}	5.95
	120	107.48 ^{abc}	0.53	213.58	84.78 ^{abc}	56.65 ^{ab}	9.38
	Average	105.62 ^{ABCD}	0.54	211.02 ^A	83.80 ^A	56.04 ^A	7.66
Hy-Line CV-22	69	104.63 ^{abcde}	0.51	198.93	79.79 ^{abc}	53.01 ^{abc}	1.19
	120	106.30 ^{abcd}	0.52	204.85	82.08 ^{abc}	54.88 ^{abc}	0
	Average	105.47 ^{ABCD}	0.52	201.89 ^{ABC}	80.93 ^{ABC}	53.94 ^{AB}	0.60
Lohmann LSL Lite	69	109.74 ^{abc}	0.52	213.67	85.28 ^{ab}	57.32 ^{ab}	5.95
	120	115.18 ^a	0.50	212.85	85.05 ^{ab}	58.07 ^a	6.25
	Average	112.46 ^A	0.51	213.26 ^A	85.16 ^A	57.69 ^A	6.10
H&N Nick Chick	69	104.81 ^{abcde}	0.54	209.00	83.35 ^{abc}	56.08 ^{abc}	5.95
	120	111.63 ^{ab}	0.53	217.42	86.86 ^a	59.07 ^a	3.12
	Average	108.22 ^{ABC}	0.54	213.21 ^A	85.11 ^A	57.58 ^A	4.54
Novogen White	69	111.24 ^{abc}	0.54	214.87	85.51 ^{ab}	59.14 ^a	1.19
	120	110.18 ^{abc}	0.54	215.65	86.49 ^a	57.95 ^a	1.56
	Average	110.71 ^{AB}	0.54	215.26 ^A	86.00 ^A	58.54 ^A	1.38
All Strains	69	104.18 ^Y	0.52	204.33	81.74	54.33	4.63
	120	108.33 ^Z	0.51	204.48	81.98	54.80	4.47

¹All strains were housed such that each strain is equally represented in each density.

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 26. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans	69	65.97 ^{bcdefg}	0	0	0.75	9.60	89.05
White	120	67.80 ^{abcd}	0	0	0.73	4.15	94.50
	Average	66.89 ^{ABC}	0	0	0.74	6.87 ^{AB}	91.78 ^{AB}
Shaver	69	65.66 ^{defg}	0	0	0.73	7.88	90.85
White	120	65.82 ^{cdefg}	0	0	0.55	7.32	92.12
	Average	65.74 ^{CDE}	0	0	0.64	7.60 ^{AB}	91.49 ^{AB}
Dekalb	69	65.75 ^{cdefg}	0	0	0.20	6.53	93.00
White	120	66.48 ^{bcdefg}	0	0.15	0	3.88	95.75
	Average	66.11 ^{BCDE}	0	0.08	0.10	5.20 ^{AB}	94.38 ^{AB}
Babcock	69	67.13 ^{abcdefg}	0	0	0.27	7.23	92.07
White	120	67.51 ^{abcde}	0	0	0	5.52	94.08
	Average	67.32 ^{AB}	0	0	0.13	6.38 ^{AB}	93.07 ^{AB}
ISA	69	65.13 ^{efg}	0	0	0	11.07	88.07
B-400	120	65.45 ^{efg}	0	0	0.62	12.58	86.58
	Average	65.29 ^{DE}	0	0	0.31	11.82 ^A	87.32 ^B
Hy-Line	69	66.04 ^{bcdefg}	0	0	0.70	7.67	91.20
W-36	120	66.36 ^{bcdefg}	0	0	0.68	7.25	91.50
	Average	66.20 ^{BCDE}	0	0	0.69	7.46 ^{AB}	91.35 ^{AB}
Hy-Line	69	64.50 ^g	0	0	0.20	9.40	89.90
CV-26	120	64.97 ^{fg}	0	0	0.42	8.52	90.25
	Average	64.74 ^E	0	0	0.31	8.96 ^{AB}	90.08 ^{AB}
Hy-Line	69	66.87 ^{abcdefg}	0	0	0.17	7.47	91.77
CV-24	120	66.84 ^{abcdefg}	0	0	0.20	7.50	92.18
	Average	66.86 ^{ABCD}	0	0	0.18	7.48 ^{AB}	91.97 ^{AB}
Hy-Line	69	66.48 ^{abcdefg}	0	0	0	5.87	93.80
CV-22	120	67.00 ^{abcdefg}	0	0	0.18	3.08	96.10
	Average	66.74 ^{ABCD}	0	0	0.09	4.47 ^B	94.95 ^A
Lohmann	69	67.18 ^{abcdefg}	0	0	0.17	5.97	93.37
LSL Lite	120	68.24 ^{ab}	0	0	0.25	7.62	92.18
	Average	67.71 ^{AB}	0	0	0.21	6.80 ^{AB}	92.77 ^{AB}
H&N	69	67.22 ^{abcdef}	0	0	0	5.90	93.17
Nick Chick	120	68.06 ^{abc}	0	0	0	4.30	95.58
	Average	67.64 ^{AB}	0	0	0	5.10 ^{AB}	94.37 ^{AB}
Novogen	69	69.12 ^a	0	0	0	2.43	97.37
White	120	66.92 ^{abcdefg}	0	0	1.08	7.10	91.82
	Average	68.02 ^A	0	0	0.54	4.77 ^{AB}	94.60 ^{AB}
All	69	66.42	0	0	0.26	7.25	91.97
Strains	120	66.79	0	0.01	0.39	6.57	92.72

¹All strains were housed such that each strain is equally represented in each density

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain*density interactions.

TABLE 27. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans	69	95.35	0.22	4.82	0.62	25.53 ^{ab}	10.22 ^{abc}
White	120	95.97	0.50	2.87	0.87	25.23 ^{ab}	10.52 ^{abc}
	Average	95.16	0.36	3.84	0.74	25.38 ^A	10.37 ^{AB}
Shaver	69	95.31	0.36	3.76	0.58	24.84 ^{ab}	9.74 ^{abc}
White	120	96.47	0	3.52	0	25.94 ^{ab}	9.96 ^{abc}
	Average	95.89	0.18	3.64	0.29	25.39 ^A	9.85 ^{ABC}
Dekalb	69	96.57	0.14	2.91	0.62	26.21 ^{ab}	10.23 ^{abc}
White	120	96.54	0.68	2.50	0.28	24.46 ^{ab}	10.57 ^{ab}
	Average	96.56	0.41	2.70	0.45	25.34 ^{AB}	10.40 ^{AB}
Babcock	69	96.70	0	2.86	0.44	24.94 ^{ab}	10.33 ^{abc}
White	120	95.93	0.80	2.78	0.48	24.33 ^{ab}	10.74 ^{ab}
	Average	96.32	0.40	2.82	0.46	24.63 ^{ABC}	10.53 ^{AB}
ISA	69	95.72	0.59	3.01	0.70	25.23 ^{ab}	9.73 ^{abc}
B-400	120	96.29	0.36	3.12	0.22	24.34 ^{ab}	9.89 ^{abc}
	Average	96.00	0.47	3.07	0.46	24.78 ^{ABC}	9.85 ^{ABC}
Hy-Line	69	96.80	0	2.78	0.61	22.28 ^{ab}	8.97 ^c
W-36	120	96.30	0.28	2.75	0.67	22.63 ^{ab}	10.11 ^{abc}
	Average	96.55	0.14	2.77	0.64	22.46 ^{BC}	9.54 ^{BC}
Hy-Line	69	95.48	0.30	3.64	0.89	22.32 ^{ab}	9.17 ^{bc}
CV-26	120	97.08	0.21	1.90	0.82	22.16 ^b	9.60 ^{abc}
	Average	96.28	0.25	2.77	0.85	22.24 ^C	9.38 ^C
Hy-Line	69	96.08	0.54	2.78	0.59	25.38 ^{ab}	10.03 ^{abc}
CV-24	120	96.57	0	3.24	0.19	26.10 ^{ab}	10.26 ^{abc}
	Average	96.33	0.27	3.01	0.39	25.74 ^A	10.14 ^{ABC}
Hy-Line	69	95.90	0.20	3.60	0.31	24.32 ^{ab}	10.05 ^{abc}
CV-22	120	94.04	0.18	5.40	0.38	25.17 ^{ab}	10.26 ^{abc}
	Average	94.97	0.19	4.50	0.35	24.74 ^{ABC}	10.16 ^{ABC}
Lohmann	69	95.06	0.29	4.17	0.49	26.17 ^{ab}	10.52 ^{abc}
LSL Lite	120	95.43	0	4.56	0.21	26.17 ^{ab}	10.96 ^a
	Average	95.25	0.14	4.36	0.35	26.17 ^A	10.74 ^A
H&N	69	96.44	0.37	2.22	0.97	25.23 ^{ab}	9.97 ^{abc}
Nick Chick	120	96.41	0.57	2.84	0.18	26.56 ^a	10.71 ^{ab}
	Average	96.42	0.47	2.53	0.57	25.89 ^A	10.34 ^{ABC}
Novogen	69	97.22	0	2.58	0.21	26.20 ^{ab}	10.66 ^{ab}
White	120	95.64	0.60	3.76	0	26.54 ^a	10.38 ^{abc}
	Average	96.43	0.30	3.17	0.10	26.37 ^A	10.52 ^{AB}
All Strains	69	95.97	0.25	3.26	0.58	24.89	9.97 ^Y
	120	96.06	0.35	3.27	0.36	24.97	10.34 ^Z

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 28. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA	69	108.30	0.42 ^{bc}	178.87	70.10 ^{ab}	45.55 ^{bc}	8.33
Amber	120	106.54	0.41 ^c	176.72	69.15 ^b	43.88 ^c	9.38
	Average	107.42	0.42 ^D	177.80	69.62 ^B	44.72 ^B	8.85
TETRA	69	106.56	0.45 ^{abc}	186.25	73.39 ^{ab}	47.03 ^{abc}	1.78
Brown	120	110.79	0.41 ^{bc}	179.60	70.46 ^{ab}	46.18 ^{abc}	6.25
	Average	108.68	0.43 ^{BCD}	182.92	71.92 ^{AB}	46.61 ^{AB}	4.01
Novogen	69	100.52	0.46 ^{abc}	175.63	68.69 ^b	46.14 ^{abc}	8.33
Brown	120	104.12	0.47 ^{abc}	185.62	72.81 ^{ab}	48.57 ^{abc}	6.25
	Average	102.32	0.46 ^{ABC}	180.63	70.75 ^B	47.36 ^{AB}	7.29
Lohmann	69	102.60	0.49 ^{ab}	191.83	75.44 ^{ab}	49.92 ^{abc}	5.95
LB-Lite	120	106.63	0.50 ^a	202.50	79.45 ^a	52.74 ^a	4.69
	Average	104.61	0.49 ^A	197.17	77.45 ^A	51.33 ^A	5.32
Hy-Line	69	106.00	0.44 ^{abc}	179.98	75.55 ^{ab}	46.61 ^{abc}	2.38
Silver Brown	120	109.82	0.41 ^c	192.00	70.78 ^{ab}	45.29 ^{bc}	3.12
	Average	107.91	0.43 ^{CD}	185.99	73.16 ^{AB}	45.95 ^B	2.75
Hy-Line	69	102.17	0.45 ^{abc}	176.57	69.32 ^b	45.62 ^{bc}	5.95
Brown	120	104.97	0.47 ^{abc}	190.40	74.62 ^{ab}	48.92 ^{abc}	9.38
	Average	103.57	0.46 ^{ABCD}	183.48	71.98 ^{AB}	47.27 ^{AB}	7.66
ISA	69	100.03	0.50 ^a	185.10	73.02 ^{ab}	49.62 ^{abc}	3.57
Brown	120	107.67	0.48 ^{ab}	196.52	77.23 ^{ab}	51.65 ^{ab}	3.12
	Average	103.85	0.49 ^A	190.81	75.12 ^{AB}	50.63 ^A	3.35
Bovans	69	106.62	0.46 ^{abc}	186.08	73.19 ^{ab}	48.53 ^{abc}	4.46
Brown	120	112.03	0.48 ^{ab}	195.35	77.07 ^{ab}	52.67 ^a	0
	Average	109.32	0.47 ^{AB}	190.71	75.13 ^{AB}	50.60 ^A	2.23
All	69	104.10 ^Y	0.46	184.04	72.34	47.38	5.10
Strains	120	107.82 ^Z	0.45	188.34	73.94	48.74	5.27

¹All strains were housed such that each strain is equally represented in each density.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 29. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69	65.00 ^{cde}	0	0	0.59	13.44 ^{ab}	85.89 ^{ab}
	120	63.49 ^{ef}	0	0	0.67	11.14 ^{ab}	87.94 ^{ab}
	Average	64.24 ^{CD}	0	0	0.63	12.29 ^{AB}	86.92 ^{AB}
TETRA Brown	69	64.07 ^{cdef}	0	0	0	13.67 ^{ab}	85.83 ^{ab}
	120	65.54 ^{abcde}	0	0.72	0.72	7.56 ^{ab}	90.44 ^{ab}
	Average	64.80 ^{CD}	0	0.36	0.36	10.61 ^{AB}	88.14 ^{AB}
Novogen Brown	69	67.17 ^{abc}	0	0	0	9.89 ^{ab}	88.41 ^{ab}
	120	66.68 ^{abc}	0	0	0	7.36 ^{ab}	91.36 ^{ab}
	Average	66.92 ^A	0	0	0	8.62 ^{AB}	89.88 ^{AB}
Lohmann LB-Lite	69	66.06 ^{abcde}	0	0	0.41	11.30 ^{ab}	88.15 ^{ab}
	120	66.43 ^{abcd}	0	0	0.39	7.11 ^{ab}	92.53 ^{ab}
	Average	66.25 ^{AB}	0	0	0.40	9.20 ^{AB}	90.34 ^{AB}
Hy-Line Silver Brown	69	61.82 ^f	0	0	0.22	16.11 ^a	83.56 ^b
	120	64.01 ^{def}	0	0	0.19	10.89 ^{ab}	87.03 ^{ab}
	Average	62.91 ^D	0	0	0.21	13.50 ^A	85.29 ^B
Hy-Line Brown	69	65.77 ^{abcde}	0	0	0.59	10.37 ^{ab}	88.59 ^{ab}
	120	65.51 ^{bcde}	0	0	0.50	8.50 ^{ab}	90.53 ^{ab}
	Average	65.64 ^{ABC}	0	0	0.55	9.44 ^{AB}	89.56 ^{AB}
ISA Brown	69	67.96 ^{ab}	0	0	0.37	10.18 ^{ab}	89.44 ^{ab}
	120	66.80 ^{abc}	0	0	0.28	10.39 ^{ab}	88.11 ^{ab}
	Average	67.38 ^A	0	0	0.32	10.29 ^{AB}	88.78 ^{AB}
Bovans Brown	69	66.27 ^{abcd}	0	0	0.53	8.56 ^{ab}	90.94 ^{ab}
	120	68.20 ^a	0	0.20	0	5.44 ^b	94.14 ^a
	Average	67.23 ^A	0	0.10	0.26	7.00 ^B	92.54 ^A
All Strains	69	65.51	0	0	0.34	11.69 ^Z	87.60 ^Z
	120	65.83	0	0.11	0.34	8.55 ^Y	90.26 ^Y

¹All strains were housed such that each strain is equally represented in each density.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdef - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 30. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69	96.27	0	3.56	0.17	21.80	10.22 ^{ab}
	120	96.47	0	3.22	0.31	21.44	10.06 ^{ab}
	Average	96.37	0	3.39	0.24	21.63	10.14
TETRA Brown	69	96.81	0	2.63	0.56	22.56	10.08 ^{ab}
	120	96.46	0.43	1.73	1.39	21.63	10.46 ^{ab}
	Average	96.63	0.21	2.18	0.97	22.09	10.28
Novogen Brown	69	95.18	0.22	2.94	1.67	21.13	9.53 ^{ab}
	120	96.48	0	2.11	1.41	22.26	9.84 ^{ab}
	Average	95.83	0.11	2.52	1.53	21.69	9.68
Lohmann LB-Lite	69	97.41	0	2.41	0.18	23.27	9.69 ^{ab}
	120	98.27	0	1.73	0	24.55	10.06 ^{ab}
	Average	97.84	0	2.07	0.09	23.91	9.88
Hy-Line Silver Brown	69	97.22	0	2.60	0.18	23.29	10.01 ^{ab}
	120	95.58	0	2.52	1.90	21.53	10.38 ^{ab}
	Average	96.40	0	2.56	1.04	22.41	10.20
Hy-Line Brown	69	96.64	0.40	2.46	0.51	21.40	9.67 ^{ab}
	120	95.44	0.28	3.80	0.48	23.21	9.92 ^{ab}
	Average	96.04	0.33	3.13	0.50	22.30	9.79
ISA Brown	69	97.00	0.85	2.16	0	22.58	9.45 ^b
	120	94.98	1.02	2.76	1.24	23.74	10.18 ^{ab}
	Average	95.99	0.93	2.46	0.62	23.16	9.82
Bovans Brown	69	96.49	0.32	3.19	0	22.73	10.08 ^{ab}
	120	96.36	0.20	2.96	0.47	23.79	10.60 ^a
	Average	96.42	0.26	3.08	0.24	23.26	10.34
All Strains	69	96.63	0.22	2.74	0.40	22.34	9.84 ^Y
	120	96.25	0.24	2.60	0.90	22.77	10.19 ^Z

¹All strains were housed such that each strain is equally represented in each density.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 31. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans White	69	101.49 ^{abcdef}	0.49	550.32	80.75	50.67	9.82
	120	107.82 ^{abc}	0.48	543.18	81.51	52.61	12.50
	Average	104.64 ^A	0.48	546.75 ^A	81.13	51.44 ^{ABC}	11.16
Shaver White	69	95.98 ^{def}	0.51	552.00	81.04	50.55	9.82
	120	99.34 ^{cdef}	0.51	557.22	83.85	52.36	6.25
	Average	97.66 ^{BC}	0.51	554.61 ^A	82.44	51.45 ^{ABC}	8.03
Dekalb White	69	102.66 ^{abcde}	0.49	559.60	82.55	52.02	12.30
	120	107.58 ^{abc}	0.47	542.00	81.44	51.72	6.25
	Average	105.12 ^A	0.48	550.80 ^A	81.99	51.87 ^{ABC}	9.28
Babcock White	69	100.66 ^{abcdef}	0.51	556.73	81.92	52.75	4.76
	120	109.72 ^{ab}	0.47	543.62	81.65	52.77 ^{AB}	9.38
	Average	105.19 ^A	0.49	550.18 ^A	81.79	52.76	7.07
ISA B-400	69	99.00 ^{cdef}	0.51	550.87	81.23	51.44	11.51
	120	105.31 ^{abcd}	0.48	533.25	80.13	51.33	4.69
	Average	102.15 ^{AB}	0.49	542.06 ^A	80.68	51.39 ^{ABC}	8.10
Hy-Line W-36	69	90.83 ^f	0.50	504.40	74.76	46.92	9.52
	120	103.64 ^{abcd}	0.44	501.70	75.70	47.05	10.42
	Average	97.24 ^{BC}	0.47	503.05 ^B	75.23	46.98 ^{BC}	9.97
Hy-Line CV-26	69	91.72 ^{ef}	0.47	501.93	73.36	44.86	7.14
	120	97.00 ^{def}	0.46	496.38	74.53	45.68	3.12
	Average	94.36 ^C	0.47	499.15 ^B	73.94	45.27 ^C	5.13
Hy-Line CV-24	69	98.27 ^{cdef}	0.50	539.80	79.37	50.37	14.68
	120	104.28 ^{abcd}	0.49	551.15	82.52	52.35	20.31
	Average	101.27 ^{ABC}	0.50	545.48 ^A	80.95	51.36 ^{ABC}	17.50
Hy-Line CV-22	69	98.00 ^{cdef}	0.49	531.67	78.25	49.52	6.35
	120	103.60 ^{abcd}	0.48	532.48	79.97	50.54	4.69
	Average	100.80 ^{ABC}	0.48	532.07 ^{AB}	79.11	50.03 ^{ABC}	5.52
Lohmann LSL Lite	69	102.08 ^{abcdef}	0.50	559.80	82.21	52.23	13.10
	120	110.11 ^a	0.47	552.48	83.03	53.79	9.38
	Average	106.10 ^A	0.49	556.14 ^A	82.62	53.01 ^{AB}	11.24
H&N Nick Chick	69	98.67 ^{cdef}	0.51	548.90	80.53	51.87	10.72
	120	108.92 ^{abc}	0.49	563.15	84.66	54.80	7.81
	Average	103.80 ^{AB}	0.50	556.02 ^A	82.60	53.33 ^{AB}	9.26
Novogen White	69	104.45 ^{abcd}	0.51	562.47	82.59	54.30	10.71
	120	109.87 ^{ab}	0.49	559.32	84.02	54.07	7.81
	Average	107.16 ^A	0.50	560.90 ^A	83.34	54.18 ^A	9.26
All Strains	69	98.65 ^Y	0.50	543.21	79.88	50.62	10.03
	120	105.60 ^Z	0.48	539.66	81.08	51.56	8.55

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdef - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 32. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69	60.50	0	5.17	5.94	18.78	67.42
	120	60.62	0	0.09	4.71	12.67	76.33
	Average	60.56	0	2.63	5.03	15.73	71.88
Shaver White	69	59.53	0	3.79	5.44	17.63	68.41
	120	59.34	0	2.90	5.48	16.75	70.54
	Average	59.44	0	3.35	5.46	17.19	69.48
Dekalb White	69	61.50	0	3.45	4.71	20.29	71.38
	120	60.94	0	3.14	3.65	14.03	75.70
	Average	61.22	0	3.30	4.18	17.16	73.54
Babcock White	69	61.79	0	2.81	4.83	15.10	74.06
	120	61.54	0.36	2.54	4.65	10.11	77.66
	Average	61.67	0.18	2.68	4.74	12.60	75.86
ISA B-400	69	58.80	0	1.17	2.94	14.87	73.35
	120	60.47	0	3.26	2.87	15.01	74.16
	Average	59.63	0	2.22	2.90	14.94	73.76
Hy-Line W-36	69	60.10	0	2.35	6.16	16.32	70.33
	120	58.29	0	1.92	7.12	16.73	68.13
	Average	59.20	0	2.14	6.64	16.02	69.23
Hy-Line CV-26	69	57.78	0	3.00	7.75	18.64	64.38
	120	58.64	0	3.64	6.62	18.01	67.06
	Average	58.21	0	3.32	7.19	18.32	65.72
Hy-Line CV-24	69	60.91	0	3.01	5.40	17.00	70.88
	120	61.17	0.24	5.34	4.01	16.62	71.74
	Average	61.04	0.72	4.18	4.71	16.81	71.31
Hy-Line CV-22	69	60.88	0	1.68	3.13	20.22	71.14
	120	60.93	0.36	2.97	4.65	14.90	73.84
	Average	60.91	0	2.32	3.89	17.56	72.49
Lohmann LSL Lite	69	60.87	0	2.96	4.65	14.75	72.96
	120	61.01	0	1.18	3.54	14.82	74.66
	Average	60.94	0	2.07	4.10	14.78	73.81
H&N Nick Chick	69	60.78	0	2.96	4.48	13.51	74.28
	120	61.06	0.54	2.79	1.63	13.58	75.80
	Average	60.92	0.27	2.88	3.05	13.54	75.04
Novogen White	69	62.69	0	2.67	2.49	9.68	80.75
	120	60.78	0	1.62	3.34	15.04	74.54
	Average	61.73	0	2.14	2.91	12.36	77.65
All Strains	69	60.51	0.10	2.92	4.83	16.40	71.61
	120	60.40	0.08	2.62	4.31	14.77	73.35

¹All strains were housed such that each strain is equally represented in each density

TABLE 33. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans	69	89.00	1.56	3.46 ^{ab}	0.54	60.72 ^{ab}	25.90 ^{abcd}
White	120	89.70	0.29	2.11 ^{ab}	0.38	61.28 ^{ab}	27.25 ^{abc}
	Average	89.35	0.92	2.79	0.46	61.00 ^A	26.58 ^{AB}
Shaver	69	87.97	0.16	2.76 ^{ab}	0	59.63 ^{ab}	25.09 ^{cd}
White	120	88.77	0.08	3.54 ^{ab}	0.42	63.13 ^a	26.08 ^{abcd}
	Average	88.37	0.12	3.15	0.21	61.38 ^A	25.59 ^{ABC}
Dekalb	69	92.80	0.06	3.93 ^{ab}	0.41	62.16 ^{ab}	26.33 ^{abcd}
White	120	88.74	0.62	2.63 ^{ab}	0.40	61.58 ^{ab}	27.10 ^{abcd}
	Average	90.77	0.34	3.28	0.40	61.87 ^A	27.72 ^{AB}
Babcock	69	92.74	0.16	2.62 ^{ab}	0.43	62.32 ^{ab}	26.17 ^{abcd}
White	120	88.95	0.76	2.51 ^{ab}	0.36	61.77 ^{ab}	27.58 ^{abc}
	Average	90.84	0.46	2.56	0.39	62.04 ^A	26.80 ^{AB}
ISA	69	87.68	0.40	2.64 ^{ab}	0.59	60.80 ^{ab}	25.74 ^{abcd}
B-400	120	88.73	0.32	2.97 ^{ab}	0.37	60.46 ^{ab}	26.32 ^{abcd}
	Average	88.20	0.36	2.80	0.48	60.63 ^A	26.03 ^{ABC}
Hy-Line	69	89.48	0	2.75 ^{ab}	0.60	55.48 ^b	24.40 ^d
W-36	120	87.28	0.30	3.09 ^{ab}	0.63	56.18 ^b	26.30 ^{abcd}
	Average	88.38	0.15	2.92	0.62	55.83 ^{BC}	25.35 ^{BC}
Hy-Line	69	89.22	0.13	2.92 ^{ab}	0.62	54.78 ^b	24.29 ^d
CV-26	120	91.36	0.26	1.50 ^b	0.36	56.10 ^b	25.15 ^{bcd}
	Average	90.29	0.19	2.21	0.49	55.43 ^C	24.72 ^C
Hy-Line	69	94.08	0.59	2.03 ^{ab}	0.40	59.10 ^{ab}	25.61 ^{abcd}
CV-24	120	91.10	0	2.23 ^{ab}	0.15	61.66 ^{ab}	26.19 ^{abcd}
	Average	92.59	0.30	2.13	0.27	60.38 ^{AB}	25.90 ^{ABC}
Hy-Line	69	90.07	0.16	3.77 ^{ab}	0.20	59.00 ^{ab}	25.78 ^{abcd}
CV-22	120	89.62	0.08	4.62 ^a	0.25	60.81 ^{ab}	26.48 ^{abcd}
	Average	89.84	0.12	4.20	0.22	59.90 ^{ABC}	26.13 ^{ABC}
Lohmann	69	92.30	0.19	2.83 ^{ab}	0.32	61.62 ^{ab}	26.90 ^{abcd}
LSL Lite	120	88.45	0	3.54 ^{ab}	0.49	63.01 ^a	27.72 ^{ab}
	Average	90.38	0.09	3.19	0.40	62.32 ^A	27.20 ^A
H&N	69	88.18	0.42	2.21 ^{ab}	0.49	60.63 ^{ab}	25.37 ^{abcd}
Nick Chick	120	89.50	0.31	2.28 ^{ab}	0.37	64.00 ^a	27.80 ^a
	Average	88.84	0.37	2.24	0.43	62.26 ^A	26.58 ^{AB}
Novogen	69	90.96	0.15	1.55 ^{ab}	0.09	61.87 ^{ab}	26.32 ^{abcd}
White	120	91.58	0.43	2.55 ^{ab}	0	63.46 ^a	26.94 ^{abcd}
	Average	91.27	0.29	2.05	0.04	62.66 ^A	26.63 ^{AB}
All	69	90.37	0.33	2.79	0.43	59.84 ^Z	25.23 ^Y
Strains	120	89.48	0.29	2.80	0.31	61.11 ^Y	26.74 ^Z

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 34. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA	69	104.13 ^{abcdef}	0.43	513.18	74.90	45.96	13.10
Amber	120	106.80 ^{abcd}	0.43	517.08	77.35	46.67	15.62
	Average	105.47 ^{AB}	0.43 ^B	515.12	76.13	46.32	14.36
TETRA	69	102.50 ^{abcdef}	0.44	521.20	75.98	46.45	19.64
Brown	120	109.99 ^{abc}	0.42	511.90	76.68	47.23	9.38
	Average	106.24 ^{AB}	0.43 ^B	516.55	76.32	46.84	14.51
Novogen	69	98.75 ^{cdef}	0.48	515.10	75.11	47.72	15.47
Brown	120	105.76 ^{abcde}	0.46	520.28	77.94	49.88	10.94
	Average	102.26 ^{AB}	0.47 ^{AB}	517.69	76.53	48.80	13.21
Lohmann	69	97.55 ^{def}	0.48	520.60	76.18	47.76	13.09
LB-Lite	120	107.16 ^{abcd}	0.48	552.42	82.95	52.56	9.38
	Average	102.35 ^{AB}	0.48 ^{AB}	536.51	79.56	50.16	11.23
Hy-Line	69	100.92 ^{bcdef}	0.45	529.33	77.61	46.21	3.57
Silver Brown	120	109.93 ^{abc}	0.42	524.72	78.69	47.64	6.25
	Average	105.43 ^{AB}	0.44 ^B	527.03	78.15	46.92	4.91
Hy-Line	69	96.03 ^{ef}	0.47	503.70	73.58	45.58	7.14
Brown	120	101.90 ^{bcdef}	0.47	523.18	78.40	48.91	10.94
	Average	98.96 ^B	0.47 ^{AB}	513.44	75.99	47.25	9.04
ISA	69	95.50 ^f	0.50	519.43	76.05	48.73	8.33
Brown	120	106.32 ^{abcd}	0.48	541.55	81.32	52.09	6.25
	Average	100.91 ^{AB}	0.49 ^A	530.49	78.68	50.41	7.29
Bovans	69	102.69 ^{abcdef}	0.46	525.98	77.01	48.71	11.61
Brown	120	111.41 ^a	0.46	534.68	80.35	51.90	3.12
	Average	107.05 ^A	0.46 ^{AB}	530.32	78.68	50.31	7.37
All	69	99.76 ^Y	0.46	518.56 ^Z	75.80 ^Z	47.14 ^Z	11.49
Strains	120	107.41 ^Z	0.46	528.22 ^Y	79.21 ^Y	49.61 ^Y	8.90

¹All strains were housed such that each strain is equally represented in each density.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdef - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 35. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69	59.89	0	3.19	8.46	25.03 ^{ab}	61.72 ^{ab}
	120	59.64	0	2.60	8.94	25.14 ^{ab}	62.92 ^{ab}
	Average	59.77 ^{AB}	0	2.89	8.70	25.08 ^{AB}	62.32 ^{AB}
TETRA Brown	69	58.69	0	2.72	5.43	22.04 ^{ab}	65.09 ^{ab}
	120	60.37	0	2.46	5.30	22.24 ^{ab}	67.48 ^{ab}
	Average	59.53 ^{AB}	0	2.59	5.37	22.14 ^{ABC}	66.28 ^{AB}
Novogen Brown	69	63.10	0.42	1.78	7.01	20.00 ^{ab}	70.04 ^{ab}
	120	63.32	0	1.63	3.88	17.73 ^{ab}	76.21 ^a
	Average	63.21 ^A	0.21	1.70	5.45	18.84 ^{ABC}	73.12 ^A
Lohmann LB-Lite	69	61.48	0	2.88	7.02	21.86 ^{ab}	66.62 ^{ab}
	120	62.58	1.63	2.08	3.25	19.76 ^{ab}	73.04 ^{ab}
	Average	62.03 ^{AB}	0.81	2.48	5.13	20.81 ^{ABC}	69.83 ^{AB}
Hy-Line Silver Brown	69	58.99	0	2.77	11.01	29.26 ^a	56.84 ^b
	120	58.92	0	1.37	6.99	24.39 ^{ab}	63.06 ^{ab}
	Average	58.96 ^B	0	2.07	9.00	26.82 ^A	59.95 ^B
Hy-Line Brown	69	60.91	0	1.23	6.09	26.01 ^{ab}	64.88 ^{ab}
	120	60.82	0	1.12	5.06	19.60 ^{ab}	71.85 ^{ab}
	Average	60.87 ^{AB}	0	1.18	5.58	22.81 ^{ABC}	68.36 ^{AB}
ISA Brown	69	62.19	1.45	2.17	3.93	19.17 ^{ab}	71.80 ^{ab}
	120	62.29	0	1.99	4.15	17.24 ^{ab}	73.92 ^{ab}
	Average	62.24 ^{AB}	0.72	2.08	4.04	18.21 ^{BC}	72.86 ^A
Bovans Brown	69	61.40	0	2.09	6.72	17.77 ^{ab}	71.21 ^{ab}
	120	61.58	0	1.24	3.16	14.65 ^b	76.53 ^a
	Average	61.49 ^{AB}	0	1.66	4.94	16.21 ^C	73.87 ^A
All Strains	69	60.83	0.23	2.35	6.96	22.64	66.02 ^Z
	120	61.19	0.20	1.81	5.09	20.09	70.63 ^Y

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 36. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69	92.42	0.17	2.67	0.36	56.12 ^{bcd}	26.27 ^{ab}
	120	94.12	0.08	2.42	0.12	57.48 ^{bcd}	26.75 ^{ab}
	Average	93.29	0.12 ^{AB}	2.54	0.24	56.80 ^{AB}	26.51
TETRA Brown	69	90.55	0.26	2.66	0.55	53.92 ^d	26.78 ^{ab}
	120	90.77	0.50	1.88	0.69	57.40 ^{bcd}	26.76 ^{ab}
	Average	90.66	0.38 ^{AB}	2.27	0.62	55.66 ^B	26.52
Novogen Brown	69	93.08	0.48	2.70	0.84	55.43 ^d	25.09 ^b
	120	92.57	0.06	2.38	0.70	59.43 ^{abcd}	26.45 ^{ab}
	Average	92.82	0.28 ^{AB}	2.54	0.77	57.43 ^{AB}	25.77
Lohmann LB-Lite	69	94.58	0	2.27	0.25	57.21 ^{bcd}	24.87 ^b
	120	94.42	0.08	1.88	0.44	63.09 ^a	26.90 ^{ab}
	Average	94.50	0.04 ^B	2.07	0.34	60.15 ^A	25.88
Hy-Line Silver Brown	69	92.47	0.13	2.92	0.18	58.47 ^{abcd}	26.04 ^{ab}
	120	90.43	0.16	2.24	0.82	58.54 ^{abcd}	27.48 ^a
	Average	91.44	0.14 ^{AB}	2.58	0.50	58.51 ^{AB}	26.76
Hy-Line Brown	69	91.10	0.27	2.44	0.38	55.62 ^{cd}	25.15 ^b
	120	89.35	0.41	4.78	0.36	59.21 ^{abcd}	26.15 ^{ab}
	Average	90.23	0.34 ^{AB}	3.61	0.37	57.42 ^{AB}	25.65
ISA Brown	69	93.95	0.81	2.29	0.11	57.86 ^{abcd}	24.92 ^b
	120	92.40	0.63	2.33	0.56	61.60 ^{ab}	26.53 ^{ab}
	Average	93.17	0.72 ^A	2.31	0.33	59.73 ^{AB}	25.73
Bovans Brown	69	90.76	0.36	2.25	0.10	57.69 ^{abcd}	25.80 ^{ab}
	120	90.62	0.43	3.32	0.19	61.39 ^{abc}	27.64 ^a
	Average	90.69	0.40 ^{AB}	2.79	0.14	59.54 ^{AB}	26.72
All Strains	69	92.37	0.31	2.53	0.35	56.54 ^Z	26.83 ^Z
	120	91.84	0.29	2.65	0.48	59.77 ^Y	25.55 ^Y

¹All strains were housed such that each strain is equally represented in each density.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 37. EFFECT OF WHITE EGG STRAIN AND DENSITY ON BODY WEIGHT OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	17 Wk Body Wt	69 Wk Body Wt	1st Cycle Wt Gain	73 Wk Body Wt	109 Wk Body Wt	2 nd Cycle Wt Gain	Total Wt Gain
(Strain)	(in ² /hen)	(kg)	(kg)	(%)	(kg)	(kg)	(kg)	(%)
Bovans	69	1.26	1.78	42.9	1.48	1.83	0.46	45.74
White	120	1.28	1.85	48.4	1.44	1.91	0.55	50.02
	Average	1.27	1.81	45.7	1.46	1.87 ^{ABC}	0.51	47.88
Shaver	69	1.31	1.80	36.6	1.45	1.83	0.38	39.69
White	120	1.31	1.98	52.7	1.50	1.95	0.44	49.13
	Average	1.31	1.89	44.3	1.48	1.89 ^{ABC}	0.41	44.41
Dekalb	69	1.30	1.77	36.2	1.40	1.94	0.58	49.25
White	120	1.30	1.80	38.5	1.52	1.91	0.33	47.42
	Average	1.30	1.78	37.7	1.46	1.92 ^{ABC}	0.46	48.34
Babcock	69	1.35	1.90	43.0	1.44	2.10	0.68	56.32
White	120	1.36	2.04	52.9	1.52	2.10	0.42	53.26
	Average	1.36	1.97	47.8	1.48	2.10 ^A	0.55	54.79
ISA	69	1.22	1.70	37.7	1.36	1.65	0.43	34.70
B-400	120	1.20	1.82	49.2	1.38	1.70	0.28	41.05
	Average	1.21	1.76	43.8	1.37	1.67 ^C	0.35	37.87
Hy-Line	69	1.56	1.82	36.5	1.46	1.81	0.26	22.23
W-36	120	1.21	1.92	61.2	1.50	2.00	0.48	66.35
	Average	1.38	1.87	47.8	1.48	1.91 ^{ABC}	0.36	44.29
Hy-Line	69	1.22	1.73	41.0	1.38	1.93	0.56	58.82
CV-26	120	1.22	1.84	50.8	1.43	1.92	0.58	57.45
	Average	1.22	1.78	45.9	1.40	1.93 ^{ABC}	0.57	58.14
Hy-Line	69	1.29	1.79	39.5	1.44	1.73	0.38	34.21
CV-24	120	1.30	1.85	44.6	1.58	1.83	0.38	40.18
	Average	1.30	1.82	41.5	1.51	1.78 ^{BC}	0.38	37.19
Hy-Line	69	1.31	1.88	42.7	1.57	1.95	0.36	48.62
CV-22	120	1.30	1.93	47.7	1.52	2.01	0.44	55.21
	Average	1.30	1.90	45.4	1.54	1.98 ^{AB}	0.40	51.92
Lohmann	69	1.27	1.72	36.2	1.34	1.92	0.58	50.81
LSL Lite	120	1.28	1.85	46.9	1.46	1.96	0.50	53.34
	Average	1.28	1.79	41.4	1.40	1.94 ^{ABC}	0.54	52.07
H&N	69	1.34	1.74	28.4	1.43	1.92	0.49	43.40
Nick Chick	120	1.29	1.84	41.9	0.72	1.92	0.60	48.75
	Average	1.31	1.79	35.1	1.07	1.92 ^{ABC}	0.54	46.07
Novogen	69	1.34	1.79	33.6	1.38	1.92	0.50	42.96
White	120	1.26	1.99	58.7	1.43	1.87	0.50	48.00
	Average	1.30	1.89	46.2	1.41	1.90 ^{ABC}	0.50	45.48
All	69	1.31	1.78 ^Y	38.2 ^Y	1.43	1.88	0.47	43.89
Strains	120	1.28	1.89 ^Z	49.2 ^Z	1.42	1.92	0.46	50.85

¹All strains were housed such that each strain is equally represented in each density.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 38. EFFECT OF BROWN EGG STRAIN AND DENSITY ON BODY WEIGHT OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN CONVENTIONAL CAGES USING THE NON-ANOREXIC MOLT PROGRAM

Breeder	Density ¹	17 Wk Body Wt	69 Wk Body Wt	1st Cycle Wt Gain	73 Wk Body Wt	109 Wk Body Wt	2 nd Cycle Wt Gain	Total Wt Gain
(Strain)	(in ² /hen)	(kg)	(kg)	(%)	(kg)	(kg)	(kg)	(%)
TETRA	69	1.67	2.30	37.1	1.90	2.20	0.29	32.28
Amber	120	1.60	2.18	36.3	1.86	2.18	0.31	36.76
	Average	1.63	2.24	36.8	1.88	2.19	0.30	34.52
TETRA	69	1.58	1.96	23.4	1.69	1.98	0.30	25.28
Brown	120	1.63	2.06	27.0	1.76	2.07	0.31	27.18
	Average	1.61	2.01	24.8	1.72	2.03	0.30	26.23
Novogen	69	1.67	2.03	21.0	1.73	2.01	0.34	20.78
Brown	120	1.59	2.06	31.4	1.78	2.06	0.22	29.43
	Average	1.63	2.05	26.4	1.76	2.03	0.28	25.10
Lohmann	69	1.63	1.96	21.5	1.61	2.03	0.36	24.42
LB-Lite	120	1.64	2.12	30.5	1.74	2.02	0.24	23.33
	Average	1.63	2.04	26.4	1.68	2.02	0.30	23.87
Hy-Line	69	1.62	2.01	25.9	1.75	2.25	0.48	38.63
Silver Brown	120	1.64	2.36	48.2	1.88	2.20	0.50	35.44
	Average	1.63	2.19	36.8	1.81	2.22	0.49	37.03
Hy-Line	69	1.61	1.96	21.7	1.66	2.05	0.26	26.78
Brown	120	1.65	2.14	30.3	1.80	2.16	0.32	31.00
	Average	1.63	2.05	25.8	1.73	2.10	0.29	28.89
ISA	69	1.53	1.91	24.8	1.64	1.98	0.35	29.13
Brown	120	1.59	2.20	37.7	1.82	2.15	0.33	35.68
	Average	1.56	2.06	31.4	1.73	2.06	0.34	32.40
Bovans	69	1.56	2.00	28.2	1.70	2.27	0.42	45.28
Brown	120	1.62	2.06	27.8	1.70	2.12	0.48	31.42
	Average	1.59	2.03	27.7	1.70	2.20	0.45	38.35
All	69	1.61	2.02	25.5 ^Y	1.71	2.10	0.35	30.32
Strains	120	1.62	2.15	34.0 ^Z	1.79	2.12	0.34	31.28

¹All strains were housed such that each strain is equally represented in each density.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 39. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
Bovans White	69 EC	10.28 ^{efg}	0.505 ^{bcdef}	311.82 ^{ab}	85.48 ^D	50.93 ^d	7.78 ^{ab}	144.90 ^{abc}
	69 ECS	10.51 ^{bcdef}	0.492 ^{def}	311.22 ^{ab}	84.74 ^{def}	50.87 ^d	17.06 ^{ab}	146.14 ^a
	Average	10.40 ^{BC}	0.499 ^{BC}	311.52 ^{ABC}	85.11 ^B	50.90 ^{BC}	12.42 ^{AB}	145.52 ^A
Shaver White	69 EC	10.16 ^{fg}	0.542 ^{abc}	307.19 ^{ab}	83.79 ^{defg}	50.89 ^d	16.67 ^{ab}	142.00 ^{bcd}
	69 ECS	10.50 ^{bcdef}	0.540 ^{abcd}	318.28 ^{ab}	85.56 ^{bcd}	51.91 ^{cd}	30.55 ^a	140.50 ^d
	Average	10.33 ^{CD}	0.541 ^A	312.74 ^{ABC}	84.68 ^B	51.40 ^B	23.61 ^A	141.25 ^C
Dekalb White	69 EC	10.56 ^{abcdef}	0.503 ^{bcdef}	309.92 ^{ab}	84.92 ^{def}	51.18 ^{cd}	7.22 ^{ab}	144.40 ^{abcd}
	69 ECS	10.48 ^{bcdefg}	0.508 ^{bcdef}	311.31 ^{ab}	85.18 ^{cdef}	51.54 ^{cd}	9.03 ^{ab}	144.25 ^{abcd}
	Average	10.52 ^{ABC}	0.506 ^B	310.62 ^{ABC}	85.05 ^B	51.36 ^B	8.12 ^{AB}	144.32 ^{AB}
Babcock White	69 EC	10.31 ^{defg}	0.560 ^a	314.52 ^{ab}	86.32 ^{bcd}	52.84 ^{abc}	3.09 ^b	141.50 ^{cd}
	69 ECS	11.09 ^a	0.539 ^{abcd}	327.38 ^a	89.54 ^a	54.27 ^{ab}	11.80 ^{ab}	140.50 ^d
	Average	10.55 ^A	0.549 ^A	320.95 ^{AB}	87.93 ^A	53.55 ^A	7.44 ^B	140.00 ^C
ISA B-400	69 EC	10.63 ^{abcde}	0.532 ^{abcd}	321.48 ^a	88.17 ^{abc}	54.63 ^a	5.96 ^{ab}	144.40 ^{abcd}
	69 ECS	10.68 ^{abcde}	0.551 ^{ab}	325.21 ^a	88.61 ^{ab}	54.75 ^a	17.50 ^{ab}	141.00 ^d
	Average	10.49 ^{ABC}	0.542 ^A	323.35 ^A	88.39 ^A	54.69 ^A	11.73 ^{AB}	142.70 ^{BC}
Hy-Line W-36	69 EC	10.11 ^{fg}	0.483 ^{ef}	297.80 ^b	81.72 ^{gh}	48.79 ^f	3.14 ^b	146.17 ^a
	69 ECS	9.89 ^g	0.493 ^{bcdef}	298.27 ^{ab}	81.81 ^{fgh}	48.22 ^f	4.63 ^{ab}	145.33 ^{abcd}
	Average	10.00 ^D	0.488 ^{BC}	298.04 ^C	81.73 ^D	48.51 ^D	3.84 ^B	145.75 ^A
Hy-Line CV-24	69 EC	10.83 ^{abc}	0.474 ^{ef}	309.24 ^{ab}	84.57 ^{def}	50.90 ^{de}	10.00 ^{ab}	144.60 ^{abcd}
	69 ECS	10.80 ^{abcd}	0.464 ^{ef}	295.24 ^b	80.70 ^h	48.93 ^{ef}	11.11 ^{ab}	145.75 ^{abc}
	Average	10.82 ^A	0.469 ^C	302.23 ^C	82.63 ^{CD}	49.91 ^C	10.56 ^{AB}	145.18 ^{AB}
Lohmann LSL Lite	69 EC	10.83 ^{abc}	0.493 ^{bcdef}	309.41 ^{ab}	84.63 ^{defg}	51.55 ^{cd}	13.20 ^{ab}	146.00 ^{ab}
	69 ECS	10.68 ^{abcde}	0.498 ^{bcdef}	310.94 ^{ab}	84.85 ^{def}	51.60 ^{cd}	13.19 ^{ab}	144.75 ^{abcd}
	Average	10.61 ^{ABC}	0.496 ^{BC}	310.18 ^{ABC}	84.74 ^B	51.57 ^B	13.20 ^{AB}	145.38 ^{AB}
H&N Nick Chick	69 EC	10.49 ^{bcdef}	0.500 ^{bcdef}	307.67 ^{ab}	84.20 ^{defg}	51.52 ^{cd}	9.26 ^{ab}	145.33 ^{ab}
	69 ECS	10.66 ^{abcde}	0.497 ^{bcdef}	309.98 ^{ab}	84.44 ^{defg}	51.63 ^{cd}	15.97 ^{ab}	145.75 ^{abc}
	Average	10.57 ^{ABC}	0.499 ^{BC}	308.82 ^{ABC}	84.32 ^{BC}	51.58 ^B	12.62 ^{AB}	145.54 ^A
Novogen White	69 EC	10.38 ^{cdefg}	0.519 ^{abcde}	309.07 ^{ab}	84.59 ^{def}	52.31 ^{bcd}	11.11 ^{ab}	144.60 ^{abcd}
	69 ECS	10.49 ^{ab}	0.480 ^{ef}	304.77 ^{ab}	82.70 ^{efgh}	50.91 ^{de}	20.55 ^{ab}	145.00 ^{abcd}
	Average	10.66 ^{AB}	0.500 ^{BC}	306.92 ^{BC}	83.64 ^{BCD}	51.61 ^B	15.83 ^{AB}	144.80 ^{AB}
All Strains	69 EC	10.43 ^Z	0.511	309.81	84.84	51.55	8.74 ^Y	144.39
	69 ECS	10.62 ^Y	0.506	311.26	84.81	51.46	15.14 ^Z	143.90

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefgh - Different letters denote significant differences (P<.01) in the strain*housing system interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 40. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 EC	58.46 ^h	0 ^{ab}	7.72	14.00 ^{ab}	25.97	51.83 ^{de}
	69 ECS	58.97 ^{efgh}	0 ^{ab}	7.74	12.37 ^{ab}	25.18	54.63 ^{cde}
	Average	58.72 ^{EF}	0	7.73 ^{AB}	13.19 ^{AB}	25.57	53.23 ^C
Shaver White	69 EC	59.64 ^{cdefg}	0.04 ^{ab}	6.28	10.38 ^{ab}	24.73	58.26 ^{abc}
	69 ECS	59.78 ^{abcdefg}	0 ^{ab}	4.75	11.86 ^{ab}	24.22	58.66 ^{abc}
	Average	59.71 ^{CD}	0.02	5.52 ^{AB}	11.12 ^{BC}	24.48	58.46 ^B
Dekalb White	69 EC	59.17 ^{efgh}	0 ^{ab}	6.33	11.48 ^{ab}	24.27	57.62 ^{abcd}
	69 ECS	59.22 ^{defgh}	0 ^{ab}	6.87	10.52 ^{ab}	23.43	58.78 ^{abc}
	Average	59.19 ^{DE}	0	6.60 ^{AB}	11.00 ^{BC}	23.85	58.20 ^B
Babcock White	69 EC	60.29 ^{abcd}	0 ^{ab}	5.32	11.06 ^{ab}	22.39	60.91 ^{ab}
	69 ECS	59.65 ^{bcdefg}	0.52 ^a	5.08	11.98 ^{ab}	23.34	58.52 ^{abc}
	Average	59.97 ^{BC}	0.26	5.20 ^{AB}	11.52 ^{ABC}	22.87	59.71 ^{AB}
ISA B-400	69 EC	60.88 ^a	0.05 ^{ab}	5.30	8.79 ^b	21.47	64.04 ^a
	69 ECS	60.81 ^{ab}	0 ^{ab}	4.35	9.28 ^{ab}	22.25	63.64 ^a
	Average	60.85 ^A	0.03	4.83 ^B	9.03 ^C	21.86	63.84 ^A
Hy-Line W-36	69 EC	58.75 ^{ef}	0 ^{ab}	7.68	14.63 ^{ab}	22.45	54.93 ^{bcd}
	69 ECS	58.02 ^h	0 ^{ab}	8.30	17.08 ^a	25.58	48.59 ^e
	Average	58.38 ^F	0	7.99 ^{AB}	15.85 ^A	24.02	51.76 ^C
Hy-Line CV-24	69 EC	58.97 ^{efgh}	0 ^{ab}	5.88	13.77 ^{ab}	22.40	57.80 ^{abc}
	69 ECS	58.86 ^{abcdefg}	0 ^{ab}	8.21	11.03 ^{ab}	22.63	57.69 ^{abcd}
	Average	59.41 ^{CD}	0	7.04 ^{AB}	12.40 ^{ABC}	22.52	57.75 ^B
Lohmann LSL Lite	69 EC	59.90 ^{abcdef}	0 ^{ab}	6.25	11.93 ^{ab}	22.48	59.26 ^{abc}
	69 ECS	59.66 ^{bcdefg}	0 ^{ab}	5.05	10.80 ^{ab}	24.03	59.66 ^{abc}
	Average	59.78 ^{BCD}	0	5.65 ^{AB}	11.36 ^{ABC}	23.26	59.46 ^{AB}
H&N Nick Chick	69 EC	59.97 ^{abcde}	0 ^{ab}	6.13	9.73 ^{ab}	22.84	61.15 ^{ab}
	69 ECS	60.14 ^{abcde}	0 ^{ab}	5.35	11.01 ^{ab}	22.45	60.82 ^{abc}
	Average	60.05 ^{BC}	0	5.74 ^{AB}	10.37 ^B	22.64	60.99 ^{AB}
Novogen White	69 EC	60.57 ^{abc}	0 ^{ab}	5.24	9.82 ^{ab}	21.76	63.07 ^a
	69 ECS	60.51 ^{abc}	0.01 ^{ab}	5.31	10.84 ^{ab}	22.13	61.34 ^{ab}
	Average	60.54 ^{AB}	0	5.28 ^{AB}	10.33 ^{BC}	21.95	62.20 ^{AB}
All Strains	69 EC	59.66	0.01	6.21	11.56	23.08	58.89
	69 ECS	59.66	0.05	6.10	11.68	23.52	58.23

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABCDEF - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefgh - Different letters denote significant differences (P<.01) in the strain*production system interactions.

TABLE 41. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans	69 EC	96.47	0.40	2.52	0.63	35.22 ^{ab}	15.37 ^{abc}
White	69 ECS	94.83	0.50	4.38	0.31	35.17 ^{ab}	15.71 ^{abc}
	Average	95.65	0.45	3.45	0.47	35.19 ^{ABC}	15.54 ^{AB}
Shaver	69 EC	95.80	0.38	3.38	0.48	34.96 ^{ab}	15.19 ^{bc}
White	69 ECS	93.62	0.42	5.00	0.95	35.79 ^{ab}	15.70 ^{abc}
	Average	94.71	0.40	4.19	0.72	35.38 ^{ABC}	15.44 ^{AB}
Dekalb	69 EC	97.16	0.40	1.88	0.58	35.48 ^{ab}	15.78 ^{abc}
White	69 ECS	92.75	0.55	6.10	0.58	35.19 ^{ab}	15.68 ^{abc}
	Average	94.96	0.48	3.99	0.58	35.34 ^{ABC}	15.73 ^{AB}
Babcock	69 EC	96.72	0.62	2.10	0.60	36.04 ^{ab}	15.42 ^{abc}
White	69 ECS	93.50	0.80	5.05	0.68	36.79 ^a	16.56 ^a
	Average	95.11	0.71	3.58	0.64	36.42 ^A	15.99 ^A
ISA	69 EC	96.12	0.32	2.82	0.72	36.96 ^a	15.91 ^{abc}
B-400	69 ECS	93.82	0.45	5.20	0.55	36.91 ^a	15.97 ^{abc}
	Average	94.97	0.38	4.01	0.64	36.94 ^A	15.94 ^A
Hy-Line	69 EC	96.64	0.54	2.46	0.34	33.70 ^b	14.95 ^c
W-36	69 ECS	93.57	0.37	5.23	0.77	32.87 ^b	14.81 ^c
	Average	95.10	0.45	3.85	0.55	33.28 ^C	14.88 ^B
Hy-Line	69 EC	96.38	0.46	2.98	0.14	35.22 ^{ab}	16.20 ^{abc}
CV-24	69 ECS	93.60	0.38	5.50	0.58	33.04 ^b	16.14 ^{abc}
	Average	94.99	0.42	4.24	0.36	34.13 ^{BC}	16.17 ^A
Lohmann	69 EC	97.15	0.42	2.30	0.15	35.55 ^{ab}	15.76 ^{abc}
LSL Lite	69 ECS	92.90	1.15	5.45	0.52	35.21 ^{ab}	15.96 ^{abc}
	Average	95.02	0.79	3.88	0.34	35.38 ^{ABC}	15.86 ^A
H&N	69 EC	95.95	0.87	2.75	0.40	35.31 ^{ab}	15.68 ^{abc}
Nick Chick	69 ECS	93.90	0.65	4.58	0.88	35.07 ^{ab}	15.94 ^{abc}
	Average	94.92	0.76	3.66	0.64	35.19 ^{ABC}	15.81 ^A
Novogen	69 EC	96.70	0.16	2.82	0.32	35.63 ^{ab}	15.50 ^{abc}
White	69 ECS	93.94	0.52	5.02	0.52	34.59 ^{ab}	16.39 ^{ab}
	Average	95.32	0.34	3.92	0.42	35.11 ^{ABC}	15.95 ^A
All	69 EC	96.51 ^Y	0.46	2.60 ^Y	0.44	35.40	15.57 ^Z
Strains	69 ECS	93.64 ^Z	0.58	5.15 ^Z	0.63	35.06	15.88 ^Y

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*production system interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 42. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
TETRA Amber	69 EC	10.46 ^{bc}	0.456 ^{bc}	293.78 ^{ab}	80.49 ^{bcde}	46.53 ^{fgh}	7.14 ^{ab}	145.00 ^{abc}
	69 ECS	10.60 ^{abc}	0.442 ^c	283.01 ^{ab}	77.69 ^{ef}	44.55 ^h	1.39 ^b	145.50 ^{abc}
	Average	10.53 ^{BC}	0.449 ^C	288.40 ^{AB}	79.09 ^{BC}	45.54 ^C	4.27 ^{AB}	145.25 ^{AB}
TETRA Brown	69 EC	10.20 ^{cd}	0.475 ^{abc}	276.75 ^b	75.57 ^f	45.21 ^{gh}	14.82 ^a	143.67 ^{abc}
	69 ECS	10.38 ^{bcd}	0.464 ^{abc}	282.18 ^{ab}	77.28 ^{ef}	45.84 ^{fgh}	5.56 ^{ab}	145.00 ^{abc}
	Average	10.29 ^{CD}	0.469 ^{BC}	279.46 ^B	76.43 ^D	45.52 ^C	10.19 ^A	144.33 ^{ABC}
Novogen Brown	69 EC	10.28 ^{cd}	0.513 ^a	293.64 ^{ab}	80.45 ^{bcde}	48.80 ^{abcde}	7.78 ^{ab}	144.00 ^{abc}
	69 ECS	10.49 ^{bc}	0.510 ^a	297.85 ^{ab}	81.75 ^{abc}	49.82 ^{abc}	3.34 ^{ab}	144.00 ^{abc}
	Average	10.39 ^{BCD}	0.512 ^A	295.74 ^{AB}	81.10 ^{AB}	49.31 ^A	5.56 ^{AB}	144.00 ^{ABC}
Lohmann LB-Lite	69 EC	10.36 ^{bcd}	0.497 ^{abc}	301.64 ^{ab}	82.72 ^{ab}	49.72 ^{abc}	4.79 ^{ab}	145.00 ^{abc}
	69 ECS	10.36 ^{bcd}	0.484 ^{abc}	294.17 ^{AB}	80.74 ^{bcde}	48.48 ^{abcdef}	4.48 ^{ab}	145.75 ^{abc}
	Average	10.36 ^{CD}	0.491 ^{AB}	297.91 ^{AB}	81.73 ^A	49.10 ^A	4.13 ^{AB}	145.38 ^{AB}
Hy-Line Silver Brown	69 EC	10.55 ^{abc}	0.483 ^{abc}	297.54 ^{ab}	81.59 ^{abcd}	47.14 ^{defg}	5.79 ^{ab}	143.50 ^{abc}
	69 ECS	10.79 ^{ab}	0.487 ^{abc}	303.42 ^{ab}	83.20 ^{ab}	47.60 ^{bcdefg}	6.94 ^{ab}	141.50 ^{bc}
	Average	10.67 ^{AB}	0.485 ^{ABC}	300.48 ^A	82.39 ^A	47.37 ^B	6.37 ^{AB}	142.50 ^{BC}
Hy-Line Brown	69 EC	10.28 ^{cd}	0.518 ^a	285.52 ^{ab}	78.46 ^{def}	47.03 ^{efg}	0.46 ^b	141.17 ^c
	69 ECS	10.65 ^{abc}	0.494 ^{abc}	289.23 ^{ab}	79.35 ^{cdef}	47.64 ^{cdefg}	4.44 ^{ab}	142.20 ^{abc}
	Average	10.47 ^{BC}	0.506 ^{AB}	287.38 ^{AB}	78.90 ^C	47.34 ^B	2.45 ^B	141.68 ^C
ISA Brown	69 EC	10.03 ^d	0.499 ^{abc}	293.07 ^{ab}	80.41 ^{bcde}	48.91 ^{abcde}	3.70 ^{ab}	147.33 ^a
	69 ECS	10.37 ^{bcd}	0.504 ^{ab}	308.91 ^a	84.74 ^a	50.87 ^a	4.45 ^{ab}	146.00 ^{abc}
	Average	10.20 ^D	0.501 ^{AB}	300.99 ^A	82.57 ^A	49.89 ^A	4.08 ^{AB}	146.67 ^A
Bovans Brown	69 EC	10.64 ^{abc}	0.473 ^{abc}	296.33 ^{ab}	81.30 ^{bcd}	49.06 ^{abcd}	3.57 ^{ab}	146.43 ^{ab}
	69 ECS	10.96 ^a	0.472 ^{abc}	299.50 ^{ab}	82.12 ^{abc}	49.92 ^{ab}	5.56 ^{ab}	146.00 ^{abc}
	Average	10.80 ^A	0.472 ^{BC}	297.92 ^{AB}	81.74 ^A	49.49 ^A	4.56 ^{AB}	146.21 ^A
All Strains	69 EC	10.58 ^Z	0.489	292.28	80.13	47.80	6.01	144.51
	69 ECS	10.35 ^Y	0.482	294.78	80.86	48.09	4.39	144.49

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde fgh - Different letters denote significant differences (P<.01) in the strain*production system interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 43. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder (Strain)	Production System	Egg Weight (g/egg)	Pee Wee (%)	Small (%)	Medium (%)	Large (%)	Extra Large (%)
TETRA Amber	69 EC	56.83 ^{cd}	0.03	7.10 ^a	21.67 ^{ab}	29.24	41.57 ^c
	69 ECS	56.51 ^{cd}	0.48	7.88 ^a	22.25 ^{abc}	28.06	41.13 ^c
	Average	56.67 ^B	0.26	7.49 ^A	21.96 ^A	28.65 ^{ABC}	41.35 ^B
TETRA Brown	69 EC	59.15 ^{ab}	0	4.77 ^{ab}	15.67 ^{bcde}	26.61	52.67 ^{ab}
	69 ECS	58.56 ^{abc}	0	4.38 ^{ab}	15.79 ^{bcde}	28.77	51.00 ^{ab}
	Average	58.86 ^A	0	4.58 ^{ABC}	15.73 ^B	27.69 ^{ABC}	51.83 ^A
Novogen Brown	69 EC	59.91 ^a	0	3.66 ^{ab}	11.89 ^{de}	26.09	57.92 ^a
	69 ECS	60.19 ^a	0	3.24 ^{ab}	13.49 ^{de}	25.86	57.09 ^a
	Average	60.05 ^A	0	3.45 ^{BC}	12.69 ^B	25.98 ^{BC}	57.51 ^A
Lohmann LB-Lite	69 EC	59.03 ^a	0	5.78 ^{ab}	12.78 ^{de}	24.97	56.24 ^a
	69 ECS	59.04 ^{ab}	0	4.96 ^{ab}	13.32 ^{de}	24.27	56.92 ^a
	Average	59.04 ^A	0	5.37 ^{AB}	13.05 ^B	24.62 ^C	56.58 ^A
Hy-Line Silver Brown	69 EC	57.12 ^{bcd}	0.12	4.50 ^{ab}	19.23 ^{abcd}	32.75	43.29 ^{bc}
	69 ECS	55.56 ^d	0.01	4.46 ^{ab}	25.31 ^a	31.84	38.47 ^c
	Average	56.49 ^B	0.06	4.48 ^{ABC}	22.27 ^A	32.30 ^A	40.88 ^B
Hy-Line Brown	69 EC	59.42 ^a	0.05	2.31 ^b	15.59 ^{de}	30.00	53.74 ^a
	69 ECS	59.50 ^a	0	1.61 ^b	12.23 ^{de}	31.83	53.86 ^a
	Average	59.46 ^A	0.08	1.96 ^C	12.91 ^B	30.92 ^{AB}	53.80 ^A
ISA Brown	69 EC	59.90 ^a	0	4.37 ^{ab}	11.43 ^e	27.59	56.40 ^a
	69 ECS	58.96 ^a	0.51	2.80 ^{ab}	14.40 ^{cde}	28.69	53.25 ^a
	Average	59.43 ^A	0.25	3.58 ^{BC}	12.92 ^B	28.14 ^{ABC}	54.82 ^A
Bovans Brown	69 EC	59.48 ^a	0.03	3.70 ^{ab}	15.53 ^{bcde}	25.57	54.80 ^a
	69 ECS	60.44 ^a	0	3.60 ^{ab}	13.83 ^{de}	26.51	55.78 ^a
	Average	59.96 ^A	0.02	3.65 ^{BC}	14.68 ^B	26.04 ^{BC}	55.29 ^A
All Strains	69 EC	58.86	0.04	4.52	15.22	27.85	52.08
	69 ECS	58.63	0.12	4.12	16.33	28.23	50.94

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*production system interactions.

TABLE 44. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder (Strain)	Production System	Grade A (%)	Grade B (%)	Cracks (%)	Loss (%)	Egg Income (\$/hen)	Feed Costs (\$/hen)
TETRA Amber	69 EC	96.13 ^{abc}	0.34	2.90 ^{bcd}	0.66	32.45 ^{abc}	15.48 ^{ab}
	69 ECS	92.58 ^d	0.42	6.62 ^a	0.38	30.55 ^c	15.68 ^{ab}
	Average	94.35	0.38	4.76	0.52	31.50 ^B	15.58 ^{AB}
TETRA Brown	69 EC	95.33 ^{abcd}	0.20	3.83 ^{abcd}	0.63	31.09 ^{bc}	15.09 ^{ab}
	69 ECS	92.87 ^{bcd}	0.60	5.87 ^{abc}	0.70	31.43 ^{abc}	15.38 ^{ab}
	Average	94.10	0.40	4.85	0.67	31.26 ^B	15.23 ^{AB}
Novogen Brown	69 EC	96.50 ^{abc}	0.34	2.58 ^{cd}	0.62	33.61 ^{abc}	15.21 ^{ab}
	69 ECS	92.70 ^d	0.62	5.94 ^{ab}	0.72	33.35 ^{abc}	15.55 ^{ab}
	Average	94.60	0.48	4.26	0.67	33.50 ^{AB}	15.38 ^{AB}
Lohmann LB-Lite	69 EC	96.70 ^{ab}	0.45	2.32 ^d	0.50	34.44 ^{ab}	15.34 ^{ab}
	69 ECS	92.92 ^{cd}	0.38	5.52 ^{abcd}	1.18	33.06 ^{abc}	15.34 ^{ab}
	Average	94.81	0.41	3.92	0.84	33.75 ^A	15.34 ^{AB}
Hy-Line Silver Brown	69 EC	97.05 ^a	0.08	2.65 ^{bcd}	0.22	33.45 ^{abc}	15.61 ^{ab}
	69 ECS	95.20 ^{abcd}	0.10	4.70 ^{abcd}	0	33.45 ^{abc}	15.97 ^{ab}
	Average	96.12	0.09	3.68	0.11	33.45 ^{AB}	15.79 ^{AB}
Hy-Line Brown	69 EC	95.85 ^{abcd}	0.37	3.27 ^{bcd}	0.57	32.45 ^{abc}	15.23 ^{ab}
	69 ECS	94.06 ^{abcd}	0.22	5.20 ^{abcd}	0.56	32.74 ^{abc}	15.76 ^{ab}
	Average	94.96	0.29	4.23	0.56	32.59 ^{AB}	15.50 ^{AB}
ISA Brown	69 EC	95.93 ^{abcd}	0.98	2.57 ^{cd}	0.52	33.59 ^{abc}	14.83 ^b
	69 ECS	94.06 ^{abcd}	0.22	5.30 ^{abcd}	0.44	34.95 ^a	15.36 ^{ab}
	Average	95.00	0.60	3.93	0.48	34.27 ^A	15.10 ^B
Bovans Brown	69 EC	95.44 ^{abcd}	0.43	3.43 ^{bcd}	0.68	33.49 ^{abc}	15.75 ^{ab}
	69 ECS	94.70 ^{abcd}	0.16	4.48 ^{abcd}	0.68	33.91 ^{abc}	16.22 ^a
	Average	95.07	0.29	3.95	0.68	33.70 ^A	15.99 ^A
All Strains	69 EC	96.12 ^Y	0.40	2.94 ^Y	0.58	33.07	15.66 ^Y
	69 ECS	93.63 ^Z	0.34	5.45 ^Z	0.55	32.94	15.32 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 45. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality*
(Strain)		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans White	69 EC	11.45 ^{abc}	0.476 ^{ab}	24.85 ^a	87.38 ^{ab}	54.56 ^{ab}	3.33
	69 ECS	11.25 ^{abc}	0.468 ^{ab}	23.68 ^{ab}	83.69 ^{ab}	52.82 ^{ab}	1.39
	Average	11.35 ^{AB}	0.472 ^{AB}	24.27	85.53	53.69	2.36
Shaver White	69 EC	10.70 ^{bc}	0.478 ^{ab}	22.22 ^{ab}	78.85 ^{ab}	50.87 ^{ab}	0.93
	69 ECS	11.89 ^{abc}	0.455 ^{ab}	24.15 ^{ab}	83.16 ^{ab}	53.75 ^{ab}	3.47
	Average	11.30 ^{AB}	0.467 ^{AB}	23.18	81.00	52.31	2.20
Dekalb White	69 EC	11.20 ^{abc}	0.490 ^{ab}	24.38 ^{ab}	86.98 ^{ab}	54.91 ^{ab}	0.56
	69 ECS	11.10 ^{abc}	0.495 ^{ab}	24.41 ^{ab}	85.55 ^{ab}	54.68 ^{ab}	1.39
	Average	11.15 ^{AB}	0.492 ^{AB}	24.40	86.76	54.79	0.97
Babcock White	69 EC	10.7 ^{bc}	0.521 ^a	25.04 ^a	87.97 ^a	56.00 ^{ab}	1.85
	69 ECS	12.82 ^a	0.415 ^{ab}	23.30 ^{ab}	82.46 ^{ab}	52.98 ^{ab}	2.08
	Average	11.78 ^A	0.468 ^{AB}	24.17	85.22	54.49	1.97
ISA B-400	69 EC	10.98 ^{abc}	0.522 ^a	24.93 ^a	89.02 ^a	57.06 ^a	0
	69 ECS	10.98 ^{abc}	0.510 ^{ab}	24.32 ^{ab}	86.43 ^{ab}	56.10 ^{ab}	0.70
	Average	10.98 ^{AB}	0.516 ^A	24.62	87.73	56.58	0.35
Hy-Line W-36	69 EC	10.34 ^c	0.497 ^{ab}	22.47 ^{ab}	80.26 ^{ab}	51.25 ^{ab}	0
	69 ECS	10.08 ^c	0.480 ^{ab}	21.59 ^{ab}	79.10 ^{ab}	48.55 ^{ab}	0
	Average	10.21 ^B	0.488 ^{AB}	22.03	78.68	49.90	0
Hy-Line CV-24	69 EC	11.09 ^{abc}	0.488 ^{ab}	22.93 ^{ab}	81.91 ^{ab}	53.42 ^{ab}	0
	69 ECS	12.16 ^{ab}	0.442 ^{ab}	22.97 ^{ab}	82.02 ^{ab}	53.75 ^{ab}	0.70
	Average	11.62 ^A	0.465 ^{AB}	22.95	81.96	53.59	0.35
Lohmann LSL Lite	69 EC	11.95 ^{abc}	0.460 ^{ab}	24.35 ^{ab}	83.23 ^{ab}	55.24 ^{ab}	4.86
	69 ECS	11.86 ^{abc}	0.388 ^b	19.92 ^b	71.13 ^b	45.64 ^b	0
	Average	11.90 ^A	0.424 ^B	22.13	77.18	50.44	2.43
H&N Nick Chick	69 EC	11.27 ^{abc}	0.493 ^{ab}	23.89 ^{ab}	85.27 ^{ab}	55.49 ^{ab}	0.46
	69 ECS	12.03 ^{abc}	0.458 ^{ab}	23.96 ^{ab}	84.76 ^{ab}	55.11 ^{ab}	0.70
	Average	11.65 ^A	0.475 ^{AB}	23.92	85.10	55.30	0.58
Novogen White	69 EC	11.28 ^{abc}	0.500 ^{ab}	24.74 ^{ab}	86.12 ^{ab}	56.27 ^{ab}	1.12
	69 ECS	11.77 ^{abc}	0.428 ^{ab}	22.04 ^{ab}	76.92 ^{ab}	50.25 ^{ab}	1.67
	Average	11.53 ^A	0.464 ^{AB}	23.16	81.52	53.26	1.39
All Strains	69 EC	11.10 ^Z	0.492 ^Z	23.93	84.70	54.51	1.31
	69 ECS	11.59 ^Y	0.454 ^Y	23.03	81.42	52.36	1.21

Enrichable Cage=EC; Enriched Colony Housing System=ECS

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 46. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 EC	62.47 ^b	0	0	0.66	32.00	56.00
	69 ECS	63.08 ^{ab}	0	0	0	31.71	58.56
	Average	62.77 ^B	0	0	0.33	31.86	57.28
Shaver White	69 EC	64.52 ^{ab}	0	0	0	24.13	68.77
	69 ECS	64.61 ^{ab}	0	0	0.83	9.28	72.24
	Average	64.56 ^{AB}	0	0	0.42	16.71	70.50
Dekalb White	69 EC	63.17 ^{ab}	0	0	0	33.63	60.90
	69 ECS	63.18 ^{ab}	0	0	0	34.29	57.44
	Average	63.17 ^{AB}	0	0	0	33.96	59.17
Babcock White	69 EC	63.66 ^{ab}	0	0	0	28.26	66.74
	69 ECS	64.23 ^{ab}	0	0	0	19.77	67.44
	Average	63.94 ^{AB}	0	0	0	24.01	67.09
ISA B-400	69 EC	64.10 ^{ab}	0	0	0	26.52	67.93
	69 ECS	64.90 ^{ab}	0	0	0	13.92	74.68
	Average	64.50 ^{AB}	0	0	0	20.22	71.30
Hy-Line W-36	69 EC	63.85 ^{ab}	0	0	0	27.21	66.50
	69 ECS	62.97 ^{ab}	0	0	0	12.39	74.28
	Average	63.41 ^{AB}	0	0	0	19.80	70.39
Hy-Line CV-24	69 EC	65.18 ^{ab}	0	0	0	26.41	68.05
	69 ECS	65.61 ^{ab}	0	0	0	13.50	76.18
	Average	65.39 ^A	0	0	0	19.96	72.11
Lohmann LSL Lite	69 EC	66.29 ^a	0	0	0	5.98	85.44
	69 ECS	64.05 ^{ab}	0	0	0	21.19	67.32
	Average	65.17 ^A	0	0	0	13.58	76.38
H&N Nick Chick	69 EC	64.98 ^{ab}	0	0	0	16.40	78.24
	69 ECS	65.07 ^{ab}	0	0	0	13.33	75.00
	Average	65.02 ^{AB}	0	0	0	14.87	76.62
Novogen White	69 EC	65.33 ^{ab}	0	0	0	19.33	76.67
	69 ECS	65.48 ^{ab}	0	0	0	8.40	76.48
	Average	65.41 ^A	0	0	0	13.87	76.57
All Strains	69 EC	64.36	0	0	0.07	23.99	69.52
	69 ECS	64.31	0	0	0.08	17.78	69.96

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain* production system interactions

TABLE 47. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 EC	88.67	0	10.66	0.67	2.74 ^{ab}	1.28 ^{abc}
	69 ECS	90.27	0	8.80	0.92	2.69 ^{ab}	1.26 ^{abc}
	Average	89.47	0	9.73	0.80	2.74	1.27 ^{AB}
Shaver White	69 EC	92.90	0	6.43	0.67	2.56 ^{ab}	1.20 ^{bc}
	69 ECS	82.36	0.86	14.28	2.50	2.60 ^{ab}	1.33 ^{abc}
	Average	87.63	0.43	10.36	1.58	2.58	1.26 ^{AB}
Dekalb White	69 EC	94.53	0	4.63	0.83	2.83 ^{ab}	1.25 ^{abc}
	69 ECS	91.73	0	7.27	1.00	2.78 ^{ab}	1.24 ^{abc}
	Average	93.13	0	5.95	0.92	2.80	1.25 ^{AB}
Babcock White	69 EC	95.00	0	4.44	0.56	2.92 ^a	1.20 ^{bc}
	69 ECS	87.22	0	11.11	1.67	2.59 ^{ab}	1.44 ^a
	Average	91.11	0	7.78	1.11	2.75	1.32 ^A
ISA B-400	69 EC	94.46	0	4.21	1.33	2.89 ^a	1.23 ^{abc}
	69 ECS	88.60	1.04	8.63	1.72	2.73 ^{ab}	1.23 ^{abc}
	Average	91.53	0.52	6.42	1.53	2.81	1.23 ^{AB}
Hy-Line W-36	69 EC	93.71	0	4.70	1.59	2.58 ^{ab}	1.16 ^c
	69 ECS	86.67	0	13.33	0	2.42 ^{ab}	1.13 ^c
	Average	90.19	0	9.02	0.80	2.50	1.14 ^B
Hy-Line CV-24	69 EC	94.46	1.25	2.86	1.43	2.65 ^{ab}	1.24 ^{abc}
	69 ECS	89.68	0	9.45	0.86	2.60 ^{ab}	1.36 ^{ab}
	Average	92.07	0.62	6.15	1.14	2.63	1.30 ^A
Lohmann LSL Lite	69 EC	91.42	0	8.58	0	2.80 ^{ab}	1.34 ^{abc}
	69 ECS	88.52	0	11.48	0	2.56 ^b	1.33 ^{abc}
	Average	89.97	0	10.03	0	2.53	1.33 ^A
H&N Nick Chick	69 EC	94.64	0.56	3.65	1.15	2.77 ^{ab}	1.26 ^{abc}
	69 ECS	88.33	0	9.79	1.88	2.69 ^{ab}	1.35 ^{abc}
	Average	91.48	0.28	6.72	1.51	2.73	1.31 ^A
Novogen White	69 EC	96.00	0	4.00	0	2.85 ^a	1.26 ^{abc}
	69 ECS	84.88	2.00	11.79	1.33	2.43 ^{ab}	1.32 ^{abc}
	Average	90.44	1.00	7.90	0.67	2.64	1.29 ^A
All Strains	69 EC	93.58 ^Y	0.18	5.42 ^Y	0.82	2.76 ^Y	1.24 ^Z
	69 ECS	87.82 ^Z	0.39	10.59 ^Z	1.19	2.58 ^Z	1.30 ^Y

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 48. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA	69 EC	10.87 ^{ab}	0.454 ^{ab}	22.79	80.67	49.36	1.59
Amber	69 ECS	11.86 ^a	0.382 ^b	20.61	73.60	45.41	0
	Average	11.36	0.418	21.70	77.14	47.39	0.79
TETRA	69 EC	11.22 ^{ab}	0.427 ^{ab}	21.57	75.11	48.03	2.78
Brown	69 ECS	11.19 ^{ab}	0.420 ^{ab}	20.77	73.92	46.74	0.93
	Average	11.21	0.423	21.17	74.52	47.38	1.85
Novogen	69 EC	11.01 ^{ab}	0.458 ^{ab}	22.50	79.45	50.31	1.67
Brown	69 ECS	11.53 ^{ab}	0.436 ^{ab}	21.96	77.24	50.26	1.67
	Average	11.27	0.447	22.23	78.35	50.29	1.67
Lohmann	69 EC	10.97 ^{ab}	0.473 ^a	23.19	82.74	52.07	0.62
LB-Lite	69 ECS	11.70 ^{ab}	0.442 ^{ab}	22.71	79.98	51.61	2.08
	Average	11.33	0.458	22.95	81.36	51.84	1.35
Hy-Line	69 EC	11.54 ^{ab}	0.447 ^{ab}	23.52	83.98	51.56	0
Silver Brown	69 ECS	11.75 ^{ab}	0.425 ^{ab}	22.87	81.68	49.88	0
	Average	11.64	0.436	23.19	82.83	50.72	0
Hy-Line	69 EC	10.33 ^b	0.455 ^{ab}	21.14	75.20	47.16	0.93
Brown	69 ECS	11.54 ^{ab}	0.414 ^{ab}	21.08	75.28	47.49	0
	Average	10.94	0.434	21.11	75.24	47.52	0.46
ISA	69 EC	10.68 ^{ab}	0.462 ^a	22.18	77.83	49.31	2.32
Brown	69 ECS	10.93 ^{ab}	0.464 ^a	22.91	81.46	50.72	0.56
	Average	10.80	0.463	22.54	79.65	50.02	1.44
Bovans	69 EC	11.48 ^{ab}	0.446 ^{ab}	22.38	79.78	50.93	0.79
Brown	69 ECS	11.86 ^a	0.438 ^{ab}	22.93	81.76	52.00	0.56
	Average	11.67	0.442	22.66	80.77	51.46	0.68
All	69 EC	11.01 ^Z	0.453 ^Z	22.41	79.35	49.84	1.34
Strains	69 ECS	11.54 ^Y	0.428 ^Y	21.98	78.12	49.31	0.72

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ab - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 49. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 EC	61.16 ^{cd}	0	0	1.12	49.85 ^a	44.23 ^b
	69 ECS	61.70 ^{bcd}	0	0	2.76	32.62 ^{ab}	55.08 ^{ab}
	Average	61.43 ^{BC}	0	0	1.94 ^A	41.23 ^{AB}	49.65 ^B
TETRA Brown	69 EC	63.96 ^{abcd}	0	0	0	22.67 ^{ab}	66.32 ^{ab}
	69 ECS	63.21 ^{abcd}	0	0	0	20.68 ^{ab}	62.92 ^{ab}
	Average	63.59 ^A	0	0	0 ^B	21.68 ^B	64.64 ^{AB}
Novogen Brown	69 EC	63.40 ^{abcd}	0	0	0	33.32 ^{ab}	64.58 ^{ab}
	69 ECS	65.05 ^a	0	0	0.71	12.32 ^b	74.63 ^a
	Average	64.23 ^A	0	0	0.36 ^{AB}	22.82 ^B	69.61 ^A
Lohmann LB-Lite	69 EC	62.93 ^{abcd}	0	0	0.46	33.40 ^{ab}	62.02 ^{ab}
	69 ECS	64.54 ^{ab}	0	0	0	18.18 ^{ab}	69.48 ^{ab}
	Average	63.73 ^A	0	0	0.28 ^{AB}	25.79 ^B	65.77 ^{AB}
Hy-Line Silver Brown	69 EC	61.39 ^{bcd}	0	0	0.83	49.84 ^a	46.82 ^{ab}
	69 ECS	61.10 ^{cd}	0	0	0	46.88 ^a	48.75 ^{ab}
	Average	61.25 ^C	0	0	0.42 ^{AB}	48.36 ^A	47.79 ^B
Hy-Line Brown	69 EC	62.72 ^{abcd}	0	0	0	35.37 ^{ab}	57.72 ^{ab}
	69 ECS	63.55 ^{abcd}	0	0	0	21.20 ^{ab}	67.49 ^{ab}
	Average	63.14 ^{AB}	0	0	0 ^B	28.29 ^{AB}	62.60 ^{AB}
ISA Brown	69 EC	63.36 ^{abcd}	0	0	0	34.47 ^{ab}	63.23 ^{ab}
	69 ECS	62.28 ^{abcd}	0	0	0	38.06 ^{ab}	50.48 ^{ab}
	Average	62.82 ^{ABC}	0	0	0 ^B	36.26 ^{AB}	56.86 ^{AB}
Bovans Brown	69 EC	63.87 ^{abcd}	0	0	0	33.25 ^{ab}	64.28 ^{ab}
	69 ECS	63.60 ^{abcd}	0	0	0	24.12 ^{ab}	63.00 ^{ab}
	Average	63.74 ^A	0	0	0 ^B	28.68 ^{AB}	63.64 ^{AB}
All Strains	69 EC	62.85	0	0	0.31	36.25 ^Y	58.66
	69 ECS	63.13	0	0	0.43	26.76 ^Z	61.48

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 50. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 EC	95.20	0	4.80	0	2.65	1.22
	69 ECS	90.45	0	8.59	0.96	2.32	1.33
	Average	92.82	0	6.69	0.48	2.48 ^{AB}	1.27
TETRA Brown	69 EC	89.00	0	11.00	0	2.44	1.26
	69 ECS	83.64	0	16.36	0	2.29	1.25
	Average	86.32	0	13.68	0	2.37 ^B	1.25
Novogen Brown	69 EC	97.91	1.33	2.09	0	2.67	1.23
	69 ECS	87.67	0	11.00	0	2.47	1.29
	Average	92.79	0.67	6.55	0	2.57 ^{AB}	1.26
Lohmann LB-Lite	69 EC	96.02	0.64	3.34	0	2.72	1.23
	69 ECS	87.66	0	12.34	0	2.56	1.31
	Average	91.84	0.32	7.84	0	2.64 ^A	1.27
Hy-Line Silver Brown	69 EC	97.50	0	2.50	0	2.76	1.29
	69 ECS	95.62	0	4.38	0	2.67	1.32
	Average	96.56	0	3.44	0	2.72 ^A	1.30
Hy-Line Brown	69 EC	93.09	0	4.01	2.90	2.40	1.16
	69 ECS	88.70	0	11.30	0	2.38	1.29
	Average	90.89	0	7.66	1.45	2.39 ^B	1.22
ISA Brown	69 EC	97.70	0.56	1.74	0	2.62	1.20
	69 ECS	88.54	0	10.79	0.66	2.57	1.22
	Average	93.12	0.28	6.27	0.33	2.60 ^{AB}	1.21
Bovans Brown	69 EC	97.53	0	1.97	0.49	2.64	1.28
	69 ECS	87.12	0	9.48	3.41	2.52	1.33
	Average	92.33	0	5.72	1.95	2.58 ^{AB}	1.30
All Strains	69 EC	95.49 ^Y	0.15	3.93 ^Y	0.42	2.61 ^Y	1.23 ^Z
	69 ECS	88.67 ^Z	0.17	10.53 ^Z	0.63	2.47 ^Z	1.29 ^Y

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 51. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder (Strain)	Production System	Feed Consumption (kg/100/hen/d)	Feed Conversion (g egg/g feed)	Eggs Per Bird Housed	Egg Production (HD%)	Egg Mass (g/HD)	Mortality*
Bovans White	69 EC	116.59 ^{bcde}	0.42 ^{abcde}	193.27	75.46 ^{ab}	48.99 ^{abc}	11.68
	69 ECS	119.53 ^{abcde}	0.40 ^{abcdef}	188.90	73.81 ^{abcd}	48.04 ^{abc}	6.95
	Average	118.06 ^{ABC}	0.41	191.09 ^A	74.63 ^A	48.51 ^A	9.31
Shaver White	69 EC	113.85 ^{def}	0.41 ^{abcdef}	177.57	68.54 ^{cdef}	45.23 ^{cde} 47.08 ^{abc}	20.84
	69 ECS	120.59 ^{abcde}	0.39 ^{abcdef}	183.66	71.19 ^{abcde}	d	12.51
	Average	117.22 ^{ABC}	0.40	180.61 ^{AB}	69.86 ^A	46.16 ^A	16.67
Dekalb White	69 EC	116.11 ^{bcde}	0.43 ^{abc}	194.69	76.08 ^a	49.38 ^{abc} 46.62 ^{abc}	7.41
	69 ECS	114.09 ^{cdef}	0.41 ^{abcdef}	185.22	72.07 ^{abcd}	d	13.90
	Average	115.10 ^{BC}	0.42	189.95 ^A	74.08 ^A	48.00 ^A	10.65
Babcock White	69 EC	115.00 ^{cdef}	0.44 ^a	194.72	76.00 ^a	49.92 ^{ab}	10.04
	69 ECS	126.39 ^{ab}	0.39 ^{cdef}	186.77	72.73 ^{abcd}	48.21 ^{abc}	17.37
	Average	120.69 ^{AB}	0.42	190.75 ^A	74.37 ^A	49.06 ^A	13.70
ISA B-400	69 EC	111.00 ^{efg}	0.43 ^{abc}	187.39	72.87 ^{abcd}	47.37 ^{abc}	11.90
	69 ECS	113.90 ^{def}	0.42 ^{abcde}	187.04	72.77 ^{abcd}	47.71 ^{abc}	13.61
	Average	112.35 ^C	0.42	187.21 ^A	72.82 ^A	47.54 ^A	12.75
Hy-Line W-36	69 EC	103.20 ^g	0.42 ^{abcdef}	156.24	64.68 ^{ef}	42.71 ^{de}	8.50
	69 ECS	103.75 ^{fg}	0.39 ^{bcdef}	165.18	61.11 ^f	40.12 ^e	9.26
	Average	103.48 ^D	0.40	160.71 ^B	62.90 ^B	41.41 ^B	8.88
Hy-Line CV-24	69 EC	117.60 ^{abde}	0.43 ^{ab}	192.71	75.14 ^{abc}	50.21 ^{ab} 45.49 ^{bcd}	15.56
	69 ECS	116.74 ^{bcde}	0.39 ^{bcdef}	171.74	67.08 ^{def}	e	11.11
	Average	117.17 ^{ABC}	0.41	182.22 ^{AB}	71.10 ^A	47.85 ^A	13.34
Lohmann LSL Lite	69 EC	118.14 ^{abcde}	0.43 ^{abcd}	190.51	74.35 ^{abcd}	49.63 ^{abc}	11.11
	69 ECS	127.83 ^a	0.37 ^f	180.36	70.17 ^{abcde}	47.37 ^{abc}	18.06
	Average	122.99 ^A	0.40	185.44 ^{AB}	72.26 ^A	48.50 ^A	14.59
H&N Nick Chick	69 EC	117.21 ^{bcde}	0.42 ^{abcde}	184.76	71.93 ^{abcd}	48.77 ^{abc}	12.51
	69 ECS	122.37 ^{abcd}	0.38 ^{ef}	176.32	68.54 ^{bcdef}	45.73 ^{bcd}	13.20
	Average	119.79 ^{AB}	0.40	180.54 ^{AB}	70.23 ^A	47.25 ^A	12.86
Novogen White	69 EC	119.80 ^{abcde}	0.42 ^{abcd}	194.34	75.43 ^{abc}	50.66 ^a	15.57
	69 ECS	124.85 ^{abc}	0.38 ^{def}	182.65	70.25 ^{abcde}	47.51 ^{abc}	16.12
	Average	122.33 ^A	0.40	188.50 ^A	72.84 ^A	49.08 ^A	15.84
All Strains	69 EC	114.83 ^Y	0.42 ^Z	187.51	73.05 ^Y	48.29 ^Y	12.68
	69 ECS	119.05 ^Z	0.39 ^Y	179.89	69.97 ^Z	46.39 ^Z	13.03

Enrichable Cage=EC; Enriched Colony Housing System=ECS

ABCD. - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain*production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 52. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder (Strain)	Production System	Egg Weight (g/egg)	Pee Wee (%)	Small (%)	Medium (%)	Large (%)	Extra Large (%)
Bovans White	69 EC	64.99 ^f	0	0	0.27	9.76 ^{ab}	85.96 ^{ab}
	69 ECS	65.11 ^{ef}	0	0.14	0.53	9.92 ^{ab}	83.61 ^{ab}
	Average	65.05 ^{DE}	0	0.07	0.40	9.83 ^{AB}	84.78 ^{BC}
Shaver White	69 EC	66.02 ^{cdef}	0	0	0.17	6.87 ^{ab}	89.07 ^{ab}
	69 ECS	66.22 ^{bcdef}	0	0	0	5.69 ^{ab}	85.33 ^{ab}
	Average	66.12 ^{BC}	0	0	0.08	6.28 ^{ABC}	87.20 ^{ABC}
Dekalb White	69 EC	64.97 ^f	0	0	0.20	11.02 ^{ab}	86.29 ^{ab}
	69 ECS	64.76 ^f	0	0.14	0	11.94 ^a	81.94 ^b
	Average	64.86 ^E	0	0.07	0.10	11.48 ^A	84.12 ^C
Babcock White	69 EC	65.83 ^{cdef}	0	0	0.26	7.59 ^{ab}	87.00 ^{ab}
	69 ECS	66.31 ^{bcdef}	0	0	0.11	5.86 ^{ab}	87.30 ^{ab}
	Average	66.07 ^{BCD}	0	0	0.18	6.73 ^{ABC}	87.15 ^{ABC}
ISA B-400	69 EC	65.06 ^f	0	0.08	0.20	8.24 ^{ab}	88.09 ^{ab}
	69 ECS	65.57 ^{def}	0	0	0.14	6.50 ^{ab}	85.64 ^{ab}
	Average	65.32 ^{CDE}	0	0.04	0.17	7.37 ^{ABC}	86.86 ^{ABC}
Hy-Line W-36	69 EC	66.10 ^{cdef}	0	0	0	7.59 ^{ab}	89.17 ^{ab}
	69 ECS	65.76 ^{cdef}	0	0	0.18	9.29 ^{ab}	84.48 ^{ab}
	Average	65.93 ^{CDE}	0	0	0.09	8.44 ^{ABC}	86.82 ^{ABC}
Hy-Line CV-24	69 EC	66.85 ^{abcd}	0	0	0	4.71 ^{ab}	92.60 ^a
	69 ECS	67.88 ^{ab}	0	0	0	5.83 ^{ab}	89.25 ^{ab}
	Average	67.36 ^A	0	0	0	5.27 ^{BC}	90.92 ^{AB}
Lohmann LSL Lite	69 EC	66.87 ^{abcd}	0	0	0	4.50 ^{ab}	90.58 ^{ab}
	69 ECS	67.39 ^{abc}	0	0	0.17	6.78 ^{ab}	87.25 ^{ab}
	Average	67.13 ^{AB}	0	0	0.08	5.64 ^{BC}	88.92 ^{ABC}
H&N Nick Chick	69 EC	67.91 ^a	0	0	0.41	4.17 ^{ab}	92.44 ^a
	69 ECS	66.79 ^{abcde}	0	0	0.14	5.17 ^{ab}	87.19 ^{ab}
	Average	67.35 ^A	0	0	0.27	4.67 ^{BC}	89.82 ^{ABC}
Novogen White	69 EC	67.22 ^{abc}	0	0	0	5.07 ^{ab}	90.53 ^{ab}
	69 ECS	67.64 ^{ab}	0	0	0	3.07 ^b	92.91 ^a
	Average	67.45 ^A	0	0	0	4.07 ^C	91.72 ^A
All Strains	69 EC	66.18	0	0.01	0.15	6.95	89.17 ^Y
	69 ECS	66.35	0	0.03	0.13	7.00	86.49 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdef - Different letters denote significant differences (P<.01) in the strain* production system interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 53. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 EC	92.52 ^{ab}	0.40	5.78 ^c	1.30	21.13	10.86
	69 ECS	89.82 ^{abc}	0.31	8.65 ^{bc}	1.22	21.07	11.11
	Average	91.17 ^{AB}	0.36	7.21 ^{BC}	1.26	21.10	10.98 ^A
Shaver White	69 EC	91.52 ^{ab}	0.33	7.71 ^c	0.44	21.15	10.58
	69 ECS	80.58 ^d	0.36	16.79 ^a	2.28	20.82	11.24
	Average	86.05 ^C	0.34	12.25 ^A	1.36	20.98	10.91 ^A
Dekalb White	69 EC	95.46 ^a	0.78	3.54 ^c	0.22	22.24	10.81
	69 ECS	88.55 ^{abc}	0.41	9.50 ^{bc}	1.55	21.74	10.60
	Average	92.00 ^{AB}	0.60	6.52 ^{BC}	0.88	21.99	10.70 ^{AB}
Babcock White	69 EC	92.22 ^{ab}	0.26	6.56 ^c	0.95	21.20	10.69
	69 ECS	87.79 ^{bcd}	0.42	9.58 ^{bc}	2.22	21.16	11.82
	Average	90.00 ^{ABC}	0.34	8.07 ^{BC}	1.58	21.18	11.25 ^A
ISA B-400	69 EC	93.76 ^{ab}	0.65	5.10 ^c	0.49	20.15	10.32
	69 ECS	83.12 ^{cd}	0.29	14.79 ^{ab}	1.81	18.35	10.60
	Average	88.44 ^{BC}	0.47	9.94 ^{AB}	1.15	19.25	10.46 ^{AB}
Hy-Line W-36	69 EC	93.42 ^{ab}	0.15	4.81 ^c	1.60	20.62	9.58
	69 ECS	88.13 ^{abcd}	0.40	10.00 ^{abc}	1.47	20.47	9.67
	Average	90.78 ^{ABC}	0.28	7.41 ^{BC}	1.54	20.54	9.62 ^B
Hy-Line CV-24	69 EC	95.49 ^a	0	4.08 ^c	0.43	22.24	10.96
	69 ECS	91.61 ^{ab}	0.33	6.80 ^c	1.26	20.40	10.90
	Average	93.55 ^A	0.16	5.44 ^C	0.84	21.32	10.93 ^A
Lohmann LSL Lite	69 EC	90.60 ^{abc}	0.91	7.08 ^c	1.41	22.06	10.99
	69 ECS	88.68 ^{abc}	0.20	9.60 ^{bc}	1.52	21.32	11.90
	Average	89.64 ^{ABC}	0.56	8.34 ^{ABC}	1.47	21.69	11.44 ^A
H&N Nick Chick	69 EC	92.99 ^{ab}	1.21	4.92 ^c	0.88	19.57	10.92
	69 ECS	87.05 ^{bcd}	0.23	9.84 ^{bc}	2.88	22.66	11.41
	Average	90.01 ^{ABC}	0.72	7.38 ^{BC}	1.88	21.12	11.17 ^A
Novogen White	69 EC	92.10 ^{ab}	0.73	5.60 ^c	1.58	19.69	11.14
	69 ECS	91.54 ^{ab}	0.79	7.08 ^c	0.59	15.61	11.62
	Average	91.82 ^{AB}	0.76	6.34 ^{BC}	1.08	17.65	11.38 ^A
All Strains	69 EC	93.00 ^Y	0.54	5.52	0.93 ^Y	21.00	10.69 ^Y
	69 ECS	87.69 ^Z	0.37	10.26	1.68 ^Z	20.36	11.09 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 54. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder (Strain)	Production	Feed	Feed	Eggs Per Bird	Egg Produc- tion	Egg Mass	Mortali- ty
	System	Consumption	Conversion	Housed	(HD%)	(g/HD)	(%)
TETRA Amber	69 EC	113.29 ^{cde}	0.35 ^{bcdef}	162.70 ^{ab}	63.44 ^{abc}	40.06 ^{bcd}	11.12
	69 ECS	123.36 ^{ab}	0.29 ^g	146.45 ^b	56.84 ^c	36.37 ^d	15.98
	Average	118.32 ^{ABC}	0.32 ^E	154.58 ^B	60.14 ^C	38.21 ^C	13.55
TETRA Brown	69 EC	117.30 ^{bcde}	0.34 ^{defg}	157.51 ^{ab}	60.85 ^{abc}	39.46 ^{bcd}	18.53
	69 ECS	124.54 ^{ab}	0.32 ^{efg}	156.54 ^{ab}	60.94 ^{abc}	39.88 ^{bcd}	14.82
	Average	120.92 ^{ABC}	0.33 ^{DE}	157.03 ^{AB}	60.89 ^{BC}	39.67 ^{BC}	16.68
Novogen Brown	69 EC	109.19 ^e	0.39 ^{ab}	164.10 ^{ab}	64.28 ^{abc}	42.40 ^{abc}	5.56
	69 ECS	122.47 ^{ab}	0.34 ^{cdef}	159.21 ^{ab}	62.07 ^{abc}	41.72 ^{abcd}	15.56
	Average	115.83 ^{BC}	0.36 ^{ABC}	161.66 ^{AB}	63.17 ^{ABC}	42.06 ^{AB}	10.56
Lohmann LB-Lite	69 EC	110.48 ^e	0.42 ^a	178.23 ^a	69.58 ^a	45.86 ^a	12.50
	69 ECS	121.95 ^{abc}	0.35 ^{bcdef}	164.92 ^{ab}	64.12 ^{abc}	42.61 ^{abc}	20.85
	Average	116.21 ^{BC}	0.38 ^A	171.57 ^{AB}	66.85 ^{AB}	44.23 ^A	16.67
Hy-Line Silver Brown	69 EC	118.69 ^{abcde}	0.36 ^{bcdef}	169.89 ^{ab}	66.51 ^{ab}	42.25 ^{abcd}	7.18
	69 ECS	127.88 ^a	0.32 ^{fg}	166.26 ^{ab}	64.84 ^{abc}	41.18 ^{abcd}	14.59
	Average	123.29 ^A	0.34 ^{CDE}	168.07 ^{AB}	65.68 ^{ABC}	41.71 ^{ABC}	10.88
Hy-Line Brown	69 EC	110.86 ^{de}	0.37 ^{bcd}	158.88 ^{ab}	62.30 ^{abc}	40.93 ^{abcd}	5.10
	69 ECS	120.13 ^{abc}	0.34 ^{def}	155.59 ^{ab}	60.77 ^{bc}	40.58 ^{bcd}	13.34
	Average	115.50 ^C	0.35 ^{BCD}	157.23 ^{AB}	61.53 ^{BC}	40.76 ^{BC}	9.22
ISA Brown	69 EC	117.59 ^{bcde}	0.37 ^{bcd}	168.99 ^{ab}	65.62 ^{ab}	43.17 ^{abc}	14.36
	69 ECS	116.22 ^{bcde}	0.36 ^{bcdef}	162.35 ^{ab}	63.37 ^{abc}	41.38 ^{abcd}	9.95
	Average	116.90 ^{BC}	0.36 ^{ABC}	165.67 ^{AB}	64.49 ^{ABC}	42.27 ^{AB}	12.15
Bovans Brown	69 EC	118.78 ^{abcd}	0.38 ^{bc}	173.68 ^{ab}	67.84 ^{ab}	44.85 ^{ab}	10.71
	69 ECS	123.45 ^{ab}	0.36 ^{bcde}	174.14 ^{ab}	67.98 ^{ab}	44.93 ^{ab}	9.45
	Average	121.12 ^{AB}	0.37 ^{AB}	173.91 ^A	67.91 ^A	44.89 ^A	10.08
All Strains	69 EC	114.52 ^Z	0.37 ^Z	166.75	65.05 ^Y	42.37	10.63
	69 ECS	122.50 ^Y	0.34 ^Y	160.68	62.61 ^Z	41.01	14.32

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 55. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 EC	63.23 ^f	0	0	0.54 ^{ab}	19.46 ^a	77.27 ^c
	69 ECS	64.09 ^{def}	0	0	0.19 ^b	14.50 ^{ab}	80.36 ^{abc}
	Average	63.66 ^C	0	0	0.37	16.98 ^A	78.81 ^C
TETRA Brown	69 EC	64.92 ^{cdef}	0	0	0.74 ^{ab}	15.52 ^{ab}	82.22 ^{abc}
	69 ECS	65.49 ^{abcd}	0	0	0 ^b	11.59 ^{abcd}	84.74 ^{abc}
	Average	65.20 ^B	0	0	0.37	13.56 ^{ABC}	83.48 ^{ABC}
Novogen Brown	69 EC	66.02 ^{abc}	0	0	0.42 ^b	9.78 ^{bcd}	87.96 ^{ab}
	69 ECS	67.26 ^a	0	0	0.16 ^b	4.84 ^{cd}	89.29 ^a
	Average	66.64 ^A	0	0	0.29	7.31 ^{CD}	88.62 ^A
Lohmann LB-Lite	69 EC	65.97 ^{abc}	0	0	0.44 ^b	8.42 ^{bcd}	89.07 ^a
	69 ECS	66.49 ^{abc}	0	0	0 ^b	4.25 ^d	89.58 ^a
	Average	66.23 ^{AB}	0	0	0.22	6.34 ^D	89.33 ^A
Hy-Line Silver Brown	69 EC	63.64 ^{ef}	0	0	1.86 ^a	16.89 ^{ab}	78.36 ^{bc}
	69 ECS	63.60 ^{ef}	0	0	0.11 ^b	13.81 ^{abc}	81.14 ^{abc}
	Average	63.62 ^C	0	0	0.99	15.35 ^{AB}	79.75 ^{BC}
Hy-Line Brown	69 EC	65.75 ^{abc}	0	0.11	0.41 ^b	10.76 ^{bcd}	85.87 ^{abc}
	69 ECS	66.85 ^{ab}	0	0	0.44 ^b	4.29 ^d	90.04 ^a
	Average	66.30 ^{AB}	0	0.06	0.42	7.52 ^{CD}	87.96 ^A
ISA Brown	69 EC	65.85 ^{abc}	0	0	0.20 ^b	11.67 ^{abcd}	84.83 ^{abc}
	69 ECS	65.42 ^{bcd}	0	0	0.33 ^b	9.07 ^{bcd}	86.11 ^{abc}
	Average	65.63 ^B	0	0	0.27	10.37 ^{BCD}	85.47 ^{AB}
Bovans Brown	69 EC	66.10 ^{abc}	0	0	0.46 ^b	9.86 ^{bcd}	85.92 ^{abc}
	69 ECS	66.23 ^{bcd}	0	0.09	0.07 ^b	7.87 ^{bcd}	87.78 ^{ab}
	Average	66.16 ^{AB}	0	0.04	0.26	8.86 ^{CD}	86.35 ^{AB}
All Strains	69 EC	65.18 ^Z	0	0.01	0.63 ^Y	12.79 ^Y	83.94
	69 ECS	65.67 ^Y	0	0.01	0.16 ^Z	8.78 ^Z	86.00

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdef - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 56. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 EC	95.48 ^{ab}	0.51	3.29 ^d	0.73	16.08	9.77
	69 ECS	91.46 ^{abcd}	0.31	6.61 ^{abcd}	1.63	17.48	10.64
	Average	93.47	0.41	4.95	1.18	16.78	10.21 ^{AB}
TETRA Brown	69 EC	95.18 ^{abc}	0.43	3.85 ^{cd}	0.53	19.35	10.11
	69 ECS	92.22 ^{abcd}	0.64	6.35 ^{abcd}	0.78	17.87	10.76
	Average	93.70	0.54	5.10	0.66	18.61	10.44 ^{AB}
Novogen Brown	69 EC	95.92 ^{ab}	0.49	3.14 ^d	0.46	16.53	9.43
	69 ECS	89.87 ^{cd}	1.00	8.04 ^{abc}	1.08	19.94	10.57
	Average	92.89	0.75	5.59	0.77	18.24	10.00 ^B
Lohmann LB-Lite	69 EC	96.38 ^a	0.40	2.47 ^d	0.75	19.70	9.52
	69 ECS	88.57 ^d	0.10	10.12 ^a	1.21	18.80	10.51
	Average	92.47	0.25	6.30	0.98	19.25	10.01 ^B
Hy-Line Silver Brown	69 EC	94.81 ^{abc}	0.71	3.46 ^{cd}	1.02	18.30	10.25
	69 ECS	89.94 ^{cd}	1.43	7.30 ^{abcd}	1.34	18.10	11.03
	Average	92.38	1.07	5.38	1.18	18.20	10.64 ^A
Hy-Line Brown	69 EC	93.41 ^{abcd}	0.80	4.36 ^{cd}	1.42	17.21	9.58
	69 ECS	90.86 ^{bcd}	0.63	8.09 ^{abc}	0.42	17.09	10.36
	Average	92.14	0.72	6.23	0.92	17.15	9.97 ^B
ISA Brown	69 EC	94.46 ^{abc}	0.69	3.40 ^{cd}	0.85	20.57	10.14
	69 ECS	91.78 ^{abcd}	0.87	6.68 ^{abcd}	0.68	19.46	10.03
	Average	93.12	0.78	5.34	0.77	20.01	10.08 ^{AB}
Bovans Brown	69 EC	93.30 ^{abcd}	0.15	5.81 ^{abcd}	0.75	20.45	10.24
	69 ECS	89.75 ^{cd}	0.10	9.04 ^{ab}	1.11	18.80	10.66
	Average	91.52	0.12	7.42	0.93	19.62	10.45 ^{AB}
All Strains	69 EC	94.87 ^Y	0.52	7.78 ^Z	0.82	18.52	9.88 ^Y
	69 ECS	90.56 ^Z	0.64	3.80 ^Y	1.03	18.44	10.57 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 57. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality*
(Strain)		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans White	69 EC	108.05 ^{abcde}	0.46	525.14	81.48	50.19	22.23 ^{ab}
	69 ECS	109.98 ^{abcd}	0.44	533.46	79.76	49.29	28.48 ^{ab}
	Average	109.02 ^A	0.45	529.30 ^A	80.62 ^{AB}	49.74 ^{AB}	25.35
Shaver White	69 EC	105.80 ^{cde}	0.46	487.14	77.61	48.68	38.43 ^{ab}
	69 ECS	110.89 ^{abc}	0.45	521.83	79.84	50.10	46.53 ^a
	Average	108.35 ^A	0.45	504.48 ^{AB}	78.72 ^{AB}	49.39 ^{AB}	42.48 ^A
Dekalb White	69 EC	109.11 ^{abcd}	0.46	515.19	81.49	50.63	15.18 ^{ab}
	69 ECS	107.73 ^{abcde}	0.47	518.35	80.18	49.75	24.31 ^{ab}
	Average	108.42 ^A	0.46	516.77 ^{AB}	80.83 ^{AB}	50.19 ^A	19.74 ^{BC}
Babcock White	69 EC	107.13 ^{bcde}	0.48	514.74	82.35	51.84	14.97 ^{ab}
	69 ECS	116.74 ^a	0.44	526.34	82.66	51.83	31.26 ^{ab}
	Average	111.93 ^A	0.46	520.54 ^{AB}	82.50 ^A	51.84 ^A	23.11 ^{ABC}
ISA B-400	69 EC	107.22 ^{bcde}	0.48	508.76	82.22	51.89	19.56 ^{ab}
	69 ECS	108.86 ^{abcde}	0.47	516.34	82.32	52.05	30.08 ^{ab}
	Average	108.04 ^A	0.47	512.55 ^{AB}	82.27 ^A	51.97 ^A	24.82 ^{ABC}
Hy-Line W-36	69 EC	100.80 ^e	0.45	484.00	74.99	46.52	11.63 ^b
	69 ECS	100.06 ^e	0.44	484.09	73.50	45.06	13.89 ^{ab}
	Average	100.43 ^B	0.45	484.04 ^B	74.24 ^B	45.79 ^B	12.76 ^C
Hy-Line CV-24	69 EC	110.37 ^{abc}	0.45	512.42	80.52	50.61	25.56 ^{ab}
	69 ECS	111.20 ^{abc}	0.43	511.59	75.67	47.92	22.92 ^{ab}
	Average	110.79 ^A	0.44	512.00 ^{AB}	78.10 ^{AB}	49.26 ^{AB}	24.24 ^{ABC}
Lohmann LSL Lite	69 EC	110.12 ^{abcd}	0.46	507.74	80.54	50.96	29.17 ^{ab}
	69 ECS	114.72 ^{ab}	0.43	513.02	78.51	49.68	31.25 ^{ab}
	Average	112.42 ^A	0.44	510.38 ^{AB}	79.53 ^{AB}	50.32 ^A	30.21 ^{ABC}
H&N Nick Chick	69 EC	109.21 ^{abcd}	0.46	499.53	79.45	50.62	22.23 ^{ab}
	69 ECS	112.54 ^{abc}	0.44	508.44	78.23	49.47	29.87 ^{ab}
	Average	110.87 ^A	0.45	504.00 ^{AB}	78.84 ^{AB}	50.04 ^A	26.05 ^{ABC}
Novogen White	69 EC	109.59 ^{abcd}	0.47	513.88	81.07	51.84	27.78 ^{ab}
	69 ECS	115.62 ^a	0.43	512.91	77.66	49.55	38.33 ^{ab}
	Average	112.61 ^A	0.45	513.39 ^{AB}	79.36 ^{AB}	50.70 ^A	33.06 ^{AB}
All Strains	69 EC	107.74 ^Y	0.46 ^Z	506.86	80.17	50.38	22.67 ^Y
	69 ECS	110.83 ^Z	0.44 ^Y	514.64	78.83	49.47	29.69 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS

ABC. - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 58. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 EC	61.11	0.87	3.37	8.27	20.14	65.47
	69 ECS	61.60	0	2.89	8.14	19.76	66.79
	Average	61.36 ^B	0.43	3.13	8.20	19.95	66.13
Shaver White	69 EC	62.35	0.02	3.55	5.90	17.98	70.80
	69 ECS	62.51	0	2.63	6.78	16.96	69.75
	Average	62.43 ^{AB}	0.01	3.09	6.34	17.47	70.26
Dekalb White	69 EC	61.66	0	3.54	6.49	19.53	69.26
	69 ECS	61.49	0	3.99	6.00	19.93	67.46
	Average	61.58 ^B	0	3.77	6.24	19.73	68.36
Babcock White	69 EC	62.61	0	3.01	6.37	16.98	71.40
	69 ECS	62.45	0.30	2.92	6.75	16.96	70.15
	Average	62.53 ^{AB}	0.15	2.97	6.56	16.97	70.77
ISA B-400	69 EC	62.66	0.02	3.03	5.04	16.71	73.64
	69 ECS	62.85	0	2.47	5.28	16.17	72.71
	Average	62.75 ^{AB}	0.01	2.75	5.16	16.44	73.18
Hy-Line W-36	69 EC	61.82	0.72	3.61	8.27	17.09	68.83
	69 ECS	61.26	0	4.70	9.74	19.19	63.71
	Average	61.54 ^B	0.36	4.15	9.00	18.14	66.27
Hy-Line CV-24	69 EC	62.36	0	3.30	7.73	15.72	72.15
	69 ECS	63.15	1.10	3.59	6.31	16.24	70.50
	Average	62.76 ^{AB}	0.55	3.45	7.02	15.98	71.33
Lohmann LSL Lite	69 EC	62.90	0	3.54	6.76	15.01	72.78
	69 ECS	62.88	0	2.87	6.18	17.63	70.82
	Average	62.89 ^{AB}	0	3.20	6.47	16.32	71.80
H&N Nick Chick	69 EC	63.30	0	3.46	5.64	15.41	74.14
	69 ECS	62.95	0	3.02	6.27	15.67	71.76
	Average	63.12 ^{AB}	0	3.24	5.95	15.54	72.95
Novogen White	69 EC	63.38	0	2.96	5.53	15.30	74.38
	69 ECS	63.54	0	2.99	6.09	14.71	74.34
	Average	63.46 ^A	0	2.98	5.81	15.00	74.36
All Strains	69 EC	62.41	0.16	3.34	6.60	16.99	71.28
	69 ECS	62.47	0.14	3.21	6.75	17.32	69.80

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 59. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 EC	94.73 ^{abcd}	0.48	3.87	0.90	60.17 ^{abc}	27.55 ^{abcd}
	69 ECS	92.92 ^{abcdef}	0.41	5.89	0.77	58.86 ^{abc}	28.00 ^{abcd}
	Average	93.83 ^A	0.44	4.88 ^B	0.84	59.51 ^A	27.78 ^A
Shaver White	69 EC	93.88 ^{abcde}	0.35	5.32	0.47	57.96 ^{abc}	26.97 ^{bcd}
	69 ECS	87.31 ^g	0.41	10.73	1.50	58.20 ^{abc}	28.28 ^{abcd}
	Average	90.60 ^B	0.38	8.02 ^A	0.98	58.08 ^{AB}	27.63 ^A
Dekalb White	69 EC	96.38 ^a	0.52	2.68	0.47	61.22 ^{ab}	27.85 ^{abcd}
	69 ECS	90.91 ^{defg}	0.46	7.66	0.99	58.86 ^{abc}	27.52 ^{abcd}
	Average	93.64 ^A	0.49	5.17 ^B	0.73	60.04 ^A	27.68 ^A
Babcock White	69 EC	94.76 ^{abcd}	0.46	3.96	0.77	61.33 ^a	27.31 ^{abcd}
	69 ECS	90.99 ^{defg}	0.62	7.20	1.24	60.28 ^{abc}	29.82 ^a
	Average	92.88 ^{AB}	0.54	5.58 ^B	1.00	60.81 ^A	28.56 ^A
ISA B-400	69 EC	94.84 ^{abcd}	0.43	4.09	0.64	61.71 ^a	27.46 ^{abcd}
	69 ECS	88.74 ^{fg}	0.40	9.82	1.10	60.13 ^{abc}	27.80 ^{abcd}
	Average	91.79 ^{AB}	0.42	6.95 ^{AB}	0.87	60.92 ^A	27.63 ^A
Hy-Line W-36	69 EC	95.12 ^{abc}	0.36	3.60	0.90	55.35 ^{bc}	25.87 ^d
	69 ECS	90.98 ^{defg}	0.36	7.65	1.03	52.88 ^c	25.60 ^d
	Average	93.05 ^{AB}	0.36	5.65 ^{AB}	0.97	54.12 ^B	25.74 ^B
Hy-Line CV-24	69 EC	95.90 ^{ab}	0.31	3.50	0.31	60.49 ^{ab}	28.41 ^{abcd}
	69 ECS	92.33 ^{abcdef}	0.33	6.49	0.87	55.37 ^{abc}	28.40 ^{abcd}
	Average	94.12 ^A	0.32	5.00 ^B	0.59	57.93 ^{AB}	28.40 ^A
Lohmann LSL Lite	69 EC	94.39 ^{abcde}	0.59	4.40	0.63	60.17 ^{abc}	28.09 ^{abcd}
	69 ECS	90.51 ^{efg}	0.72	7.88	0.89	57.83 ^{abc}	29.19 ^{ab}
	Average	92.45 ^{AB}	0.65	6.14 ^{AB}	0.76	59.00 ^A	28.64 ^A
H&N Nick Chick	69 EC	94.68 ^{abcd}	0.98	3.70	0.62	59.58 ^{abc}	27.86 ^{abcd}
	69 ECS	90.76 ^{defg}	0.46	7.10	1.66	57.38 ^{abc}	28.70 ^{abc}
	Average	92.72 ^{AB}	0.72	5.40 ^B	1.14	58.47 ^{AB}	28.28 ^A
Novogen White	69 EC	94.76 ^{abcd}	0.37	4.07	0.79	60.86 ^{ab}	27.91 ^{abcd}
	69 ECS	92.05 ^{bcdef}	0.68	6.70	0.56	58.12 ^{abc}	29.33 ^{ab}
	Average	93.40 ^A	0.53	5.39 ^B	0.68	59.49 ^A	28.62 ^A
All Strains	69 EC	94.94 ^Y	0.48	3.92 ^Y	0.65 ^Y	59.88 ^Y	27.53 ^Y
	69 ECS	90.75 ^Z	0.48	7.71 ^Z	1.06 ^Z	57.79 ^Z	28.26 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 60. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder (Strain)	Production System	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Produc- tion	Egg Mass	Mortality
		(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA Amber	69 EC	107.32	0.41	485.17	73.83	44.12	19.85 ^{ab}
	69 ECS	112.52	0.37	482.84	69.35	41.38	17.37 ^{ab}
	Average	109.92 ^{AB}	0.39 ^B	484.00	71.59	42.75 ^C	18.61 ^{AB}
TETRA Brown	69 EC	107.60	0.40	471.90	69.79	43.08	36.11 ^a
	69 ECS	111.46	0.39	484.05	70.74	43.55	21.30 ^{ab}
	Average	109.53 ^{AB}	0.39 ^{AB}	477.97	70.27	43.31 ^{BC}	28.71 ^A
Novogen Brown	69 EC	104.78	0.44	490.33	74.08	46.36	15.00 ^{ab}
	69 ECS	111.42	0.42	496.85	73.85	46.67	20.56 ^{ab}
	Average	108.10 ^{AB}	0.43 ^A	493.59	73.97	46.52 ^{AB}	17.78 ^{AB}
Lohmann LB-Lite	69 EC	105.60	0.45	492.53	77.58	48.31	17.90 ^{ab}
	69 ECS	110.52	0.42	503.69	74.20	46.32	26.40 ^a
	Average	108.06 ^{AB}	0.43 ^A	498.11	75.89	47.32 ^{AB}	22.15 ^{AB}
Hy-Line Silver Brown	69 EC	110.08	0.41	488.96	75.93	45.47	12.97 ^{ab}
	69 ECS	115.38	0.39	491.64	75.81	45.14	21.53 ^{ab}
	Average	112.73 ^A	0.40 ^{AB}	490.30	75.87	45.30 ^{ABC}	17.25 ^{AB}
Hy-Line Brown	69 EC	105.17	0.42	481.45	71.99	44.65	6.48 ^b
	69 ECS	111.40	0.40	487.25	71.90	44.89	17.79 ^{ab}
	Average	108.29 ^{AB}	0.41 ^{AB}	484.35	71.95	44.77 ^{ABC}	12.13 ^B
ISA Brown	69 EC	106.38	0.44	483.80	74.52	46.66	20.38 ^{ab}
	69 ECS	108.12	0.43	492.30	76.21	47.17	14.94 ^{ab}
	Average	107.25 ^B	0.43 ^A	488.05	75.36	46.92 ^{AB}	17.66 ^{AB}
Bovans Brown	69 EC	110.71	0.43	502.47	75.97	47.49	15.08 ^{ab}
	69 ECS	114.59	0.41	500.96	76.57	48.06	15.56 ^{ab}
	Average	112.65 ^A	0.42 ^{AB}	501.71	76.27	47.78 ^A	15.32 ^{AB}
All Strains	69 EC	107.20 ^Y	0.42 ^Z	487.45	74.21	45.77	17.97
	69 ECS	111.93 ^Z	0.40 ^Y	487.08	73.58	45.40	19.43

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain* production system interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 61. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)		(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 EC	59.53	0.02	4.01	12.51	26.52	55.66
	69 ECS	59.70	0.27	4.46	12.77	23.33	57.09
	Average	59.61 ^C	0.14	4.23	12.64 ^A	24.92 ^{AB}	56.37 ^B
TETRA Brown	69 EC	61.62	0	2.70	9.14	22.58	64.83
	69 ECS	61.47	0	2.48	8.93	22.40	64.72
	Average	61.54 ^{AB}	0	2.59	9.04 ^{AB}	22.49 ^{ABC}	64.78 ^{AB}
Novogen Brown	69 EC	62.46	0	2.07	6.89	20.11	69.96
	69 ECS	63.17	0	1.83	7.72	17.59	70.44
	Average	62.81 ^A	0	1.95	7.30 ^B	18.85 ^{BC}	70.20 ^A
Lohmann LB-Lite	69 EC	61.92	0	3.27	7.42	19.07	69.34
	69 ECS	62.20	0	2.80	7.53	16.71	70.25
	Average	62.06 ^A	0	3.04	7.48 ^{AB}	17.87 ^C	69.80 ^A
Hy-Line Silver Brown	69 EC	59.80	0.06	2.57	11.76	27.55	56.82
	69 ECS	59.68	0	2.61	14.05	25.39	56.10
	Average	59.74 ^{BC}	0.03	2.59	12.91 ^A	26.47 ^A	56.46 ^B
Hy-Line Brown	69 EC	62.04	0.08	1.35	7.84	22.88	66.49
	69 ECS	62.56	0	0.91	7.09	21.08	68.62
	Average	62.30 ^A	0.04	1.13	7.46 ^{AB}	21.98 ^{ABC}	67.56 ^A
ISA Brown	69 EC	62.33	0	2.49	6.59	21.86	67.64
	69 ECS	61.68	0.28	1.57	8.20	21.76	66.21
	Average	62.01 ^A	0.14	2.03	7.39 ^B	21.81 ^{ABC}	66.93 ^A
Bovans Brown	69 EC	62.26	0.02	2.09	8.96	19.84	67.39
	69 ECS	62.84	0	2.07	7.84	19.52	68.23
	Average	62.55 ^A	0.01	2.08	8.40 ^{AB}	19.68 ^{BC}	67.81 ^A
All Strains	69 EC	61.49	0.02	2.57	8.89	22.55	64.77
	69 ECS	61.66	0.07	2.34	9.27	20.97	65.21

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 62. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEMS

Breeder	Production System	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)		(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 EC	95.88 ^{ab}	0.38	3.08	0.64	54.12 ^{ab}	26.37
	69 ECS	91.84 ^{de}	0.35	6.92	0.88	49.60 ^b	27.55
	Average	93.86	0.37	5.00	0.76	51.86 ^B	26.96 ^{ABC}
TETRA Brown	69 EC	95.00 ^{abcd}	0.26	4.16	0.58	51.99 ^{ab}	26.37
	69 ECS	91.96 ^{de}	0.59	6.71	0.71	51.79 ^{ab}	27.30
	Average	93.48	0.43	5.43	0.64	51.89 ^B	26.83 ^{ABC}
Novogen Brown	69 EC	96.37 ^{ab}	0.36	2.73	0.51	55.59 ^{ab}	25.78
	69 ECS	91.17 ^e	0.79	7.22	0.82	54.05 ^{ab}	27.32
	Average	93.77	0.58	4.98	0.66	54.82 ^{AB}	26.55 ^{BC}
Lohmann LB-Lite	69 EC	96.59 ^a	0.47	2.41	0.58	58.16 ^a	26.00
	69 ECS	90.89 ^e	0.25	7.68	1.16	54.24 ^{ab}	27.06
	Average	93.74	0.35	5.05	0.87	56.20 ^A	26.53 ^{BC}
Hy-Line Silver Brown	69 EC	96.24 ^{ab}	0.32	2.89	0.53	56.03 ^a	27.05
	69 ECS	93.17 ^{bcde}	0.62	5.70	0.53	55.09 ^{ab}	28.22
	Average	94.71	0.47	4.29	0.53	55.56 ^{AB}	27.63 ^{AB}
Hy-Line Brown	69 EC	94.76 ^{abcd}	0.51	3.73	0.98	53.27 ^{ab}	25.88
	69 ECS	92.36 ^{de}	0.37	6.80	0.46	52.99 ^{ab}	27.32
	Average	93.56	0.44	5.27	0.72	53.13 ^{AB}	26.60 ^{BC}
ISA Brown	69 EC	95.48 ^{abc}	0.84	3.01	0.64	55.90 ^a	26.07
	69 ECS	92.78 ^{cde}	0.47	6.22	0.52	56.24 ^a	26.52
	Average	94.13	0.66	4.61	0.58	56.07 ^A	26.30 ^C
Bovans Brown	69 EC	94.70 ^{abcd}	0.29	4.30	0.71	56.27 ^a	27.18
	69 ECS	92.28 ^{de}	0.12	6.68	0.96	56.19 ^a	28.11
	Average	93.49	0.21	5.49	0.84	56.23 ^A	27.65 ^A
All Strains	69 EC	95.63 ^Y	0.43	3.29 ^Y	0.65	55.17	26.34 ^Y
	69 ECS	92.06 ^Z	0.45	6.74 ^Z	0.76	53.77	27.42 ^Z

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

abcde - Different letters denote significant differences (P<.01) in the strain* production system interactions.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

YZ - Different letters denote significant differences (P<.01), comparisons made among production system average values.

TABLE 63. EFFECT OF WHITE EGG STRAIN AND PRODUCTION SYSTEM ON BODY WEIGHT OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEM: NON-MOLTED PROGRAM

Breeder	Production System	17 Wk* Body Wt	69 Wk** Body Wt	1st Cycle Wt Gain	73 Wk** Body Wt	109 Wk Body Wt	2 nd Cycle Wt Gain	Total Wt Gain
(Strain)		(kg)	(kg)	(%)	(kg)	(kg)	(kg)	(%)
Bovans White	69 EC	1.22	1.74	42.6	1.78	1.83	0.04	49.16
	69 ECS	1.22	1.64	34.4	1.72	1.80	0.10	48.82
	Average	1.22 ^{AB}	1.69	38.5	1.75	1.82 ^{ABC}	0.07	48.99 ^{AB}
Shaver White	69 EC	1.35	1.66	23.0	1.71	1.84	0.12	41.13
	69 ECS	1.28	1.73	34.4	1.74	1.80	0.08	44.46
	Average	1.32 ^A	1.70	28.8	1.72	1.82 ^{AB}	0.10	43.29 ^{AB}
Dekalb White	69 EC	1.20	1.63	36.7	1.73	1.81	0.10	50.27
	69 ECS	1.20	1.61	35.0	1.72	1.75	0	48.06
	Average	1.20 ^{AB}	1.62	35.8	1.72	1.78 ^{BC}	0.06	49.16 ^{AB}
Babcock White	69 EC	1.28	1.78	39.1	1.79	1.89	0.10	45.71
	69 ECS	1.31	1.74	32.1	1.73	1.95	0.16	50.72
	Average	1.30 ^{AB}	1.76	35.4	1.76	1.92 ^A	0.14	48.21 ^{AB}
ISA B-400	69 EC	1.22	1.67	36.9	1.68	1.75	0.10	44.77
	69 ECS	1.18	1.63	39.0	2.09	1.61	-0.15	34.66
	Average	1.20 ^{AB}	1.65	37.5	1.88	1.68 ^C	-0.02	39.71 ^B
Hy-Line W-36	69 EC	1.22	1.65	35.2	1.72	1.86	0.16	54.31
	69 ECS	1.20	1.60	33.3	1.57	1.81	0.18	52.41
	Average	1.21 ^{AB}	1.62	33.9	1.64	1.83 ^{AB}	0.17	53.36 ^A
Hy-Line CV-24	69 EC	1.22	1.62	32.8	1.62	1.79	0.18	44.31
	69 ECS	1.22	1.65	36.1	1.69	1.84	0.15	47.75
	Average	1.22 ^{AB}	1.64	34.4	1.66	1.81 ^{ABC}	0.17	46.03 ^{AB}
Lohmann LSL Lite	69 EC	1.27	1.68	32.3	1.63	1.86	0.20	49.26
	69 ECS	1.22	1.58	29.5	1.97	1.84	-0.07	48.58
	Average	1.24 ^{AB}	1.63	30.6	1.80	1.85 ^{ABC}	0.06	48.92 ^{AB}
H&N Nick Chick	69 EC	1.21	1.66	36.4	1.71	1.84	0.08	43.82
	69 ECS	1.24	1.66	33.9	1.67	1.76	0.07	41.57
	Average	1.23 ^{AB}	1.66	35.0	1.69	1.80 ^{ABC}	0.07	42.70 ^{AB}
Novogen White	69 EC	1.24	1.71	37.9	1.75	1.85	0.09	45.52
	69 ECS	1.22	1.64	34.4	1.64	1.81	0.14	46.90
	Average	1.23 ^{AB}	1.68	35.8	1.70	1.83 ^{AB}	0.12	46.21 ^{AB}
All Strains	69 EC	1.24	1.68	35.5	1.71	1.83	0.12	46.96
	69 ECS	1.22	1.66	36.1	1.76	1.81	0.06	48.47

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

(*) All replicates in all strains were weight at 17 wks.

(**) Only a sample of replicates (2 per strain treatment) in each strain were weighted at 69 and 73 wks.

TABLE 64. EFFECT OF BROWN EGG STRAIN AND PRODUCTION SYSTEM ON BODY WEIGHT OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN ENRICHABLE AND ENRICHED COLONY HOUSING SYSTEM: NON-MOLTED PROGRAM

Breeder	Production System	17 Wk* Body Wt	69 Wk** Body Wt	1st Cycle Wt Gain	73 Wk** Body Wt	109 Wk Body Wt	2 nd Cycle Wt Gain	Total Wt Gain
(Strain)		(kg)	(kg)	(%)	(kg)	(kg)	(kg)	(%)
TETRA Amber	69 EC	1.50	2.10	40.0	2.07	1.89	0.02	22.48
	69 ECS	1.50	1.89	26.7	2.02	1.84	0.04	20.75
	Average	1.50	1.99	33.3	2.04	1.86	0.03	21.61
TETRA Brown	69 EC	1.64	1.94	18.3	1.81	1.90	0.25	16.24
	69 ECS	1.54	1.88	22.1	1.87	1.92	0.18	26.18
	Average	1.58	1.91	20.3	1.84	1.91	0.22	21.71
Novogen Brown	69 EC	1.60	1.79	11.9	1.95	1.94	-0.02	24.52
	69 ECS	1.55	1.94	25.2	1.91	1.82	0.07	16.02
	Average	1.58	1.86	18.4	1.93	1.88	0.02	20.27
Lohmann LB-Lite	69 EC	1.44	1.79	25.0	1.82	1.86	0.10	23.22
	69 ECS	1.52	2.00	31.6	1.91	1.93	0.07	26.16
	Average	1.48	1.89	28.4	1.86	1.90	0.08	24.69
Hy-Line Silver Brown	69 EC	1.64	1.86	13.4	2.02	2.10	0.14	20.72
	69 ECS	1.42	1.98	39.4	1.98	1.80	0.16	21.66
	Average	1.53	1.92	25.5	2.00	1.95	0.15	26.19
Hy-Line Brown	69 EC	1.62	2.00	23.5	2.00	1.91	0.13	20.16
	69 ECS	1.53	1.87	22.2	2.38	1.96	-0.26	29.75
	Average	1.58	1.94	22.8	2.19	1.94	-0.07	24.96
ISA Brown	69 EC	1.47	1.92	31.3	1.92	2.04	0.06	38.97
	69 ECS	1.45	1.82	25.5	1.87	1.87	0.08	30.73
	Average	1.46	1.87	28.8	1.89	1.95	0.07	34.85
Bovans Brown	69 EC	1.52	1.98	30.3	1.99	1.93	0.03	28.34
	69 ECS	1.60	1.94	21.3	1.94	1.89	0.04	22.78
	Average	1.56	1.96	25.6	1.96	1.91	0.04	25.56
All Strains	69 EC	1.55	1.92	23.9	1.95	1.94	0.09	25.71
	69 ECS	1.51	1.92	26.5	1.98	1.88	0.04	24.25

Enrichable Cage=EC; Enriched Colony Housing System=ECS.

(*) All replicates in all strains were weight at 17 wks.

(**) Only a sample of replicates (2 per strain treatment) in each strain were weighted at 69 and 73 wks.

TABLE 65. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
Bovans White	69 ECS	10.51 ^{abc}	0.492 ^{cde}	311.22 ^{abc}	84.47 ^{efg}	50.87 ^{fg}	17.06	146.14 ^a
	138 ECS	10.80 ^{ab}	0.501 ^{abcde}	317.25 ^{abc}	86.78 ^{abcde}	53.13 ^{bcde}	9.52	143.14 ^{abcd}
	Average	10.66 ^{AB}	0.496 ^C	314.24 ^{ABC}	85.76 ^C	52.00 ^B	13.30	144.64 ^A
Shaver White	69 ECS	10.50 ^{abc}	0.540 ^{ab}	318.28 ^{abc}	85.56 ^{bcdef}	51.91 ^{cdef}	30.55	140.50 ^{cd}
	138 ECS	10.53 ^{abc}	0.527 ^{abcd}	319.98 ^{abc}	87.67 ^{abcd}	53.52 ^{abcde}	2.78	142.00 ^{abcd}
	Average	10.52 ^B	0.534 ^{AB}	318.83 ^{AB}	86.61 ^{BC}	52.72 ^B	16.17	141.25 ^B
Dekalb White	69 ECS	10.48 ^{bc}	0.508 ^{abcde}	311.31 ^{abc}	85.18 ^{cdefg}	51.54 ^{ef}	9.03	144.25 ^{abcd}
	138 ECS	10.66 ^{ab}	0.500 ^{abcde}	322.68 ^{ab}	88.51 ^{ab}	53.33 ^{bcde}	4.17	142.25 ^{abcd}
	Average	10.57 ^{AB}	0.504 ^{ABC}	316.99 ^{ABC}	86.84 ^{BC}	52.44 ^B	6.60	143.25 ^{AB}
Babcock White	69 ECS	11.09 ^a	0.539 ^{abc}	327.78 ^a	89.54 ^a	54.27 ^{ab}	11.80	140.50 ^{cd}
	138 ECS	10.72 ^{ab}	0.538 ^{abc}	325.23 ^{ab}	89.33 ^a	55.64 ^a	3.33	141.20 ^{bcd}
	Average	10.90 ^A	0.538 ^A	326.30 ^A	89.44 ^A	54.96 ^A	7.57	140.85 ^B
ISA B-400	69 ECS	10.68 ^{ab}	0.551 ^a	325.21 ^{ab}	88.61 ^{ab}	54.74 ^{ab}	17.50	141.00 ^{bcd}
	138 ECS	10.95 ^{ab}	0.518 ^{abcde}	321.20 ^{abc}	87.96 ^{abc}	54.84 ^{ab}	6.95	139.75 ^d
	Average	10.81 ^{AB}	0.534 ^{AB}	323.20 ^A	88.28 ^{AB}	54.79 ^A	12.22	140.38 ^B
Hy-Line W-36	69 ECS	9.89 ^{cd}	0.493 ^{abcde}	298.27 ^{bc}	81.81 ^{gh}	48.22 ^h	4.63	145.33 ^{abc}
	138 ECS	9.79 ^d	0.503 ^{abcde}	299.15 ^{abc}	82.07 ^{fgh}	49.02 ^{gh}	1.85	145.67 ^{abc}
	Average	9.84 ^C	0.498 ^{BC}	298.71 ^C	81.94 ^D	48.62 ^D	3.24	145.50 ^A
Hy-Line CV-24	69 ECS	10.80 ^{ab}	0.464 ^e	295.21 ^c	80.70 ^h	48.93 ^{gh}	11.11	145.75 ^{ab}
	138 ECS	10.67 ^{ab}	0.483 ^{de}	310.53 ^{abc}	85.17 ^{cdefg}	51.67 ^{def}	6.95	145.25 ^{abc}
	Average	10.74 ^{AB}	0.474 ^C	302.87 ^{BC}	82.93 ^D	50.30 ^C	9.03	145.50 ^A
Lohmann LSL Lite	69 ECS	10.68 ^{ab}	0.498 ^{abcde}	310.94 ^{abc}	84.85 ^{cdefg}	51.60 ^{def}	13.19	144.75 ^{abc}
	138 ECS	10.77 ^{ab}	0.503 ^{abcde}	319.88 ^{abc}	87.34 ^{abcde}	54.10 ^{abc}	11.11	144.25 ^{abcd}
	Average	10.73 ^{AB}	0.500 ^{BC}	315.14 ^{ABC}	86.10 ^C	52.85 ^B	12.15	144.50 ^A
H&N Nick Chick	69 ECS	10.66 ^{ab}	0.497 ^{abcde}	309.98 ^{abc}	84.44 ^{defg}	51.63 ^{def}	15.97	145.75 ^{ab}
	138 ECS	10.85 ^{ab}	0.495 ^{abcde}	315.70 ^{abc}	86.60 ^{abcde}	53.82 ^{abcd}	6.94	144.25 ^{abcd}
	Average	10.76 ^{AB}	0.496 ^C	312.84 ^{ABC}	85.52 ^C	52.73 ^B	11.46	145.00 ^A
Novogen White	69 ECS	10.95 ^{ab}	0.480 ^{de}	304.77 ^{abc}	82.70 ^{fgh}	50.90 ^{fg}	20.55	145.00 ^{abc}
	138 ECS	10.77 ^{ab}	0.513 ^{abcde}	318.44 ^{abc}	87.24 ^{abcde}	54.47 ^{ab}	7.78	143.67 ^{abcd}
	Average	10.86 ^{AB}	0.496 ^C	316.61 ^{ABC}	84.97 ^C	52.69 ^B	14.17	144.33 ^A
All Strains	69 ECS	10.62	0.506	311.26 ^Z	84.81 ^Z	51.46 ^Z	15.14 ^Z	143.90
	138 ECS	10.65	0.508	316.94 ^Y	86.86 ^Y	53.36 ^Y	6.14 ^Y	143.14

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefgh - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 66. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 ECS	58.90 ^{ghi}	0.31	8.10	12.04 ^{abc}	25.21	54.58 ^{de}
	138 ECS	60.02 ^{abcdefg}	3.44	3.53	7.97 ^c	25.48	59.46 ^{bcd}
	Average	59.46 ^{DE}	1.87	5.81	10.00 ^B	25.35 ^{AB}	57.02 ^C
Shaver White	69 ECS	59.77 ^{cdefgh}	0	4.75	11.85 ^{abc}	24.27	58.67 ^{cd}
	138 ECS	60.05 ^{abcdefg}	0.96	4.31	9.36 ^{abc}	23.31	61.90 ^{abcd}
	Average	59.91 ^{BCD}	0.48	4.53	10.61 ^B	23.79 ^{AB}	60.29 ^{ABC}
Dekalb White	69 ECS	59.22 ^{efghi}	0	6.88	10.50 ^{abc}	23.42	58.83 ^{cd}
	138 ECS	58.48 ^{hi}	2.23	3.02	13.67 ^{abc}	24.33	56.54 ^{cde}
	Average	58.85 ^E	1.11	4.95	12.08 ^{AB}	23.87 ^{AB}	57.68 ^{BC}
Babcock White	69 ECS	59.65 ^{defghi}	0.52	5.08	11.96 ^{abc}	23.36	58.54 ^{cd}
	138 ECS	61.31 ^a	0.37	4.55	7.01 ^c	20.01	67.86 ^a
	Average	60.48 ^{AB}	0.44	4.82	9.49 ^B	21.69 ^{AB}	63.20 ^{AB}
ISA B-400	69 ECS	60.81 ^{abcd}	0	4.34	9.26 ^{abc}	22.29	63.62 ^{abc}
	138 ECS	61.33 ^{ab}	1.44	3.02	8.03 ^{bc}	19.43	67.63 ^{ab}
	Average	61.07 ^A	0.72	3.68	8.65 ^B	20.86 ^{AB}	65.63 ^A
Hy-Line W-36	69 ECS	58.02 ⁱ	0	8.31	17.10 ^a	25.54	48.51 ^e
	138 ECS	59.01 ^{fghi}	2.99	1.41	16.13 ^{ab}	25.51	54.00 ^{de}
	Average	58.52 ^E	1.49	5.86	16.61 ^A	25.52 ^{AB}	51.26 ^D
Hy-Line CV-24	69 ECS	59.86 ^{abcdefgh}	0.27	8.20	11.04 ^{abc}	22.62	57.67 ^{cde}
	138 ECS	59.09 ^{fghi}	1.74	3.58	13.70 ^{abc}	22.15	59.00 ^{bcd}
	Average	59.47 ^{CDE}	1.00	5.89	12.37 ^{AB}	22.38 ^{AB}	58.34 ^{BC}
Lohmann LSL Lite	69 ECS	59.66 ^{defghi}	0	5.06	10.83 ^{abc}	23.98	59.71 ^{abcd}
	138 ECS	60.58 ^{ef}	2.25	3.48	9.51 ^{abc}	19.48	65.08 ^{abc}
	Average	60.12 ^{ABCD}	1.12	4.26	10.17 ^B	21.73 ^{AB}	62.39 ^{AB}
H&N Nick Chick	69 ECS	60.14 ^{abcdefg}	0	5.35	11.00 ^{abc}	22.38	60.83 ^{abcd}
	138 ECS	60.71 ^{abcde}	3.85	2.33	10.56 ^{abc}	18.85	64.54 ^{abc}
	Average	60.42 ^{ABC}	1.92	3.84	10.78 ^B	20.62 ^{AB}	62.68 ^{AB}
Novogen White	69 ECS	60.51 ^{abcdef}	0.20	5.34	10.82 ^{abc}	22.13	61.31 ^{abcd}
	138 ECS	61.20 ^{abc}	1.53	5.91	6.16 ^c	18.39	67.38 ^{ab}
	Average	60.86 ^{AB}	0.86	5.62	8.49 ^B	20.26 ^B	64.34 ^A
All Strains	69 ECS	59.65 ^Z	0.13 ^Z	6.14 ^Y	11.54	23.52 ^Y	58.23 ^Z
	138 ECS	60.18 ^Y	2.08 ^Y	3.51 ^Z	10.21	21.69 ^Z	62.34 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefghi - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 67. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 ECS	94.83	0.50	4.38	0.31	35.17 ^{abc}	15.71 ^{abc}
	138 ECS	91.60	0.44	7.66	0.31	35.67 ^{abc}	16.17 ^{abc}
	Average	93.21	0.47	6.02	0.31	35.42 ^{AB}	15.94 ^A
Shaver White	69 ECS	93.62	0.42	5.00	0.95	35.79 ^{abc}	15.70 ^{abc}
	138 ECS	89.80	0.68	9.05	0.48	35.45 ^{abc}	15.77 ^{abc}
	Average	91.71	0.55	7.02	0.71	35.62 ^{AB}	15.74 ^{AB}
Dekalb White	69 ECS	92.75	0.55	6.10	0.58	35.19 ^{abc}	15.68 ^{abc}
	138 ECS	93.98	0	5.22	0.80	36.23 ^{abc}	15.96 ^{abc}
	Average	93.36	0.28	5.66	0.69	35.17 ^{AB}	15.82 ^{AB}
Babcock White	69 ECS	93.50	0.80	5.05	0.68	36.79 ^{ab}	16.56 ^a
	138 ECS	92.38	0.40	6.76	0.48	36.77 ^{ab}	16.06 ^{abc}
	Average	92.94	0.60	5.90	0.58	36.78 ^A	16.31 ^A
ISA B-400	69 ECS	93.82	0.45	5.20	0.55	36.91 ^a	15.97 ^{abc}
	138 ECS	90.00	0.28	8.90	0.82	35.89 ^{abc}	16.39 ^{abc}
	Average	91.91	0.36	7.05	0.69	36.40 ^A	16.18 ^A
Hy-Line W-36	69 ECS	93.57	0.37	5.23	0.77	32.87 ^c	14.81 ^{bc}
	138 ECS	93.13	0	6.17	0.73	33.30 ^{bc}	14.66 ^c
	Average	93.35	0.18	5.70	0.75	33.08 ^C	14.73 ^B
Hy-Line CV-24	69 ECS	93.60	0.38	5.50	0.58	33.04 ^c	16.14 ^{abc}
	138 ECS	93.38	0.15	6.35	0.12	34.87 ^{abc}	15.97 ^{abc}
	Average	93.49	0.26	5.92	0.35	33.96 ^{BC}	16.07 ^A
Lohmann LSL Lite	69 ECS	92.90	1.15	5.45	0.52	35.21 ^{abc}	15.96 ^{abc}
	138 ECS	93.18	0.20	6.35	0.25	36.24 ^{abc}	16.11 ^{abc}
	Average	93.04	0.68	5.90	0.39	35.72 ^{AB}	16.04 ^A
H&N Nick Chick	69 ECS	93.90	0.65	4.58	0.88	35.07 ^{abc}	15.94 ^{abc}
	138 ECS	93.58	0.50	5.80	0.12	35.77 ^{abc}	16.25 ^{abc}
	Average	93.74	0.58	5.19	0.50	35.42 ^{AB}	16.09 ^A
Novogen White	69 ECS	93.74	0.52	5.02	0.52	34.59 ^{abc}	16.08 ^{abc}
	138 ECS	90.56	0.66	8.00	0.78	35.58 ^{abc}	16.39 ^{ab}
	Average	92.25	0.59	6.51	0.65	35.09 ^{ABC}	16.24 ^A
All Strains	69 ECS	93.64	0.58	5.15 ^Y	0.63	35.06	15.88
	138 ECS	92.16	0.33	7.02 ^Z	0.49	35.58	15.94

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 68. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality	Age at 50% Production
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)	(Days)
TETRA	69 ECS	10.60 ^{cdef}	0.442 ^{bc}	283.01	77.69 ^{bc}	44.55 ^d	1.39	145.50 ^{ab}
Amber	138 ECS	11.04 ^{abcd}	0.432 ^c	290.02	79.44 ^{abc}	46.57 ^{bcd}	6.94	143.00 ^{ab}
	Average	10.82 ^B	0.437 ^D	286.52	78.57 ^{BC}	45.56 ^C	4.17	144.25 ^{ABC}
TETRA	69 ECS	10.38 ^{ef}	0.464 ^{abc}	282.18	77.28 ^{bc}	45.84 ^{cd}	5.56	145.00 ^{ab}
Brown	138 ECS	10.98 ^{abcde}	0.438 ^{bc}	278.31	76.31 ^{bc}	45.49 ^{cd}	1.85	140.33 ^{ab}
	Average	10.68 ^{BC}	0.450 ^{CD}	280.24	76.79 ^C	45.66 ^C	3.70	142.67 ^{ABC}
Novogen	69 ECS	10.49 ^{def}	0.510 ^a	297.85	81.75 ^{abc}	49.82 ^{abc}	3.34	144.00 ^{ab}
Brown	138 ECS	11.18 ^{ab}	0.439 ^{bc}	276.89	75.91 ^c	46.91 ^{bcd}	8.89	141.80 ^{ab}
	Average	10.84 ^B	0.475 ^{ABCD}	287.37	78.83 ^{BC}	48.36 ^{ABC}	6.11	142.90 ^{ABC}
Lohmann	69 ECS	10.36 ^f	0.485 ^{abc}	294.17	80.74 ^{abc}	48.49 ^{abcd}	3.48	145.75 ^{ab}
LB-Lite	138 ECS	10.44 ^{ef}	0.515 ^a	310.48	85.34 ^a	51.87 ^a	0	142.25 ^{ab}
	Average	10.40 ^C	0.500 ^A	302.32	83.04 ^{AB}	50.18 ^{AB}	1.74	144.00 ^{ABC}
Hy-Line	69 ECS	10.79 ^{bcd}	0.487 ^{abc}	303.42	83.20 ^{ab}	47.73 ^{abcd}	6.94	141.50 ^{ab}
Silver Brown	138 ECS	10.981 ^{abcde}	0.463 ^{abc}	304.12	83.43 ^{ab}	48.37 ^{abcd}	4.17	140.00 ^{ab}
	Average	10.85 ^{AB}	0.475 ^{ABCD}	303.77	83.32 ^A	48.05 ^{ABC}	5.56	140.75 ^{BC}
Hy-Line	69 ECS	10.65 ^{cdef}	0.494 ^{ab}	289.23	78.35 ^{abc}	47.64 ^{abcd}	4.44	142.20 ^{ab}
Brown	138 ECS	11.11 ^{abc}	0.458 ^{abc}	284.33	78.11 ^{bc}	47.81 ^{abcd}	0	139.40 ^b
	Average	10.88 ^{AB}	0.476 ^{BC}	286.78	78.73 ^{BC}	47.73 ^{BC}	2.22	140.80 ^C
ISA	69 ECS	10.37 ^f	0.504 ^a	308.91	84.73 ^a	50.87 ^{ab}	4.45	146.00 ^a
Brown	138 ECS	10.76 ^{bcd}	0.480 ^{abc}	296.40	81.29 ^{abc}	50.71 ^{ab}	2.22	144.60 ^{ab}
	Average	10.56 ^{BC}	0.492 ^{AB}	302.65	83.01 ^A	50.79 ^A	3.34	145.30 ^A
Bovans	69 ECS	10.96 ^{abcde}	0.472 ^{abc}	299.50	81.12 ^{abc}	49.93 ^{abc}	5.56	146.00 ^a
Brown	138 ECS	11.40 ^a	0.444 ^{bc}	286.05	78.51 ^{abc}	49.28 ^{abc}	4.17	144.00 ^{ab}
	Average	10.18 ^A	0.458 ^{BCD}	292.78	80.31 ^{ABC}	49.60 ^{AB}	4.86	145.00 ^{AB}
All	69 ECS	10.58 ^Z	0.482 ^Z	294.78	80.86	48.11	4.39	144.49 ^Y
Strains	138 ECS	10.98 ^Y	0.459 ^Y	290.82	79.79	48.38	3.53	141.92 ^Z

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABCd - Different letters denote significant differences (P<.01), comparisons made among strain average values

abcde - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 69. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 ECS	56.51 ^h	0.48	7.89 ^a	22.24 ^{ab}	28.06 ^{abc}	41.14 ^{ef}
	138 ECS	57.60 ^{fgh}	1.92	4.88 ^{ab}	16.00 ^{bcd}	29.67 ^{ab}	47.38 ^{def}
	Average	57.05 ^D	1.20	6.38 ^A	19.12 ^A	28.86 ^{ABC}	44.26 ^C
TETRA Brown	69 ECS	58.56 ^{efg}	0	4.37 ^{ab}	15.81 ^{bcd}	28.77 ^{abc}	51.00 ^{cde}
	138 ECS	59.09 ^{def}	3.41	3.54 ^{ab}	9.17 ^{cde}	29.02 ^{abc}	54.69 ^{cd}
	Average	58.83 ^C	1.71	3.96 ^{AB}	12.49 ^B	28.90 ^{ABC}	52.85 ^B
Novogen Brown	69 ECS	60.19 ^{bcd}	0	3.24 ^{ab}	13.49 ^{cde}	25.86 ^{abc}	57.09 ^{bcd}
	138 ECS	60.97 ^{abc}	0.68	2.66 ^b	7.05 ^e	20.46 ^{bc}	69.00 ^a
	Average	60.58 ^{AB}	0.34	2.95 ^B	10.27 ^B	23.16 ^C	63.01 ^A
Lohmann LB-Lite	69 ECS	59.04 ^{def}	0	4.68 ^{ab}	13.23 ^{cde}	24.27 ^{abc}	56.92 ^{bcd}
	138 ECS	60.01 ^{bcd}	0.56	4.13 ^{ab}	8.98 ^{de}	25.41 ^{abc}	60.75 ^{abc}
	Average	59.53 ^{BC}	0.28	4.40 ^{AB}	11.10 ^B	24.85 ^{BC}	58.84 ^{AB}
Hy-Line Silver Brown	69 ECS	56.18 ^{gh}	0.13	4.44 ^{ab}	25.28 ^a	31.65 ^a	38.71 ^f
	138 ECS	57.28 ^{fgh}	1.08	4.96 ^{ab}	17.99 ^{abc}	33.81 ^a	41.54 ^{ef}
	Average	57.05 ^D	0.60	4.70 ^{AB}	21.63 ^A	32.73 ^A	40.12 ^C
Hy-Line Brown	69 ECS	59.50 ^{cde}	0	1.61 ^b	12.23 ^{cde}	31.83 ^a	53.86 ^{cd}
	138 ECS	60.51 ^{abcd}	1.09	3.34 ^{ab}	7.42 ^e	27.64 ^{abc}	60.37 ^{abc}
	Average	60.00 ^{BC}	0.55	2.48 ^B	9.82 ^B	29.74 ^{AB}	57.11 ^{AB}
ISA Brown	69 ECS	58.96 ^{def}	0.51	2.81 ^{ab}	14.40 ^{bcd}	29.69 ^{abc}	53.25 ^{cd}
	138 ECS	61.67 ^{ab}	0	4.64 ^{ab}	7.17 ^e	18.91 ^c	69.09 ^a
	Average	60.31 ^{AB}	0.25	3.72 ^{AB}	10.78 ^B	23.80 ^C	61.17 ^A
Bovans Brown	69 ECS	60.44 ^{abcd}	0	3.60 ^{ab}	13.83 ^{cde}	26.51 ^{abc}	55.78 ^{cd}
	138 ECS	61.90 ^a	0.13	4.06 ^{ab}	8.14 ^{de}	20.42 ^{bc}	67.40 ^{ab}
	Average	61.17 ^A	0.07	3.83 ^{AB}	10.99 ^B	23.46 ^C	61.59 ^A
All Strains	69 ECS	58.75 ^Z	0.14 ^Z	4.08	16.31 ^Y	28.20 ^Y	50.97 ^Z
	138 ECS	59.88 ^Y	1.11 ^Y	4.02	10.24 ^Z	25.67 ^Z	58.78 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABCD - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde fgh - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 70. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-483 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 ECS	92.58	0.42	6.62	0.38	30.55	15.68 ^{ab}
	138 ECS	91.18	1.40	7.20	0.25	31.87	16.37 ^{ab}
	Average	91.88	0.91	6.91	0.31	31.21	16.02 ^{AB}
TETRA Brown	69 ECS	92.87	0.60	5.87	0.70	31.43	15.38 ^{ab}
	138 ECS	93.67	0.17	6.03	0.17	31.26	16.29 ^{ab}
	Average	93.27	0.38	5.95	0.43	31.34	15.83 ^{AB}
Novogen Brown	69 ECS	92.70	0.62	5.94	0.72	33.39	15.55 ^{ab}
	138 ECS	92.62	0.54	6.38	0.50	31.44	16.59 ^{ab}
	Average	92.66	0.58	6.16	0.61	32.42	16.07 ^{AB}
Lohmann LB-Lite	69 ECS	92.92	0.38	5.52	1.18	33.06	15.34 ^b
	138 ECS	91.10	0.52	8.12	0.25	34.83	15.47 ^{ab}
	Average	92.10	0.45	6.82	0.71	33.95	15.40 ^B
Hy-Line Silver Brown	69 ECS	95.20	0.10	4.70	0	33.46	15.97 ^{ab}
	138 ECS	94.10	0.32	4.82	0.72	33.62	16.17 ^{ab}
	Average	94.65	0.21	4.76	0.36	33.54	16.07 ^{AB}
Hy-Line Brown	69 ECS	94.06	0.22	5.20	0.56	32.74	15.76 ^{ab}
	138 ECS	91.92	0.48	7.10	0.52	31.91	16.47 ^{ab}
	Average	92.99	0.35	6.15	0.54	32.33	16.12 ^{AB}
ISA Brown	69 ECS	94.06	0.22	5.30	0.44	34.96	15.36 ^b
	138 ECS	92.42	0.10	7.26	0.20	33.69	15.95 ^{ab}
	Average	93.24	0.16	6.28	0.32	34.32	15.65 ^B
Bovans Brown	69 ECS	94.70	0.16	4.48	0.68	33.91	16.22 ^{ab}
	138 ECS	92.78	0.25	6.82	0.15	32.57	16.90 ^a
	Average	93.74	0.20	5.65	0.41	33.24	16.56 ^A
All Strains	69 ECS	93.63	0.34	5.45 ^Y	0.58	32.94	15.66 ^Z
	138 ECS	92.47	0.47	6.72 ^Z	0.34	32.65	16.28 ^Y

¹All strains were housed such that each strain is equally represented in each density.
Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 71. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans White	69 ECS	11.25 ^{abc}	0.468 ^{abcd}	23.68 ^{ab}	83.69 ^{ab}	52.82 ^{ab}	1.39
	138 ECS	11.38 ^{abc}	0.495 ^{abcd}	25.22 ^a	89.36 ^a	56.26 ^{ab}	1.39
	Average	11.32 ^{BC}	0.481	24.45 ^{AB}	86.52	54.54	1.39
Shaver White	69 ECS	11.89 ^{abc}	0.455 ^{abcd}	24.15 ^{ab}	83.16 ^{ab}	53.75 ^{ab}	3.47
	138 ECS	10.83 ^{abc}	0.545 ^a	25.45 ^a	90.89 ^a	58.97 ^a	0
	Average	11.36 ^{AB}	0.500	24.80 ^{AB}	87.02	56.36	1.74
Dekalb White	69 ECS	11.10 ^{abc}	0.495 ^{abcd}	24.41 ^{ab}	86.55 ^{ab}	54.68 ^{ab}	1.39
	138 ECS	10.96 ^{abc}	0.522 ^{abc}	25.84 ^a	91.21 ^a	57.38 ^{ab}	2.78
	Average	11.03 ^{AB}	0.509	25.12 ^A	88.84	56.03	2.08
Babcock White	69 ECS	12.82 ^a	0.415 ^{cd}	23.30 ^{ab}	82.46 ^{ab}	52.98 ^{ab}	2.08
	138 ECS	11.16 ^{abc}	0.532 ^{ab}	25.33 ^a	90.49 ^a	59.24 ^a	0
	Average	11.99 ^A	0.474	24.32 ^{AB}	86.48	56.11	1.04
ISA B-400	69 ECS	10.98 ^{abc}	0.510 ^{abc}	24.32 ^{ab}	86.44 ^{ab}	56.10 ^{ab}	0.69
	138 ECS	11.06 ^{abc}	0.512 ^{abc}	24.16 ^{ab}	85.52 ^{ab}	56.55 ^{ab}	1.39
	Average	11.02 ^{AB}	0.511	24.24 ^{AB}	85.98	56.32	1.04
Hy-Line W-36	69 ECS	10.08 ^{bc}	0.480 ^{abcd}	21.59 ^{ab}	77.10 ^{ab}	48.55 ^{ab}	0
	138 ECS	9.98 ^c	0.497 ^{abcd}	21.59 ^{ab}	77.11 ^{ab}	49.37 ^{ab}	0
	Average	10.03 ^B	0.488	21.59 ^B	77.10	48.96	0
Hy-Line CV-24	69 ECS	12.16 ^{ab}	0.442 ^{abcd}	22.97 ^{ab}	82.02 ^{ab}	53.75 ^{ab}	0.69
	138 ECS	10.89 ^{abc}	0.530 ^{abc}	25.17 ^a	89.90 ^a	57.55 ^{ab}	0
	Average	11.52 ^A	0.486	24.07 ^{AB}	85.96	55.65	0.35
Lohmann LSL Lite	69 ECS	11.86 ^{abc}	0.388 ^d	19.92 ^b	71.13 ^b	45.64 ^b	0
	138 ECS	11.28 ^{abc}	0.528 ^{abc}	25.03 ^a	89.40 ^a	59.23 ^a	0
	Average	11.57 ^A	0.458	22.47 ^{AB}	80.26	52.44	0
H&N Nick Chick	69 ECS	12.03 ^{abc}	0.458 ^{abcd}	23.96 ^{ab}	87.76 ^{ab}	55.11 ^{ab}	0.69
	138 ECS	11.29 ^{abc}	0.520 ^{abc}	25.11 ^a	88.65 ^{ab}	58.72 ^a	1.39
	Average	11.66 ^A	0.489	24.54 ^{AB}	86.81	56.91	1.04
Novogen White	69 ECS	11.77 ^{abc}	0.428 ^{bcd}	22.04 ^{ab}	76.92 ^{ab}	50.25 ^{ab}	1.67
	138 ECS	11.76 ^{abc}	0.514 ^{abc}	25.48 ^a	91.01 ^a	59.99 ^a	0
	Average	11.77 ^A	0.471	23.76 ^{AB}	83.96	55.12	0.83
All Strains	69 ECS	11.59 ^Y	0.454 ^Y	23.03 ^Z	81.42 ^Z	52.36 ^Z	1.21
	138 ECS	11.06 ^Z	0.519 ^Z	24.84 ^Y	88.36 ^Y	57.33 ^Y	0.69

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 72. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 ECS	63.08	0	0	0	31.71	58.56
	138 ECS	62.98	0	0	0	27.16	59.69
	Average	63.03 ^B	0	0	0	29.43 ^A	59.12
Shaver White	69 ECS	64.61	0	0	0.83	9.28	72.24
	138 ECS	64.90	0	0	0	1.32	76.54
	Average	64.75 ^{AB}	0	0	0.42	5.30 ^{AB}	74.39
Dekalb White	69 ECS	63.18	0	0	0	34.29	57.44
	138 ECS	62.99	0	0	0	23.10	65.41
	Average	63.08 ^{AB}	0	0	0	28.70 ^A	61.42
Babcock White	69 ECS	64.23	0	0	0	19.77	67.44
	138 ECS	65.46	0	0	0	6.35	66.32
	Average	64.84 ^{AB}	0	0	0	13.06 ^{AB}	66.88
ISA B-400	69 ECS	64.90	0	0	0	13.92	74.68
	138 ECS	66.08	0	0	0	0	71.44
	Average	65.49 ^{AB}	0	0	0	6.96 ^{AB}	73.06
Hy-Line W-36	69 ECS	62.97	0	0	0	12.39	74.28
	138 ECS	64.06	0	0	0	19.46	68.93
	Average	63.52 ^{AB}	0	0	0	15.92 ^{AB}	71.60
Hy-Line CV-24	69 ECS	65.61	0	0	0	13.50	76.18
	138 ECS	64.07	0	0	1.92	16.42	62.48
	Average	64.84 ^{AB}	0	0	0.96	14.96 ^{AB}	69.33
Lohmann LSL Lite	69 ECS	64.05	0	0	0	21.19	67.32
	138 ECS	66.25	0	0	0	6.61	82.82
	Average	65.15 ^{AB}	0	0	0	13.90 ^{AB}	75.07
H&N Nick Chick	69 ECS	64.98	0	0	0	13.33	75.00
	138 ECS	66.10	0	0	0	6.36	79.46
	Average	65.54 ^{AB}	0	0	0	9.84 ^{AB}	77.23
Novogen White	69 ECS	65.48	0	0	0	8.40	76.48
	138 ECS	65.93	0	0	0	0	76.75
	Average	65.70 ^A	0	0	0	4.20 ^B	76.61
All Strains	69 ECS	64.31	0	0	0.08	17.78	69.96
	138 ECS	64.88	0	0	0.19	10.68	70.98

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

TABLE 73. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 ECS	90.28	0	8.80	0.92	2.69	1.26 ^{abc}
	138 ECS	86.85	0	11.83	1.32	2.81	1.28 ^{abc}
	Average	88.56	0	10.31	1.12	2.75	1.27 ^{AB}
Shaver White	69 ECS	82.36	0.86	14.28	2.50	2.60	1.33 ^{abc}
	138 ECS	77.86	0	22.14	0	2.73	1.22 ^{abc}
	Average	80.11	0.43	18.21	1.25	2.66	1.27 ^{AB}
Dekalb White	69 ECS	91.73	0	7.27	1.00	2.78	1.24 ^{abc}
	138 ECS	88.51	0	11.49	0	2.92	1.23 ^{abc}
	Average	90.12	0	9.38	0.50	2.85	1.24 ^{AB}
Babcock White	69 ECS	87.22	0	11.11	1.67	2.559	1.44 ^a
	138 ECS	72.67	0	27.33	0	2.63	1.25 ^{abc}
	Average	79.95	0	19.22	0.83	2.61	1.34 ^A
ISA B-400	69 ECS	88.60	1.04	8.63	1.72	2.73	1.23 ^{abc}
	138 ECS	71.44	0	27.16	1.39	2.48	1.24 ^{abc}
	Average	80.02	0.52	17.90	1.56	2.61	1.23 ^{AB}
Hy-Line W-36	69 ECS	86.67	0	13.33	0	2.42	1.13 ^{bc}
	138 ECS	88.39	0	11.61	0	2.44	1.12 ^c
	Average	87.53	0	12.47	0	2.43	1.12 ^B
Hy-Line CV-24	69 ECS	89.68	0	9.45	0.86	2.60	1.36 ^{ab}
	138 ECS	80.82	0	18.05	1.14	2.71	1.22 ^{abc}
	Average	85.25	0	13.75	1.00	2.66	1.29 ^A
Lohmann LSL Lite	69 ECS	88.52	0	11.48	0	2.26	1.33 ^{abc}
	138 ECS	89.42	0	10.58	0	2.85	1.26 ^{abc}
	Average	88.97	0	11.03	0	2.55	1.30 ^A
H&N Nick Chick	69 ECS	88.33	0	9.79	1.88	2.69	1.35 ^{abc}
	138 ECS	85.82	0	14.18	0	2.81	1.26 ^{abc}
	Average	87.08	0	11.99	0.94	2.75	1.30 ^A
Novogen White	69 ECS	84.88	2.00	11.79	1.33	2.43	1.32 ^{abc}
	138 ECS	76.75	0	22.07	1.18	2.70	1.32 ^{abc}
	Average	80.81	1.00	16.93	1.25	2.57	1.32 ^A
All Strains	69 ECS	87.82 ^Y	0.39	10.59 ^Y	1.19	2.58	1.30 ^Y
	138 ECS	81.85 ^Z	0	17.64 ^Z	0.50	2.71	1.24 ^Z

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*density interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 74. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA Amber	69 ECS	11.86	0.382 ^b	20.61	73.60	45.41	0
	138 ECS	11.23	0.420 ^{ab}	21.61	77.18	47.35	0
	Average	11.55	0.401 ^B	21.11	75.39	46.38	0
TETRA Brown	69 ECS	11.19	0.420 ^{ab}	20.77	73.92	46.74	0.93
	138 ECS	11.51	0.420 ^{ab}	21.48	76.70	48.25	0
	Average	11.35	0.420 ^{AB}	21.12	75.31	47.49	0.46
Novogen Brown	69 ECS	11.53	0.436 ^{ab}	21.96	78.41	50.26	1.67
	138 ECS	10.97	0.454 ^{ab}	22.19	77.24	49.62	2.22
	Average	11.25	0.445 ^{AB}	22.07	77.83	49.94	1.95
Lohmann LB-Lite	69 ECS	11.70	0.442 ^{ab}	22.71	79.98	51.16	2.08
	138 ECS	10.92	0.505 ^a	24.32	85.87	55.16	1.39
	Average	11.31	0.474 ^{AB}	23.51	82.93	53.39	1.74
Hy-Line Silver Brown	69 ECS	11.75	0.425 ^{ab}	22.87	81.68	49.88	0
	138 ECS	12.07	0.425 ^{ab}	23.50	83.94	51.52	0
	Average	11.91	0.425 ^{AB}	23.19	82.81	50.69	0
Hy-Line Brown	69 ECS	11.54	0.414 ^{ab}	21.08	75.28	47.89	0
	138 ECS	11.26	0.412 ^{ab}	20.50	73.22	46.46	0
	Average	11.40	0.413 ^{AB}	20.79	74.25	47.17	0
ISA Brown	69 ECS	10.93	0.464 ^{ab}	24.09	81.46	50.72	0.56
	138 ECS	11.58	0.492 ^{ab}	22.91	85.54	56.78	1.11
	Average	11.26	0.078 ^A	23.50	83.50	53.75	0.83
Bovans Brown	69 ECS	11.86	0.438 ^{ab}	22.93	81.76	52.00	0.56
	138 ECS	11.28	0.480 ^{ab}	23.38	83.11	53.50	1.39
	Average	11.57	0.459 ^{AB}	23.15	82.43	52.75	0.97
All Strains	69 ECS	11.54	0.428	21.98	78.12	49.31	0.72
	138 ECS	11.35	0.451	22.63	80.50	51.08	0.76

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 75. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 ECS	61.70 ^{cd}	0	0	2.76	32.62 ^{ab}	55.08
	138 ECS	61.40 ^{cd}	0	0	0	42.12 ^a	48.07
	Average	61.55 ^{BC}	0	0	1.38	37.37 ^{AB}	51.58
TETRA Brown	69 ECS	63.21 ^{abcd}	0	0	0	20.68 ^{ab}	62.96
	138 ECS	62.92 ^{bcd}	0	0	0	31.84 ^{ab}	61.01
	Average	63.06 ^{ABC}	0	0	0	26.26 ^{AB}	61.99
Novogen Brown	69 ECS	65.05 ^{ab}	0	0	2.50	12.32 ^{ab}	74.63
	138 ECS	63.19 ^{bcd}	0	0	0.71	22.37 ^{ab}	65.19
	Average	64.12 ^A	0	0	1.61	17.35 ^B	69.91
Lohmann LB-Lite	69 ECS	64.54 ^{abc}	0	0	0	18.18 ^{ab}	69.71
	138 ECS	64.25 ^{abcd}	0	0	0	9.92 ^{ab}	68.71
	Average	64.39 ^A	0	0	0	14.04 ^B	69.09
Hy-Line Silver Brown	69 ECS	61.10 ^d	0	0	0	46.88 ^a	48.75
	138 ECS	61.34 ^{cd}	0	0	0	40.38 ^a	52.73
	Average	61.22 ^C	0	0	0	43.63 ^A	50.74
Hy-Line Brown	69 ECS	63.55 ^{abcd}	0	0	0	21.20 ^{ab}	67.49
	138 ECS	63.39 ^{abcd}	0	0	1.43	22.96 ^{ab}	63.89
	Average	63.47 ^{AB}	0	0	0.71	22.08 ^{AB}	65.69
ISA Brown	69 ECS	66.41 ^a	0	0	0	38.06 ^a	50.48
	138 ECS	62.28 ^{bcd}	0	0	0	3.33 ^b	74.28
	Average	64.34 ^A	0	0	0	20.70 ^B	62.38
Bovans Brown	69 ECS	63.60 ^{abcd}	0	0	0	24.12 ^{ab}	63.00
	138 ECS	64.26 ^{abcd}	0	0	0	26.50 ^{ab}	64.35
	Average	63.93 ^A	0	0	0	25.31	63.68
All Strains	69 ECS	63.13	0	0	0.43	26.76	61.48
	138 ECS	63.39	0	0	0.49	24.93	62.28

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

TABLE 76. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (483-511 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 ECS	90.45	0	8.59	0.96	2.32	1.33
	138 ECS	90.20	1.67	5.86	2.27	2.43	1.26
	Average	90.32	0.83	7.22	1.62	2.38	1.29
TETRA Brown	69 ECS	83.64	0	16.36	0	2.29	1.25
	138 ECS	92.86	0	4.76	2.38	2.45	1.29
	Average	88.25	0	10.56	1.19	2.37	1.27
Novogen Brown	69 ECS	87.67	1.33	11.00	0	2.47	1.29
	138 ECS	90.06	0	9.94	0	2.50	1.23
	Average	88.86	0.67	10.47	0	2.49	1.26
Lohmann LB-Lite	69 ECS	87.66	0	12.34	0	2.56	1.31
	138 ECS	78.62	0	21.38	0	2.61	1.22
	Average	83.14	0	16.86	0	2.58	1.27
Hy-Line Silver Brown	69 ECS	95.62	0	4.38	0	2.67	1.32
	138 ECS	93.11	0	6.89	0	2.70	1.35
	Average	94.37	0	5.63	0	2.69	1.34
Hy-Line Brown	69 ECS	88.70	0	11.30	0	2.38	1.29
	138 ECS	88.28	0	8.65	3.08	2.27	1.26
	Average	88.49	0	9.98	1.54	2.33	1.28
ISA Brown	69 ECS	88.54	0	10.79	0.66	2.57	1.22
	138 ECS	77.61	0	22.39	0	2.58	1.30
	Average	83.08	0	16.59	0.33	2.58	1.26
Bovans Brown	69 ECS	87.12	0	9.48	3.41	2.52	1.33
	138 ECS	90.85	0	6.65	2.50	2.64	1.26
	Average	88.98	0	8.06	2.95	2.58	1.30
All Strains	69 ECS	88.67	0.16	10.53	0.63	2.47	1.29
	138 ECS	87.70	0.21	10.82	1.28	2.52	1.27

¹All strains were housed such that each strain is equally represented in each density.
Enriched Colony Housing System=ECS.

TABLE 77. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans White	69 ECS	119.53 ^{abcd}	0.40 ^{efg}	188.90 ^{ab}	73.81 ^{abcd}	48.04 ^{abc}	6.95
	138 ECS	111.16 ^{cdef}	0.46 ^{abcde}	199.30 ^{ab}	77.76 ^{abc}	50.03 ^{abc}	6.95
	Average	115.34 ^{AB}	0.43 ^{AB}	194.10 ^A	75.79 ^A	49.03 ^A	6.95
Shaver White	69 ECS	121.11 ^{abc}	0.39 ^{fg}	183.66 ^{ab}	70.92 ^{abcde}	46.92 ^{bcd}	12.51
	138 ECS	108.14 ^{efg}	0.46 ^{abcd}	192.03 ^{ab}	75.17 ^{abcd}	49.86 ^{abc}	8.34
	Average	114.62 ^{AB}	0.43 ^{AB}	187.84 ^A	73.05 ^A	48.39 ^A	10.42
Dekalb White	69 ECS	113.95 ^{bcdef}	0.41 ^{efg}	185.22 ^{ab}	72.02 ^{abcde}	46.47 ^{bcd}	13.90
	138 ECS	105.80 ^{fg}	0.48 ^{ab}	202.78 ^a	79.03 ^{ab}	51.04 ^{abc}	9.73
	Average	109.87 ^B	0.45 ^A	194.00 ^A	75.53 ^A	48.78 ^A	11.81
Babcock White	69 ECS	126.07 ^a	0.39 ^{fg}	186.77 ^{ab}	72.73 ^{abcd}	48.19 ^{abc}	17.37
	138 ECS	105.80 ^{fg}	0.47 ^{abc}	193.28 ^{ab}	75.62 ^{abcd}	50.30 ^{abc}	6.67
	Average	115.94 ^{AB}	0.43 ^{AB}	190.02 ^A	74.17 ^A	49.24 ^A	12.02
ISA B-400	69 ECS	114.31 ^{bcdef}	0.42 ^{defg}	187.04 ^{ab}	73.09 ^{abcd}	47.91 ^{abc}	11.90
	138 ECS	109.13 ^{def}	0.43 ^{bcdef}	182.15 ^{ab}	71.22 ^{abcde}	47.48 ^{bcd}	12.50
	Average	111.72 ^{AB}	0.43 ^{AB}	184.59 ^{AB}	72.16 ^A	47.69 ^A	12.20
Hy-Line W-36	69 ECS	103.75 ^{fg}	0.39 ^{fg}	156.24 ^b	61.11 ^f	40.12 ^e	9.26
	138 ECS	96.27 ^g	0.42 ^{cdefg}	158.71 ^{ab}	62.20 ^{ef}	40.97 ^{de}	5.56
	Average	100.01 ^C	0.41 ^B	157.48 ^B	61.66 ^B	40.54 ^B	7.41
Hy-Line CV-24	69 ECS	118.21 ^{abcde}	0.39 ^{fg}	171.74 ^{ab}	68.02 ^{def}	46.11 ^{cde}	11.12
	138 ECS	106.53 ^{fg}	0.47 ^{abc}	191.24 ^{ab}	75.58 ^{abcd}	49.92 ^{abc}	12.51
	Average	112.37 ^{AB}	0.43 ^{AB}	181.49 ^{AB}	71.30 ^A	48.02 ^A	11.81
Lohmann LSL Lite	69 ECS	127.83 ^a	0.37 ^g	180.36 ^{ab}	70.17 ^{bcdef}	47.37 ^{bcd}	18.06
	138 ECS	108.79 ^{def}	0.45 ^{abcde}	185.99 ^{ab}	72.20 ^{abcde}	48.61 ^{abc}	12.51
	Average	118.31 ^A	0.41 ^B	183.18 ^{AB}	71.19 ^A	47.99 ^A	15.29
H&N Nick Chick	69 ECS	122.37 ^{ab}	0.38 ^g	176.32 ^{ab}	68.54 ^{cdef}	45.73 ^{cde}	13.20
	138 ECS	109.42 ^{def}	0.49 ^a	203.66 ^a	79.08 ^a	53.03 ^a	19.46
	Average	115.90 ^{AB}	0.43 ^{AB}	189.99 ^A	73.81 ^A	49.51 ^A	16.33
Novogen White	69 ECS	124.85 ^a	0.38 ^g	182.65 ^{ab}	70.24 ^{bcdef}	47.51 ^{bc}	16.12
	138 ECS	111.33 ^{cdef}	0.47 ^{abc}	196.51 ^{ab}	76.64 ^{abcd}	51.92 ^{ab}	11.12
	Average	118.09 ^A	0.43 ^{AB}	189.58 ^A	73.44 ^A	49.72 ^A	13.62
All Strains	69 ECS	119.20 ^Z	0.39 ^Y	179.89 ^Z	70.07 ^Z	46.44 ^Z	13.04
	138 ECS	107.24 ^Y	0.46 ^Z	190.56 ^Y	74.35 ^Y	49.34 ^Y	10.53

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 78. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 ECS	65.11 ^{efgh}	0	0.14	0.53 ^{ab}	9.92 ^{abc}	83.61 ^{abc}
	138 ECS	64.35 ^h	0	0	1.25 ^a	12.97 ^a	78.80 ^c
	Average	64.73 ^E	0	0.07	0.89 ^A	11.44 ^A	81.21 ^C
Shaver White	69 ECS	66.24 ^{abcdefg}	0	0	0 ^b	5.54 ^{abc}	85.73 ^{abc}
	138 ECS	66.45 ^{abcdef}	0	0	0.19 ^{ab}	4.30 ^{abc}	88.39 ^{abc}
	Average	66.34 ^{BCD}	0	0	0.10 ^B	4.92 ^{BC}	87.06 ^{ABC}
Dekalb White	69 ECS	64.68 ^{fgh}	0	0.14	1 ^b	12.28 ^{ab}	81.43 ^{bc}
	138 ECS	64.65 ^{gh}	0	0	0.19 ^{ab}	9.22 ^{abc}	85.47 ^{abc}
	Average	64.66 ^E	0	0.07	0.10 ^B	10.75 ^{AB}	83.45 ^{BC}
Babcock White	69 ECS	66.29 ^{abcdefg}	0	0	0.11 ^b	6.14 ^{abc}	87.25 ^{abc}
	138 ECS	66.53 ^{abcde}	0	0	0 ^b	3.09 ^c	93.22 ^a
	Average	66.41 ^{BCD}	0	0	0.06 ^B	4.61 ^C	90.24 ^{AB}
ISA B-400	69 ECS	65.56 ^{defgh}	0	0	0.13 ^b	6.63 ^{abc}	85.92 ^{abc}
	138 ECS	66.68 ^{abcde}	0	0	0 ^b	4.28 ^{abc}	88.37 ^{abc}
	Average	66.12 ^{CD}	0	0	0.06 ^B	5.46 ^{BC}	87.15 ^{ABC}
Hy-Line W-36	69 ECS	65.76 ^{cdefgh}	0	0	0.18 ^{ab}	9.30 ^{abc}	84.48 ^{abc}
	138 ECS	65.98 ^{bcddefgh}	0	0	0 ^b	5.55 ^{abc}	88.70 ^{abc}
	Average	65.87 ^{DE}	0	0	0.09 ^B	7.42 ^{ABC}	86.59 ^{ABC}
Hy-Line CV-24	69 ECS	67.86 ^a	0	0	0 ^b	5.89 ^{abc}	89.32 ^{abc}
	138 ECS	67.02 ^{abcd}	0	0	0.39 ^{ab}	3.86 ^{bc}	86.56 ^{abc}
	Average	67.44 ^{AB}	0	0	0.19 ^B	4.88 ^C	87.94 ^{ABC}
Lohmann LSL Lite	69 ECS	67.39 ^{abc}	0	0	0.17 ^b	6.78 ^{abc}	87.25 ^{abc}
	138 ECS	67.36 ^{abc}	0	0	0 ^b	3.53 ^{bc}	91.14 ^{ab}
	Average	67.38 ^{AB}	0	0	0.08 ^B	5.15 ^{BC}	89.19 ^{AB}
H&N Nick Chick	69 ECS	67.79 ^{abcde}	0	0	0.14 ^b	5.17 ^{abc}	87.19 ^{abc}
	138 ECS	67.46 ^{abc}	0	0	0.50 ^{ab}	4.83 ^{abc}	88.19 ^{abc}
	Average	67.12 ^{ABC}	0	0	0.32 ^{AB}	5.00 ^{BC}	87.69 ^{ABC}
Novogen White	69 ECS	67.68 ^{ab}	0	0	0 ^b	3.07 ^c	92.91 ^a
	138 ECS	67.85 ^a	0	0	0 ^b	3.70 ^{bc}	88.55 ^{abc}
	Average	67.77 ^A	0	0	0 ^B	3.38 ^C	90.70 ^A
All Strains	69 ECS	66.34	0	0	0.13	7.07 ^Y	86.51
	138 ECS	66.43	0	0	0.25	5.54 ^Z	87.74

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefg - Different letters denote significant differences (P<.01) in the strain*density interactions.

TABLE 79. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 ECS	89.82 ^{ab}	0.31	8.65 ^{ab}	1.22	22.15 ^a	11.11
	138 ECS	88.74 ^{ab}	0.68	9.44 ^{ab}	1.13	20.55 ^a	10.34
	Average	89.29 ^A	0.50	9.04 ^B	1.17	21.35 ^A	10.73 ^A
Shaver White	69 ECS	80.58 ^b	0.36	16.79 ^a	2.28	21.62 ^a	11.24
	138 ECS	85.74 ^{ab}	0	12.90 ^{ab}	1.36	21.01 ^a	10.08
	Average	83.16 ^{AB}	0.18	14.85 ^{AB}	1.82	21.32 ^A	10.66 ^A
Dekalb White	69 ECS	88.55 ^{ab}	0.41	9.50 ^{ab}	1.55	22.64 ^a	10.60
	138 ECS	87.23 ^{ab}	0.86	11.20 ^{ab}	0.70	21.30 ^a	9.85
	Average	87.89 ^{AB}	0.64	10.35 ^{AB}	1.13	21.97 ^A	10.22 ^{AB}
Babcock White	69 ECS	87.79 ^{ab}	0.42	9.58 ^{ab}	2.22	20.66 ^a	11.82
	138 ECS	89.09 ^{ab}	0.37	9.82 ^{ab}	0.72	22.08 ^a	9.87
	Average	88.44 ^{AB}	0.40	9.70 ^B	1.47	21.37 ^A	10.84 ^A
ISA B-400	69 ECS	83.12 ^{ab}	0.29	14.79 ^{ab}	1.81	20.96 ^a	10.60
	138 ECS	82.320 ^{ab}	0.19	16.83 ^a	0.67	17.97 ^{ab}	10.12
	Average	82.72 ^B	0.24	15.80 ^A	1.24	19.47 ^A	10.36 ^{AB}
Hy-Line W-36	69 ECS	88.13 ^{ab}	0.40	10.00 ^{ab}	1.47	19.02 ^a	9.67
	138 ECS	86.88 ^{ab}	0	11.60 ^{ab}	1.52	18.39 ^a	8.97
	Average	87.51 ^{AB}	0.20	10.80 ^{AB}	1.50	18.70 ^{AB}	9.32 ^B
Hy-Line CV-24	69 ECS	91.60 ^a	0.33	6.81 ^b	1.26	20.78 ^a	10.90
	138 ECS	84.54 ^{ab}	1.30	12.58 ^{ab}	1.58	20.49 ^a	9.92
	Average	88.07 ^{AB}	0.81	9.70 ^{AB}	1.42	20.63 ^A	10.41 ^{AB}
Lohmann LSL Lite	69 ECS	88.68 ^{ab}	0.20	9.60 ^{ab}	1.52	22.28 ^a	11.90
	138 ECS	88.77 ^{ab}	0.82	10.01 ^{ab}	0.40	21.28 ^a	10.13
	Average	88.73 ^{AB}	0.51	9.80 ^{AB}	0.96	21.78 ^A	11.01 ^A
H&N Nick Chick	69 ECS	87.05 ^{ab}	0.23	9.84 ^{ab}	2.88	21.98 ^a	11.41
	138 ECS	87.73 ^{ab}	0.25	10.48 ^{ab}	1.54	21.12 ^a	10.12
	Average	87.39 ^{AB}	0.24	10.16 ^{AB}	2.21	21.55 ^A	10.77 ^A
Novogen White	69 ECS	91.54 ^a	0.79	7.08 ^b	0.59	12.86 ^b	11.62
	138 ECS	86.58 ^{ab}	0.28	12.06 ^{ab}	1.08	18.15 ^a	10.36
	Average	89.06 ^A	0.53	9.57 ^B	0.83	15.50 ^B	10.99 ^A
All Strains	69 ECS	87.69	0.37	10.26	1.68	20.50	11.09 ^Z
	138 ECS	86.76	0.47	11.69	1.07	20.23	9.98 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 80. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA Amber	69 ECS	123.28 ^{ab}	0.30 ^e	146.45	56.97 ^{cd}	36.46 ^{bc}	15.98
	138 ECS	101.36 ^e	0.35 ^{de}	147.80	57.23 ^{bcd}	35.68 ^c	16.68
	Average	112.32 ^{AB}	0.32 ^C	147.13	57.10 ^C	36.07 ^E	16.33
TETRA Brown	69 ECS	124.54 ^{ab}	0.32 ^{de}	156.54	60.94 ^{abcd}	39.88 ^{abc}	14.82
	138 ECS	106.75 ^{de}	0.32 ^{de}	137.11	53.67 ^d	34.72 ^c	5.56
	Average	115.65 ^{AB}	0.32 ^C	146.83	57.31 ^{BC}	37.30 ^{DE}	10.19
Novogen Brown	69 ECS	122.43 ^{ab}	0.34 ^{de}	159.21	62.36 ^{abcd}	41.91 ^{abc}	15.56
	138 ECS	104.27 ^e	0.42 ^{abc}	171.75	66.41 ^{abc}	43.16 ^{ab}	17.20
	Average	113.35 ^{AB}	0.38 ^{AB}	165.48	64.39 ^{AB}	42.53 ^{ABC}	16.38
Lohmann LB-Lite	69 ECS	121.95 ^{abc}	0.35 ^{de}	164.92	64.12 ^{abcd}	42.61 ^{abc}	20.85
	138 ECS	103.35 ^e	0.45 ^a	180.95	70.78 ^a	46.58 ^a	5.56
	Average	112.65 ^{AB}	0.40 ^A	172.94	67.45 ^A	44.60 ^{AB}	13.20
Hy-Line Silver Brown	69 ECS	127.66 ^a	0.32 ^{de}	166.26	64.79 ^{abcd}	41.17 ^{abc}	14.59
	138 ECS	109.34 ^{de}	0.37 ^{bcd}	164.07	64.25 ^{abcd}	40.50 ^{abc}	6.95
	Average	118.50 ^A	0.34 ^{BC}	165.17	64.52 ^{AB}	40.84 ^{BCD}	10.77
Hy-Line Brown	69 ECS	120.13 ^{abc}	0.34 ^{de}	155.59	60.77 ^{abcd}	40.58 ^{abc}	13.34
	138 ECS	104.51 ^e	0.35 ^{de}	144.30	55.91 ^d	36.63 ^{bc}	5.56
	Average	112.32 ^{AB}	0.34 ^C	149.94	58.34 ^{BC}	38.61 ^{CDE}	9.45
ISA Brown	69 ECS	116.62 ^{bcd}	0.36 ^d	162.35	63.62 ^{abcd}	41.59 ^{abc}	9.95
	138 ECS	106.72 ^e	0.41 ^{abc}	172.13	66.48 ^{abc}	44.38 ^a	21.12
	Average	111.67 ^B	0.38 ^A	167.24	65.05 ^A	42.98 ^{AB}	15.53
Bovans Brown	69 ECS	123.45 ^{ab}	0.36 ^{cd}	174.14	67.98 ^{ab}	44.93 ^a	9.45
	138 ECS	111.26 ^{cde}	0.42 ^{ab}	178.14	69.51 ^a	47.05 ^a	15.28
	Average	117.35 ^{AB}	0.39 ^A	176.14	68.74 ^A	45.99 ^A	12.37
All Strains	69 ECS	122.51 ^Y	0.34 ^Y	160.68	62.69	41.14	14.32
	138 ECS	105.95 ^Z	0.39 ^Z	162.03	63.03	40.09	11.74

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABCDE - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcde - Different letters denote significant differences (P<.01) in the strain*density interactions.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 81. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 ECS	64.10 ^{efgh}	0	0	0.19 ^{ab}	14.57 ^{ab}	80.43 ^{ab}
	138 ECS	62.43 ^h	0	0	2.60 ^a	16.08 ^a	73.60 ^b
	Average	63.27 ^C	0	0	1.39	15.33 ^A	77.01 ^B
TETRA Brown	69 ECS	65.49 ^{bcdef}	0	0	0 ^b	11.59 ^{abc}	84.74 ^{ab}
	138 ECS	64.69 ^{defg}	0	0	1.00 ^b	8.18 ^{abc}	87.48 ^a
	Average	65.09 ^B	0	0	0.50	9.89 ^{ABC}	86.11 ^A
Novogen Brown	69 ECS	67.25 ^{ab}	0	0	0.15 ^{ab}	4.84 ^{bc}	84.51 ^{ab}
	138 ECS	64.96 ^{defg}	0	0	0.24 ^{ab}	10.33 ^{abc}	89.41 ^a
	Average	66.10 ^{AB}	0	0	0.20	7.59 ^{BC}	86.96 ^A
Lohmann LB-Lite	69 ECS	66.49 ^{abcd}	0	0	0 ^b	4.25 ^c	89.58 ^a
	138 ECS	65.87 ^{bcde}	0	0	0.25 ^{ab}	5.61 ^{abc}	87.56 ^a
	Average	66.18 ^{AB}	0	0	0.12	4.93 ^C	88.57 ^A
Hy-Line Silver Brown	69 ECS	63.63 ^{fgh}	0	0	0.11 ^{ab}	13.78 ^{abc}	81.32 ^{ab}
	138 ECS	63.15 ^{gh}	0	0	1.44 ^{ab}	14.14 ^{abc}	82.69 ^{ab}
	Average	63.39 ^C	0	0	0.77	13.96 ^{AB}	82.00 ^{AB}
Hy-Line Brown	69 ECS	66.86 ^{abc}	0	0	0.44 ^{ab}	4.29 ^c	90.04 ^a
	138 ECS	65.78 ^{bcde}	0	0	0.42 ^{ab}	13.37 ^{abc}	82.19 ^{ab}
	Average	66.32 ^{AB}	0	0	0.43	8.83 ^{ABC}	86.11 ^A
ISA Brown	69 ECS	65.48 ^{cde}	0	0	0.32 ^{ab}	9.17 ^{abc}	86.21 ^a
	138 ECS	66.72 ^{abcd}	0	0	1.07 ^{ab}	6.38 ^{abc}	88.80 ^a
	Average	66.10 ^{AB}	0	0	0.69	7.77 ^{BC}	87.51 ^A
Bovans Brown	69 ECS	66.23 ^{abcd}	0	0.08	0.07 ^b	7.87 ^{abc}	86.78 ^a
	138 ECS	67.89 ^a	0	0	0 ^b	3.50 ^c	92.69 ^a
	Average	67.06 ^A	0	0.04	0.03	5.68 ^C	89.74 ^A
All Strains	69 ECS	65.69 ^Y	0	0.01	0.16 ^Y	8.80	86.07
	138 ECS	65.18 ^Z	0	0	0.88 ^Z	9.70	84.94

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcdefgh - Different letters denote significant differences (P<.01) in the strain*density interactions.

TABLE 82. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (511-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 ECS	91.46 ^{ab}	0.31	6.61 ^{ab}	1.63	16.41	10.64
	138 ECS	85.27 ^b	0	14.26 ^a	0.51	17.70	8.76
	Average	88.34	0.15	10.44 ^{AB}	1.07	17.06	9.70
TETRA Brown	69 ECS	92.22 ^{ab}	0.64	6.35 ^{ab}	0.78	18.82	10.76
	138 ECS	92.60 ^{ab}	0	6.46 ^{ab}	0.93	18.07	9.21
	Average	92.41	0.32	6.40 ^{AB}	0.86	18.45	9.98
Novogen Brown	69 ECS	89.87 ^{ab}	1.00	8.04 ^{ab}	1.08	18.40	10.57
	138 ECS	90.52 ^{ab}	0.42	8.69 ^{ab}	0.37	17.61	9.00
	Average	90.20	0.71	8.37 ^{AB}	0.73	18.01	9.79
Lohmann LB-Lite	69 ECS	88.57 ^{ab}	0.10	10.12 ^{ab}	1.21	19.66	10.51
	138 ECS	85.89 ^{ab}	0.46	12.60 ^a	1.04	18.53	8.92
	Average	87.23	0.28	11.36 ^A	1.13	19.10	9.72
Hy-Line Silver Brown	69 ECS	89.94 ^{ab}	1.43	7.30 ^{ab}	1.34	18.21	11.03
	138 ECS	91.43 ^{ab}	0.25	7.85 ^{ab}	0.47	17.57	9.46
	Average	90.69	0.84	7.57 ^{AB}	0.90	17.89	10.24
Hy-Line Brown	69 ECS	90.86 ^{ab}	0.63	8.09 ^{ab}	0.42	15.87	10.36
	138 ECS	93.44 ^a	0.22	4.61 ^b	1.73	18.23	9.09
	Average	92.15	0.43	6.35 ^B	1.07	17.05	9.73
ISA Brown	69 ECS	91.78 ^{ab}	0.87	6.68 ^{ab}	0.68	19.17	10.03
	138 ECS	89.53 ^{ab}	0	9.76 ^{ab}	0.71	18.64	9.22
	Average	90.63	0.43	8.22 ^{AB}	0.70	18.90	9.63
Bovans Brown	69 ECS	89.75 ^{ab}	0.10	9.04 ^{ab}	1.11	19.50	10.66
	138 ECS	90.04 ^{ab}	0.96	9.00 ^{ab}	0	20.17	9.61
	Average	89.89	0.53	9.02 ^{AB}	0.56	19.83	10.14
All Strains	69 ECS	90.55	0.64	7.78	1.03	18.26	10.57 ^Z
	138 ECS	89.83	0.29	9.16	0.72	18.32	9.16 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 83. EFFECT OF WHITE EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
Bovans	69 ECS	109.98	0.44	520.46 ^{abc}	79.76	49.29	28.48
White	138 ECS	107.71	0.47	537.26 ^{abc}	82.66	51.55	19.45
	Average	108.85 ^A	0.45	528.95 ^{AB}	81.21 ^{AB}	50.42 ^A	23.96
Shaver	69 ECS	110.89	0.45	526.09 ^{abc}	79.83	50.10	46.53
White	138 ECS	104.88	0.49	536.87 ^{abc}	82.92	52.32	11.12
	Average	107.88 ^A	0.47	531.48 ^{AB}	81.38 ^{AB}	51.21 ^A	28.82
Dekalb	69 ECS	107.73	0.47	520.94 ^{abc}	80.18	49.75	24.31
White	138 ECS	104.77	0.49	551.30 ^a	84.91	52.58	16.67
	Average	106.25 ^A	0.48	536.12 ^{AB}	82.55 ^A	51.16 ^A	20.49
Babcock	69 ECS	116.74	0.44	537.44 ^{abc}	82.66	51.83	31.26
White	138 ECS	105.18	0.50	543.84 ^{ab}	84.01	53.64	10.00
	Average	110.96 ^A	0.47	540.64 ^{AB}	83.33 ^A	52.74 ^A	20.63
ISA	69 ECS	108.86	0.47	536.58 ^{abc}	82.32	52.05	30.08
B-400	138 ECS	107.61	0.48	527.07 ^{abc}	81.30	52.03	20.84
	Average	108.24 ^A	0.47	531.82 ^{AB}	81.81 ^{AB}	52.04 ^A	25.46
Hy-Line	69 ECS	100.06	0.44	476.10 ^c	73.50	45.06	13.89
W-36	138 ECS	95.66	0.47	479.45 ^{bc}	74.08	45.86	7.41
	Average	97.86 ^B	0.46	477.78 ^B	73.79 ^B	45.46 ^B	10.65
Hy-Line	69 ECS	111.20	0.42	489.92 ^{abc}	75.67	47.92	22.92
CV-24	138 ECS	105.09	0.48	527.18 ^{abc}	81.23	51.25	19.46
	Average	108.15 ^A	0.45	508.55 ^{AB}	78.45 ^{AB}	49.58 ^{AB}	21.19
Lohmann	69 ECS	114.72	0.43	511.22 ^{abc}	78.51	49.68	31.25
LSL Lite	138 ECS	106.68	0.48	531.01 ^{abc}	81.51	52.18	23.62
	Average	110.70 ^A	0.46	521.12 ^{AB}	80.01 ^{AB}	50.93 ^A	27.43
H&N	69 ECS	112.54	0.43	510.26 ^{abc}	78.23	49.47	28.87
Nick Chick	138 ECS	107.40	0.49	544.47 ^{ab}	83.76	53.83	27.79
	Average	109.97 ^A	0.46	527.36 ^{AB}	81.00 ^{AB}	51.65 ^A	28.83
Novogen	69 ECS	115.62	0.43	509.46 ^{abc}	77.66	49.55	38.33
White	138 ECS	107.78	0.49	540.55 ^{abc}	83.27	53.73	18.90
	Average	111.70 ^A	0.46	525.01 ^{AB}	80.46 ^{AB}	51.64 ^A	28.62
All Strains	69 ECS	110.83 ^Z	0.48 ^Z	513.86 ^Z	78.83 ^Z	49.47 ^Z	29.69 ^Z
	138 ECS	105.28 ^Y	0.44 ^Y	531.90 ^Y	81.96 ^Y	52.00 ^Y	17.52 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyses was transformed in Square Root Asin

TABLE 84. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
Bovans White	69 ECS	61.60	0	2.89	8.14	19.76	66.79
	138 ECS	61.76	3.40	1.20	4.53	22.06	65.90
	Average	61.68 ^B	1.70	2.05	6.34	20.91 ^A	66.35
Shaver White	69 ECS	62.51	0	2.63	6.78	16.96	69.75
	138 ECS	62.76	0.54	2.44	5.37	15.66	73.12
	Average	62.63 ^{AB}	0.27	2.54	6.08	16.31 ^{AB}	71.43
Dekalb White	69 ECS	61.49	0	3.99	6.00	16.93	67.46
	138 ECS	61.65	1.09	1.16	7.26	19.22	68.98
	Average	61.57 ^B	0.54	2.58	6.63	19.58 ^{AB}	68.22
Babcock White	69 ECS	62.45	0.29	2.92	6.75	16.96	70.15
	138 ECS	63.48	0.96	2.14	3.53	13.38	78.32
	Average	62.96 ^{AB}	0.62	2.53	5.14	15.17 ^{AB}	74.24
ISA B-400	69 ECS	62.85	0	2.47	5.28	16.17	72.71
	138 ECS	63.62	0.81	1.64	4.60	13.14	76.73
	Average	63.23 ^{AB}	0.41	2.05	4.94	14.66 ^{AB}	74.72
Hy-Line W-36	69 ECS	61.26	0	4.70	9.74	19.19	63.71
	138 ECS	61.98	1.45	1.04	8.64	18.19	68.23
	Average	61.62 ^B	0.72	2.87	9.19	18.69 ^{AB}	65.97
Hy-Line CV-24	69 ECS	63.15	1.10	3.59	6.31	16.24	70.50
	138 ECS	62.68	0.81	0.93	8.72	15.80	70.01
	Average	62.92 ^{AB}	0.96	2.26	7.51	16.02 ^{AB}	70.26
Lohmann LSL Lite	69 ECS	62.88	0	2.87	6.18	17.63	70.81
	138 ECS	63.48	1.27	1.97	5.38	13.13	76.05
	Average	63.18 ^{AB}	0.64	2.41	5.78	15.38 ^{AB}	73.43
H&N Nick Chick	69 ECS	62.95	0	3.21	6.27	15.67	71.76
	138 ECS	63.58	2.17	1.31	6.16	13.09	74.79
	Average	63.27 ^{AB}	1.09	2.17	6.21	14.38 ^{AB}	73.28
Novogen White	69 ECS	63.54	0	2.99	6.09	14.71	74.34
	138 ECS	63.94	1.74	2.50	3.47	12.30	76.60
	Average	63.74 ^A	0.87	2.74	4.78	13.50 ^B	75.47
All Strains	69 ECS	62.47	0.14 ^Y	3.21 ^Z	6.76	17.32	69.80
	138 ECS	62.89	1.43 ^Z	1.63 ^Y	5.76	15.60	72.87

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 85. EFFECT OF WHITE EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
Bovans White	69 ECS	92.92	0.41	5.89	0.77	58.86 ^{abc}	28.00
	138 ECS	90.03	0.48	8.76	0.76	60.27 ^{abc}	27.75
	Average	91.48 ^A	0.44	7.32 ^B	0.77	59.56 ^A	27.88 ^A
Shaver White	69 ECS	87.31	0.41	10.73	1.50	58.20 ^{abc}	28.28
	138 ECS	85.72	0.32	13.18	0.79	59.47 ^{abc}	27.06
	Average	86.52 ^C	0.36	11.96 ^A	1.15	58.83 ^A	27.67 ^A
Dekalb White	69 ECS	90.91	0.46	7.66	0.99	58.86 ^{abc}	27.52
	138 ECS	90.67	0.34	8.28	0.73	62.01 ^a	27.04
	Average	90.79 ^A	0.40	7.97 ^B	0.86	60.43 ^A	27.28 ^{AB}
Babcock White	69 ECS	90.99	0.62	7.20	1.24	60.28 ^{abc}	29.82
	138 ECS	88.91	0.37	10.23	0.55	61.41 ^{ab}	27.18
	Average	89.95 ^{ABC}	0.50	8.72 ^{AB}	0.89	60.85 ^A	28.50 ^A
ISA B-400	69 ECS	88.74	0.40	9.82	1.10	60.13 ^{abc}	27.80
	138 ECS	84.80	0.24	14.25	0.78	58.36 ^{abc}	27.75
	Average	86.77 ^{BC}	0.32	12.03 ^A	0.94	59.25 ^A	27.78 ^A
Hy-Line W-36	69 ECS	90.98	0.36	7.65	1.03	52.88 ^C	25.60
	138 ECS	90.03	0	8.97	1.01	53.48 ^{bc}	24.75
	Average	90.51 ^{ABC}	0.18	8.31 ^{AB}	1.02	53.18 ^B	25.17 ^B
Hy-Line CV-24	69 ECS	92.33	0.33	6.50	0.87	55.37 ^{abc}	28.40
	138 ECS	88.74	0.59	9.94	0.75	58.75 ^{abc}	27.11
	Average	90.53 ^{AB}	0.46	8.22 ^B	0.81	57.06 ^{AB}	27.76 ^A
Lohmann LSL Lite	69 ECS	90.51	0.72	7.88	0.89	57.83 ^{abc}	29.19
	138 ECS	91.12	0.43	8.20	0.29	60.26 ^{abc}	27.51
	Average	90.82 ^A	0.58	8.04 ^B	0.59	59.04 ^A	28.35 ^A
H&N Nick Chick	69 ECS	90.76	0.46	7.10	1.66	57.38 ^{abc}	28.70
	138 ECS	90.43	0.38	8.55	0.66	61.46 ^{ab}	27.63
	Average	90.60 ^A	0.42	7.82 ^B	1.16	59.42 ^A	28.17 ^A
Novogen White	69 ECS	92.05	0.68	6.70	0.56	58.12 ^{abc}	29.30
	138 ECS	87.34	0.48	11.31	0.91	60.27 ^{abc}	27.76
	Average	89.70 ^{ABC}	0.58	9.01 ^{AB}	0.74	59.20 ^A	28.54 ^A
All Strains	69 ECS	90.75 ^Y	0.48	7.71 ^Y	1.06 ^Z	57.79 ^Z	28.26 ^Z
	138 ECS	88.78 ^Z	0.36	10.17 ^Z	0.72 ^Y	59.57 ^Y	27.16 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abc - Different letters denote significant differences (P<.01) in the strain*density interactions

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 86. EFFECT OF BROWN EGG STRAIN AND DENSITY ON PERFORMANCE OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Feed Consumption	Feed Conversion	Eggs Per Bird Housed	Egg Production	Egg Mass	Mortality
(Strain)	(in ² /hen)	(kg/100/hen/d)	(g egg/g feed)		(HD%)	(g/HD)	(%)
TETRA Amber	69 ECS	112.52	0.37 ^d	450.08	69.35	41.38	17.37
	138 ECS	105.28	0.39 ^{bcd}	459.43	70.77	42.42	23.62
	Average	108.90 ^{AB}	0.38 ^C	454.76	70.06	41.90 ^B	20.49
TETRA Brown	69 ECS	111.46	0.39 ^{bcd}	459.49	70.74	43.55	21.30
	138 ECS	107.18	0.38 ^{cd}	436.90	67.47	41.39	7.41
	Average	109.32 ^{AB}	0.38 ^{BC}	448.19	69.10	42.47 ^B	14.36
Novogen Brown	69 ECS	111.42	0.42 ^{bcd}	479.02	73.85	46.67	20.56
	138 ECS	107.09	0.42 ^{bcd}	470.82	72.30	45.56	28.31
	Average	109.25 ^{AB}	0.42 ^{AB}	474.92	73.08	46.11 ^{AB}	24.44
Lohmann LB-Lite	69 ECS	110.52	0.42 ^{bcd}	481.80	74.20	46.32	26.40
	138 ECS	102.55	0.48 ^a	515.76	79.66	49.95	6.95
	Average	106.53 ^B	0.45 ^A	498.78	76.93	48.13 ^A	16.67
Hy-Line Silver Brown	69 ECS	115.38	0.39 ^{bcd}	492.55	75.81	45.14	21.53
	138 ECS	108.03	0.41 ^{bcd}	491.70	75.95	45.43	11.12
	Average	111.70 ^{AB}	0.40 ^{BC}	492.13	75.88	45.28 ^{AB}	16.32
Hy-Line Brown	69 ECS	111.40	0.40 ^{bcd}	465.90	71.90	44.89	17.79
	138 ECS	107.22	0.40 ^{bcd}	449.13	69.48	43.58	5.56
	Average	109.31 ^{AB}	0.40 ^{BC}	457.52	70.69	44.23 ^{AB}	11.67
ISA Brown	69 ECS	108.12	0.43 ^{abc}	494.17	76.21	47.17	14.94
	138 ECS	105.93	0.45 ^{ab}	492.61	75.68	48.50	24.45
	Average	107.02 ^A	0.44 ^A	493.39	75.95	47.84 ^A	19.70
Bovans Brown	69 ECS	114.59	0.41 ^{bcd}	496.57	76.57	48.06	15.56
	138 ECS	111.23	0.43 ^{abc}	487.56	75.18	48.57	20.84
	Average	112.91 ^A	0.42 ^{AB}	492.06	75.88	48.32 ^A	18.20
All Strains	69 ECS	111.93 ^Z	0.40 ^Y	477.45	73.58	45.40	19.43
	138 ECS	106.81 ^Y	0.42 ^Z	475.49	73.31	45.67	16.03

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

Mortality percentage prior to analyzes was transformed in Square Root Asin

TABLE 87. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG WEIGHT AND EGG SIZE DISTRIBUTION OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Egg Weight	Pee Wee	Small	Medium	Large	Extra Large
(Strain)	(in ² /hen)	(g/egg)	(%)	(%)	(%)	(%)	(%)
TETRA Amber	69 ECS	59.70 ^d	0.27	4.46	12.77	23.33	57.09 ^{abc}
	138 ECS	59.66 ^d	1.09	2.75	10.04	25.04	57.96 ^{ab}
	Average	59.68 ^C	0.68	3.60	11.41 ^{AB}	24.18 ^{AB}	57.52 ^B
TETRA Brown	69 ECS	61.47 ^{abcd}	0	2.48	8.93	22.40	64.72 ^{abc}
	138 ECS	61.45 ^{abcd}	1.93	2.00	5.58	21.20	67.80 ^{abc}
	Average	61.46 ^{BC}	0.96	2.24	7.25 ^{ABC}	21.80 ^{ABC}	66.26 ^{AB}
Novogen Brown	69 ECS	63.17 ^{ab}	0	1.83	7.72	17.59	70.44 ^{abc}
	138 ECS	62.62 ^{ab}	0.38	1.50	4.19	17.00	75.91 ^{abc}
	Average	62.90 ^{AB}	0.19	1.67	5.96 ^C	17.30 ^C	72.68 ^A
Lohmann LB-Lite	69 ECS	62.20 ^{abcd}	0	2.80	7.53	16.71	70.25 ^{abc}
	138 ECS	62.49 ^{abc}	0.32	2.34	5.17	17.76	71.75 ^{abc}
	Average	62.34 ^{AB}	0.16	2.57	6.35 ^{BC}	17.23 ^C	71.00 ^A
Hy-Line Silver Brown	69 ECS	59.68 ^d	0	2.61	14.05	25.39	56.10 ^a
	138 ECS	59.75 ^{cd}	0.61	2.81	10.73	26.70	58.13 ^a
	Average	59.72 ^C	0.30	2.71	12.39 ^A	26.04 ^A	57.11 ^B
Hy-Line Brown	69 ECS	62.56 ^{ab}	0	0.91	7.09	21.08	68.62 ^{abc}
	138 ECS	62.73 ^{ab}	0.62	1.89	4.41	22.17	69.20 ^{abc}
	Average	62.64 ^{AB}	0.31	1.40	5.75 ^C	21.63 ^{ABC}	68.91 ^A
ISA Brown	69 ECS	61.69 ^{abcd}	0.28	1.57	8.20	21.76	66.22 ^{abc}
	138 ECS	63.85 ^{ab}	0	2.62	4.47	13.85	77.48 ^c
	Average	62.77 ^{AB}	0.14	2.09	6.33 ^{BC}	17.80 ^{BC}	71.85 ^A
Bovans Brown	69 ECS	62.84 ^{ab}	0	2.07	7.84	19.52	68.23 ^{abc}
	138 ECS	64.28 ^a	1.09	1.22	4.60	14.36	77.16 ^{bc}
	Average	63.56 ^A	0.54	1.64	6.22 ^{BC}	16.94 ^C	72.70 ^A
All Strains	69 ECS	61.66	0.07 ^Y	2.34	9.27 ^Y	19.76	65.21 ^Z
	138 ECS	62.11	0.75 ^Z	2.14	6.15 ^Z	20.97	69.30 ^Y

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

ABC - Different letters denote significant differences (P<.01), comparisons made among strain average values.

abcd - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 88. EFFECT OF BROWN EGG STRAIN AND DENSITY ON EGG QUALITY, INCOME AND FEED COSTS OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEMS

Breeder	Density ¹	Grade A	Grade B	Cracks	Loss	Egg Income	Feed Costs
(Strain)	(in ² /hen)	(%)	(%)	(%)	(%)	(\$/hen)	(\$/hen)
TETRA Amber	69 ECS	91.84 ^{ab}	0.35	6.92 ^{ab}	0.88	49.60	27.55
	138 ECS	88.71 ^{ab}	0.87	10.04 ^{ab}	0.44	50.72	26.30
	Average	90.27 ^{AB}	0.61	8.48 ^{AB}	0.66	50.16	26.92 ^{AB}
TETRA Brown	69 ECS	91.96 ^{ab}	0.59	6.71 ^{ab}	0.71	51.79	27.30
	138 ECS	93.04 ^a	0.10	6.35 ^{ab}	0.56	49.50	26.69
	Average	92.50 ^{AB}	0.35	6.52 ^{AB}	0.64	50.64	26.99 ^{AB}
Novogen Brown	69 ECS	91.17 ^{ab}	0.79	7.23 ^{ab}	0.82	54.05	27.32
	138 ECS	91.63 ^{ab}	0.47	7.55 ^{ab}	0.44	53.59	26.73
	Average	91.40 ^{AB}	0.63	7.39 ^{AB}	0.63	53.82	27.02 ^{AB}
Lohmann LB-Lite	69 ECS	90.89 ^{ab}	0.25	7.68 ^{ab}	1.16	54.24	27.06
	138 ECS	88.05 ^b	0.48	10.97 ^a	0.55	57.63	25.52
	Average	89.47 ^B	0.36	9.33 ^A	0.86	55.94	26.29 ^B
Hy-Line Silver Brown	69 ECS	93.17 ^a	0.62	5.70 ^b	0.54	55.09	28.22
	138 ECS	93.01 ^a	0.29	6.14 ^b	0.60	55.14	26.87
	Average	93.09 ^A	0.46	5.92 ^B	0.57	55.12	27.54 ^{AB}
Hy-Line Brown	69 ECS	92.36 ^{ab}	0.37	6.80 ^{ab}	0.46	52.99	27.32
	138 ECS	92.27 ^{ab}	0.36	6.26 ^b	1.12	50.90	26.73
	Average	92.31 ^{AB}	0.36	6.53 ^{AB}	0.79	51.95	27.03 ^{AB}
ISA Brown	69 ECS	92.78 ^a	0.47	6.22 ^b	0.52	56.24	26.52
	138 ECS	89.78 ^{ab}	0.06	9.80 ^{ab}	0.39	55.85	26.37
	Average	91.28 ^{AB}	0.27	8.01 ^{AB}	0.45	56.04	26.44 ^{AB}
Bovans Brown	69 ECS	92.28 ^{ab}	0.12	6.68 ^{ab}	0.96	56.19	28.11
	138 ECS	91.03 ^{ab}	0.52	8.29 ^{ab}	0.20	55.69	27.68
	Average	91.66 ^{AB}	0.32	7.48 ^{AB}	0.58	55.94	27.89 ^A
All Strains	69 ECS	92.06	0.45	6.74 ^Y	0.76	53.77	27.42 ^Z
	138 ECS	90.94	0.40	8.17 ^Z	0.54	53.63	26.61 ^Y

¹All strains were housed such that each strain is equally represented in each density.
Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values.

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

TABLE 89. EFFECT OF WHITE EGG STRAIN AND DENSITY ON BODY WEIGHT OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEM THAT WERE NON-MOLTED

Breeder	Density ¹	17 Wk* Body Wt	69 Wk** Body Wt	1st Cycle Wt Gain	73 Wk** Body Wt	109 Wk Body Wt	2 nd Cycle Wt Gain	Total Wt Gain
(Strain)	(in ² /hen)	(kg)	(kg)	(%)	(kg)	(kg)	(kg)	(%)
Bovans White	69 ECS	1.22	1.64 ^{ab}	34.4	1.72	1.84	0.10	50.60
	138 ECS	1.32	1.82 ^{ab}	37.9	1.86	1.80	-0.02	38.08
	Average	1.27	1.73	36.2	1.79	1.82 ^{AB}	0.04	44.34
Shaver White	69 ECS	1.28	1.73 ^{ab}	34.4	1.74	1.80	0.08	44.46
	138 ECS	1.34	1.92 ^{ab}	43.3	1.96	1.90	-0.04	46.11
	Average	1.32	1.82	38.6	1.85	1.85 ^{AB}	0.02	45.28
Dekalb White	69 ECS	1.20	1.61 ^{ab}	45.0	1.72	1.75	0	48.06
	138 ECS	1.20	1.74 ^{ab}	35.0	1.72	1.77	0.05	43.60
	Average	1.20	1.68	40.0	1.72	1.76 ^B	0.03	45.83
Babcock White	69 ECS	1.31	1.74 ^{ab}	48.1	1.73	1.95	0.16	50.72
	138 ECS	1.36	1.99 ^a	30.9	2.02	1.99	0.08	50.90
	Average	1.34	1.86	39.6	1.88	1.97 ^A	0.12	50.81
ISA B-400	69 ECS	1.18	1.63 ^{ab}	39.0	2.09	1.61	-0.36	34.66
	138 ECS	1.23	1.72 ^{ab}	39.8	1.78	1.84	0.10	50.82
	Average	1.20	1.68	39.2	1.94	1.72 ^B	-0.13	42.74
Hy-Line W-36	69 ECS	1.20	1.60 ^{ab}	33.3	1.57	1.81	0.18	52.41
	138 ECS	1.22	1.70 ^{ab}	39.3	1.70	1.86	0.20	54.47
	Average	1.21	1.65	36.4	1.64	1.83 ^{AB}	0.19	53.44
Hy-Line CV-26	69 ECS	1.12	1.74 ^{ab}	54.5	1.87	1.90	0.03	69.31
	138 ECS	1.19	1.74 ^{ab}	46.2	1.79	1.81	0.02	51.90
	Average	1.16	1.74	50.0	1.83	1.86 ^{AB}	0.02	60.60
Hy-Line CV-24	69 ECS	1.22	1.66 ^{ab}	36.1	1.69	1.84	0.17	47.75
	138 ECS	1.24	1.82 ^{ab}	46.8	1.75	1.82	0.10	44.71
	Average	1.23	1.74	41.5	1.72	1.83 ^{AB}	0.13	46.23
Lohmann LSL Lite	69 ECS	1.22	1.58 ^b	29.5	1.97	1.84	-0.15	48.58
	138 ECS	1.24	1.65 ^{ab}	33.1	1.69	1.80	0.08	44.68
	Average	1.23	1.62	30.9	1.83	1.82 ^{AB}	-0.03	46.63
H&N Nick Chick	69 ECS	1.24	1.66 ^{ab}	33.9	1.67	1.76	0.07	41.57
	138 ECS	1.24	1.83 ^{ab}	46.8	1.80	1.82	0.08	44.80
	Average	1.24	1.74	40.3	1.74	1.79 ^{AB}	0.08	43.18
Novogen White	69 ECS	1.22	1.64 ^{ab}	34.4	1.64	1.81	0.14	46.90
	138 ECS	1.26	1.83 ^{ab}	46.0	1.87	1.86	-0.06	45.59
	Average	1.24	1.74	40.3	1.76	1.84 ^{AB}	0.04	46.24
All Strains	69 ECS	1.22	1.66 ^Y	36.1 ^Y	1.76	1.81	0.04	48.64
	138 ECS	1.26	1.80 ^Z	42.9 ^Z	1.81	1.84	0.05	46.88

¹All strains were housed such that each strain is equally represented in each density.

Enriched Colony Housing System=ECS.

AB - Different letters denote significant differences (P<.01), comparisons made among strain average values

ab - Different letters denote significant differences (P<.01) in the strain*density interactions.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.

(*) All replicates in all strains were weight at 17 wks,

(**) Only a sample of replicates (2 per strain treatment) in each strain were weighted at 69 and 73 wks.

TABLE 90. EFFECT OF BROWN EGG STRAIN AND DENSITY ON BODY WEIGHT OF HENS IN THE 39th NCLP&MT (119-763 DAYS) IN THE ENRICHED COLONY HOUSING SYSTEM THAT WERE NON-MOLTED

Breeder	Density ¹	17 Wk* Body Wt	69 Wk** Body Wt	1st Cycle Wt Gain	73 Wk** Body Wt	109 Wk Body Wt	2 nd Cycle Wt Gain	Total Wt Gain
(Strain)	(in ² /hen)	(kg)	(kg)	(%)	(kg)	(kg)	(kg)	(%)
TETRA Amber	69 ECS	1.50	1.89	26.7	2.02	1.84	0.10	20.75
	138 ECS	1.55	2.04	31.0	2.15	1.77	0	13.08
	Average	1.52	1.96	28.9	2.07	1.80	0.05	17.92
TETRA Brown	69 ECS	1.54	1.88	22.1	1.87	1.92	0.18	26.18
	138 ECS	1.58	1.95	22.8	2.02	2.07	0.04	30.66
	Average	1.56	1.92	22.4	1.94	2.00	0.11	28.42
Novogen Brown	69 ECS	1.55	1.94	24.5	1.91	1.82	0.04	16.02
	138 ECS	1.58	1.94	22.8	2.02	1.93	-0.05	21.45
	Average	1.56	1.94	23.7	1.97	1.88	0	18.73
Lohmann LB-Lite	69 ECS	1.52	2.00	31.6	1.90	1.93	0.08	26.16
	138 ECS	1.57	1.92	22.9	2.04	2.02	-0.06	29.12
	Average	1.54	1.96	27.3	1.97	1.98	0.01	27.64
Hy-Line Silver Brown	69 ECS	1.42	1.98	39.4	1.98	1.80	0.16	21.66
	138 ECS	1.70	2.16	27.1	2.18	2.25	-0.06	27.25
	Average	1.56	2.07	32.7	2.08	1.93	0.05	24.45
Hy-Line Brown	69 ECS	1.53	1.87	22.2	2.38	1.96	-0.26	29.75
	138 ECS	1.64	2.08	28.0	2.09	1.84	0	13.80
	Average	1.58	1.98	25.3	2.24	1.90	-0.13	21.78
ISA Brown	69 ECS	1.45	1.82	25.5	1.87	1.87	0.08	30.73
	138 ECS	1.48	2.06	39.2	2.06	1.88	-0.04	26.04
	Average	1.46	1.94	32.9	1.96	1.87	0.02	28.38
Bovans Brown	69 ECS	1.60	1.94	21.3	1.94	1.89	0.05	22.78
	138 ECS	1.60	2.04	27.5	2.06	1.90	-0.04	19.16
	Average	1.60	1.99	24.4	2.00	1.89	0	20.97
All Strains	69 ECS	1.51	1.92 ^Z	26.5	1.98	1.88	0.05	24.25
	138 ECS	1.58	2.02 ^Y	27.8	2.07	1.93	-0.02	22.57

Enriched Colony Housing System=ECS.

YZ - Different letters denote significant differences (P<.01), comparisons made among density average values.


(*) All replicates in all strains were weighed at 17 wks,

(**) Only a sample of replicates (2 per strain treatment) in each strain were weighed at 69 and 73 wks.

Production Graphs for Laying Hens


- Enriched Colony Housing System
 - 445.2 cm²/hen (69 in²)
 - 890.3 cm²/hen (138 in²)
- Enrichable Cages at
 - 445.2 cm²/hen (69 in²)

Figure 1. Bovans White, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 2. Shaver, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 3. Dekalb, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 4. Babcock, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 5. B-400, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 6. Hy-Line W-36, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 7. Hy-Line CV-24, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 8. Lohmann, LSL-Lite, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹
by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS)
and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 9. Lohmann, “Nick Chick”, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 10. Novogen White, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 11. TETRA Amber, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 12. TETRA Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 13. Novogen Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 14. Lohmann, LB-Lite, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹
by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS)
and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 15. Hy-Line Silver Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 16. Hy-Line Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Figure 17. ISA Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²



¹ kg per 100 Hens

Figure 18. Bovans Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 138 in²) in Enriched Colony Housing System(ECS) and the Enrichable Cage (EC) at 69 in²




¹ kg per 100 Hens

Production Graphs for Laying Hens


- Conventional Cages
 - 445.2 cm²/hen (69 in²)
 - 774.2 cm²/hen (120 in²)

Figure 19. Bovans White, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 20. Shaver, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 21. Dekalb, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 22. Babcock, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 23. B-400, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 24. Hy-Line W-36, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 25. Hy-Line CV-26, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 26. Hy-Line CV-22, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by
hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 27. LSL-Lite, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 28. H&N “Nick Chick”, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹
by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 29. Novogen White, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 30. Tetra Amber, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 31. Tetra Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 32. Novogen Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 33. Lohman LB-Lite, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 34. Hy-Line Silver Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 35. Hy-Line Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage




¹ kg per 100 Hens

Figure 36. ISA Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage



¹ kg per 100 Hens

Figure 37. Bovans Brown, Bi-weekly Hen-day Egg Production and Period Feed Consumption¹ by hen density (69 and 120 in²) in Conventional Cage



¹ kg per 100 Hens

Table 91. Entries in the 39th NCLP&MT by Breeder, Stock Suppliers, and Categories			
Breeder	Stock	Category ¹	Source
Hy-Line International 2583 240 th Street Dallas Center, IA 50063	W-36	I-A	Hy-Line North America 4432 Highway 213, Box 309 Mansfield, GA 30255
	Hy-Line Brown	I-A	(Same)
	Hy-Line Silver Brown	III-A	(Same)
	CV22	II-A	(Same)
	CV24	II-A	(Same)
	CV26	II-A	(Same)
Lohmann Tierzucht GmbH Am Seedeich 9-11 . P.O.Box 460 D-27454 Cuxhaven, Germany	Lohmann LSL-Lite	I-A	Hy-Line North America Elizabeth- town 79 Industrial Rd Elizabethtown, PA 17022
	Lohmann LB-Lite	I-A	(Same)
H&N International 321 Burnett Ave South, Suite 300 Renton, Washington 98055	H&N “Nick Chick”	I-A	Feather Land Farms 32832 E. Peral Road Coberg, OR 97408
Instiut de Selection Animale (A Hendrix Genetic Company) ISA North America 650 Riverbend Drive, Suite C Kitchener, Ontario N2K 3S2 Canada	Bovans White	I-A	CPI-South Central Hatchery 5087 County Road 35 Bremen, AL 35033
	Dekalb White	I-A	(Same)
	Bovans Brown	I-A	(Same)
	Babcock White	II-A	Institute de Sélection Animale 650 Riverbend Dr. Suite C Kitchener, Ontario N2K 3S2 Canada
	B 400	II-A	(Same)
Tetra Americana, LLC 1105 Washington Road Lexington, GA 30648	Shaver White	I-A	Midwest Farms, LLC. 135 S. Epes St. Blackstone, VA 23824
	ISA Brown	I-A	(Same)
	TETRA Brown	I-A	CPI-MidAmerica Hatchery Lexington, GA 30648
NOVOGEN S.A.S. Mauguérand – Le Foeil BP 265 22 800 QUINTIN - FRANCE	TETRA Amber	I-A	(Same)
	NOVOgen BROWN	I-A	Morris Hatchery 18370 SW 232 Street, Goulds, FL 33170-5399
	NOVOgen WHITE	I-A	Pennovo Hatchery 621 Stevens Road Ephrata, PA 17522

¹ I = Extensive distribution in southeast United States

A = Entry requested

II = Little or no distribution in southeast United States

III = Unavailable for commercial distribution in United States